The second message of the study is that the term persistent vegetative state is widely misused, even in recent papers.³ No clinician should accept a diagnosis of such a state within 12 months of the onset of coma. I strongly recommend use of the term "continuing vegetative state," or even the simpler term "coma," for all patients who are comatose over the first 12 months after the onset of coma. In this way clinicians and relatives will avoid drawing premature or incorrect conclusions from loosely applied diagnostic labels.

DERICK T WADE

Consultant in neurological disability

Rivermead Rehabilitation Centre,
Oxford OX1 4XD

- 1 Andrews K, Murphy L, Munday R, Littlewood C. Misdiagnosis of the vegetative state: retrospective study in a rehabilitation unit. *BMJ* 1996;313:13-6. (6 July.) [With commentary by R Smith.]
- 2 The permanent vegetative state. *J R Coll Physicians Lond* 1996; 30:119-21.
- 3 Grubb A, Walsh P, Lambe N, Murrells T, Robinson S. Survey of British clinicians' views on management of patients in persistent vegetative state. *Lancet* 1996;348:35-40.

Misdiagnosis certainly occurs

EDITOR,—It is gratifying, though alarming, that Keith Andrews and colleagues' study¹ so closely duplicates our findings regarding misdiagnosis of the vegetative state.² The accompanying editorial by Ronald Cranford, which suggests that the rate of misdiagnosis is a new finding and that Andrews and colleagues' methods were questionable, merits critical comment.³

Aside from the implication that physicians who provide rehabilitative care are scientifically suspect, Cranford shows his inexperience in the neurorehabilitation of brain injured patients when he doubts the veracity of the buzzer communication system.³ Prosthetic communication and environmental control systems, including simple on-off switching devices like the buzzer, are commonly used in rehabilitation.⁴ Obviously, the patients in Andrews and colleagues' study were showing the responsiveness required to negate a diagnosis of vegetative state before they could use a communication device with 90% consistency.

Cranford—a member of the Multi-Society Task Force on Persistent Vegetative State—says that neurodiagnostic tests have "some use in the diagnosis" of vegetative state; we find this perplexing. While the task force stated that structural and functional neuroimaging "may provide useful information when used in conjunction with clinical evaluation," it also said that "neurodiagnostic tests alone can neither confirm the diagnosis of vegetative state nor predict the potential for recovery of awareness."⁵ Contradictory statements such as these need to be reconciled. In addition, if these tests are diagnostic, as Cranford suggests, then it was the referring physicians who were remiss not Andrews and colleagues. Whether neurodiagnostic tests were used has little relevance to Andrews and colleagues since their purpose was to validate or invalidate the preadmission diagnoses.

Cranford zealously promotes the concept of "permanent vegetative state" in the face of concern that this implies unquantified finality. Yet he suggests that some patients recovered awareness under the observation and care of Andrews and colleagues. Thus the patients were not misdiagnosed: their diagnosis changed. On the basis of the Multi-Society Task Force's criteria,⁵ 70% of these patients were "permanently" vegetative. How amazing that most patients suddenly emerged from this state, and their diagnosis changed, 16 days after their admission to the Royal Hospital for Neurodisability.

Despite Cranford's scepticism we continue to encounter misdiagnosis of the vegetative state in our practice. Further study of patients diagnosed as being in this state, dissemination of current diagnostic criteria, and education of doctors about the difficulties of evaluation are sorely needed.

NANCY L CHILDS

Medical director, inpatient brain injury programme

WALT N MERCER

Director of research, psychology department

Healthcare Rehabilitation Center,
1106 West Dittmar Road,
Austin, TX 78745,
USA

- 1 Andrews K, Murphy L, Munday R, Littlewood C. Misdiagnosis of the vegetative state: retrospective study in a rehabilitation unit. *BMJ* 1996;313:13-6. (6 July.) [With commentary by R Smith.]
- 2 Childs NL, Mercer WN, Childs HW. Accuracy of diagnosis of persistent vegetative state. *Neurology* 1993;43:1465-7.
- 3 Cranford R. Misdiagnosing the persistent vegetative state. *BMJ* 1996;313:5-6. (6 July.)
- 4 Miller RM, Groher ME, Yorkston KM, Rees TS. Speech, language, swallowing, and auditory rehabilitation. In: DeLisa JA, ed. *Rehabilitation medicine: principles and practice*. Philadelphia: Lippincott, 1988.
- 5 Multi-Society Task Force on PVS. Medical aspects of the persistent vegetative state. *N Engl J Med* 1994;330:1499-508.

Young female doctors report achieving fewer surgical skills than young male doctors

EDITOR,—In their paper on the career preferences of British doctors Trevor W Lambert and colleagues highlight the different choices of career between men and women.¹ As in the United Kingdom, in Norway the proportion of female graduates has increased in recent decades and is currently 50% (roughly 200 graduates); this change greatly influences recruitment to the different specialties. For surgery the same gender imbalance reported by Lambert and colleagues is observed in Norway, where only 27 of 973 specialists in general surgery are women. An interesting finding in Lambert and colleagues' paper is the important role of self appraisal of skills and aptitudes in the choice of a future medical career.

As part of an assessment of the Norwegian internship we sent a questionnaire to a cohort (n = 133) of Norwegian preregistration house officers who had recently finished their hospital service; 103 (77%) replied.² We asked them whether they thought they had achieved particular skills in several procedures. The female doctors reported having achieved fewer surgical skills than did their male colleagues (table 1).

Female house officers have the same potential as men to achieve surgical skills and eventually to start a surgical career. Hostile attitudes towards women have been reported among surgeons and could account for the women's limited skills, as could the absence of women in surgery.³ Because self appraisal of clinical skills is important in doctors' later choice of career, and because the

recruitment of women to surgery is low, those who teach surgery at both undergraduate and postgraduate levels should address this problem.

GEIR FALCK

Research fellow

Norwegian University for Science and Technology,
Department of Pharmacology and Toxicology,
Medical Technical Centre,
N-7005 Trondheim,
Norway

GUTTORM BRATTEBØ

Senior registrar

Norwegian Medical Association,
N-1324 Lysaker,
Norway

- 1 Lambert TW, Goldacre MJ, Edwards C, Parkhouse J. Career preferences of doctors who qualified in the United Kingdom in 1993 compared with those doctors qualifying in 1974, 1977, 1980, and 1983. *BMJ* 1996;313:19-24. (6 July.)
- 2 Falck G, Brattebø G, Aarseth O. Er turnuskandidater opplæring i praktiske prosedyrer god nok? [Is the training of preregistration house officers in practical clinical skills adequate?] *Tidsskr Nor Lægeforen* 1995;115:2091-5.
- 3 Caiach S. Training doctors. *BMJ* 1995;301:1473.

Advertisement for toremifene

Playing on women's fears is unethical

EDITOR,—We are concerned by the advertisement for Fareston (toremifene) published in the clinical research edition of the *BMJ* on 6 July (between p 24 and p 25) and in other clinical journals. The advertisement focuses on patients' perceived concerns about the toxicity of tamoxifen—specifically, its carcinogenic potential—and proposes toremifene as an alternative that could allay those fears. The issue of endometrial cancer and tamoxifen has stimulated much debate.¹ It should, however, be seen in context. The national surgical adjuvant breast and bowel project, a trial of adjuvant tamoxifen, showed roughly a twofold increased risk of endometrial cancer, but overall benefit far outweighed this risk.²

Whatever the concerns about tamoxifen, to use them in promoting toremifene is inappropriate. Firstly, toremifene does not have a licence for use as adjuvant treatment, the setting in which the question of endometrial cancer has been raised. Toremifene is licensed for use only in patients with advanced breast cancer. There is no evidence that the occurrence of endometrial cancer affects the outcome for this group of women treated with tamoxifen. Secondly, although there may be experimental grounds for believing that toremifene is less likely to be carcinogenic than tamoxifen, there are no clinical data to support this contention. Not surprisingly, in a randomised comparison of tamoxifen and toremifene no patient developed endometrial cancer.³ Indeed, there was no difference in either activity or adverse events between the two treatments. Thirdly, the datasheet for toremifene specifically describes "a risk of endometrial changes including hyperplasia, polyps and cancer," and toremifene is contraindicated in women with pre-existing endometrial hyperplasia. Finally, the potential risks associated with tamoxifen come with prolonged use, yet the datasheet for toremifene warns that experience of use of toremifene for more than one year is limited. By contrast, there has been more than seven million woman years of use of tamoxifen.

As clinical trialists we support the development of better treatments for women with cancer. Toremifene may find a place in the treatment of breast cancer. This advertisement, however, is unbalanced and misleading, uses preclinical data unsupported by clinical studies, and seems to contravene the Association of the British Pharmaceutical Industry's guidelines. Patients' concerns are important, but it is surely unethical to play on these concerns when there

Table 1—Number (percentage) of procedures in which men and women reported having achieved skill

	Men (n = 56)	Women (n = 47)
Resuscitation (10 procedures)	109 (19)	85 (18)
Medical emergencies (11 procedures)	305 (50)	221 (43)
Laboratory procedures (8 procedures)	213 (48)	162 (43)
Surgical procedures (20 procedures)	774 (69)	556 (59)*
All procedures (n = 49)	1401 (51)	1024 (44) †

* $\chi^2 = 22.2$, 1 df, $P < 0.01$. † $\chi^2 = 21.8$, 1 df, $P < 0.01$.