When the potential side effects of physical therapy (such as dependency) and the cost are taken into account there is not enough evidence to promote the maintenance of physical activity in the acute phase of low back pain.

JOOP BRUSSEN

Medical researcher, National Casemix Office

Churchfarm,
8A Station Road,
Morton PE10 0NN

- 1 Jayson MIV. ABC of work related disorders: back pain. *BMJ* 1996;313:355-8. (10 August.)
- 2 Malmivaara A, Hakkinen U, Aro T, Heinrichs ML, Koskenniemi L, Kuosma E, *et al.* The treatment of acute low back pain—bed rest, exercises, or ordinary activity. *N Engl J Med* 1995;332:351-5.
- 3 Faas A, Van Eijk JTHM, Chevannes AW, Gubbels JW. A randomised trial of exercise therapy in patients with acute low back pain. *Spine* 1995;20:941-7.
- 4 Koes BW, Van Tulder MW, Van der Windt DAWM, Bouter LM. The efficacy of back schools: a review of randomised clinical trials. *J Clin Epidemiol* 1994;47:851-62.
- 5 Shekelle PG, Adams AH, Chassin MR, Hurwitz EL, Brook RH. Spinal manipulation for low-back pain. *Ann Intern Med* 1992;117:590-8.

Use of interferon beta 1b for multiple sclerosis

Patients will put up with side effects because of their desire to beat the disease

EDITOR,—As a physician who has both multiple sclerosis and a doctorate in clinical pharmacology, I am probably in a unique position to answer Peter Harvey's question of why interferon beta 1b was licensed.¹

Multiple sclerosis, particularly the relapsing-remitting form, is a notoriously difficult illness on which to perform clinical trials. This probably accounts for the large number of initially promising treatments that later proved to be useless. To my knowledge, interferon beta 1b is the only treatment that has been shown to have a significant beneficial effect on multiple sclerosis in large randomised controlled trials. If one reads the results of the trials carefully (as Harvey obviously has) it is apparent that the drug is not the panacea that the popular press would lead us to believe that it is. It also has unpleasant side effects. These are the reasons why I do not take it myself. Its development and the trial methods, which used magnetic resonance imaging, have, however, led to hope of better treatments to come and better ways of evaluating them.

Harvey has not taken into account another important factor, which is the desperate desire of people with multiple sclerosis (and their families) to do something to treat the disease. This desire becomes especially strong during an acute relapse. Unless something better is discovered before then, I will probably end up taking interferon beta 1b after my next bad relapse.

MIRIAM CHELLINGSWORTH
Retired consultant geriatrician

1 Montville Drive,
Stafford ST17 9XJ

- 1 Harvey P. Why interferon beta 1b was licensed is a mystery. *BMJ* 1996;313:297-8. (3 August.)

Manufacturer defends the drug

EDITOR,—It was disappointing to read Peter Harvey's letter dismissing the value of interferon beta 1b in the treatment of patients with relapsing-remitting multiple sclerosis.¹ Harvey takes no account of the results of controlled clinical trials showing a significant reduction in severity and frequency of relapses (up to 30-50%),^{2,3} with the most severe cases benefiting most, or of the accumulating data showing that generally there is a reasonable correlation between findings on magnetic resonance imag-

ing and clinical status.³⁻⁵ Individual cases do not render the overall evidence invalid.

Harvey fails to recognise that a reduction in the frequency and severity of attacks can make an immense difference to the quality of life of a patient with multiple sclerosis. It can mean the difference between being able to work or not, being able to go shopping or not, and between independence or reliance on others. To dismiss these improvements as inconsequential does a disservice to both patients and the drug.

While it is true that some patients experience side effects, such as flu-like symptoms in the first few weeks of treatment, simple measures can be taken to minimise this; in our experience the dropout rate due to side effects in the first six months of treatment is of the order of only 3%. The issue of neutralising antibodies is not a simple one as suggested by Harvey, and antibody titres do not correlate strongly with a reduction in the drug's effectiveness. Above all, it must be a clinical judgment whether an individual patient should continue treatment, depending on the benefit received.

As clinical experience with interferon beta 1b increases I hope that those who adopt polarised positions will come to modify them. In the meantime, Schering will put the evidence forward by whatever means possible, including in the *BMJ*; this is especially important when contacts of any kind with the company are spurned, as in Harvey's case. It is patients who stand to suffer by being denied an effective treatment.

P N LONGTHORNE
Medical director

Schering Health Care,
Burgess Hill RH15 9NE

- 1 Harvey P. Why interferon beta 1b was licensed is a mystery. *BMJ* 1996;313:297-8. (3 August.)
- 2 IFNB Multiple Sclerosis Study Group. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. I. Clinical results of a multicentre, randomised, double-blind, placebo-controlled trial. *Neurology* 1993;43:655-61.
- 3 IFNB Multiple Sclerosis Study Group and the University of British Columbia MS/MRI Analysis Group. Interferon beta-1b in the treatment of multiple sclerosis: final outcome of the randomised controlled trial. *Neurology* 1995;45:1277-85.
- 4 Filippi M, Horsfield MA, Morrissey SP, MacManus DG, Rudge P, McDonald WI, *et al.* Quantitative brain MRI lesion load predicts the course of clinically isolated syndromes suggestive of multiple sclerosis. *Neurology* 1994;44:635-41.
- 5 Van Walderveen MAA, Barkhof F, Hommes OR, Polman CH, Tabi H, Frequin STFM, *et al.* Correlating MRI and clinical disease activity in multiple sclerosis: relevance of hypointense lesions on short-T2/short-TE (T1-weighted) spin-echo images. *Neurology* 1995;45:1684-90.

Importance of relapses must not be underestimated

EDITOR,—Peter Harvey asserts that patients with multiple sclerosis are interested in disability and, by implication, are not interested in relapses.¹ While many patients with multiple sclerosis cope successfully with relapses, for a considerable proportion of patients relapses are frequent and severe, causing great distress and disruption to normal life. It is strange to condemn the first licensed product to show an effect on relapses because it has not also been shown to affect disability without acknowledging the importance of relapses in patients' lives.

Equally, it is rather negative to place so much emphasis on the side effects of this drug without giving equal weight to the potential benefits. Experience even before the product was licensed in 1995 shows that patients with multiple sclerosis are capable of weighing the evidence on treatments and reaching rational conclusions; hence the lack of the feared stampede for interferon beta 1b. The Multiple Sclerosis Society has put great effort into ensuring that the evidence has been available to patients in a balanced and intelligible form and has looked to doctors to do likewise.

Harvey's view is incapable of defence without better evidence than anecdote from the narrow context of the clinic. In Britain the shortage of neurologists means that few neurologists can afford extensive contact with their patients with multiple sclerosis, and the waiting times for outpatient appointments and admissions ensure that patients are rarely seen by their consultant during an exacerbation. It is not surprising, therefore, that the importance of relapses to patients is often underrated by British neurologists, and this may go some way to explaining the differences in levels of prescribing of interferon beta 1b between Britain and other countries.

Harvey falls into the same trap as the authors of the landmark studies of interferon beta^{2,3} in making assumptions about the treatment priorities of patients with multiple sclerosis. In the context of carefully designed, multimillion dollar clinical trials it seems strange to those of us in daily contact with large numbers of patients that neither investigators nor pharmaceutical companies have sought firm evidence on this. Given the current range of possibilities, would patients with multiple sclerosis prefer their relapses to be treated or not? This is an important issue that has not yet been investigated, even though further costly large scale trials continue to be advocated. The Multiple Sclerosis Society is interested in potential collaborations on this question and would be pleased to hear from others of like mind.

PETER CARDY
Chief executive

Multiple Sclerosis Society,
London SW6 1EE

- 1 Harvey P. Why interferon beta 1b was licensed is a mystery. *BMJ* 1996;313:297-8. (3 August.)
- 2 IFNB Multiple Sclerosis Study Group. Interferon beta-1b is effective in relapsing-remitting multiple sclerosis. Clinical results of a multicentre, randomised, double-blind placebo-controlled trial. *Neurology* 1993;43:655-61.
- 3 Jacobs LD, Cookfair DL, Rudick RA, Herndon RM, Richert JR, Salazar AM, *et al.* Intramuscular interferon beta-1a for disease progression in relapsing multiple sclerosis. *Ann Neurol* 1996;39:285-94.

Practice in endoscopic cholangiopancreatography in Denmark differs from that in Britain

EDITOR,—Jacob Rosenberg and colleagues' conclusion that premedication with metoprolol prevents myocardial ischaemia during endoscopic cholangiopancreatography has important implications for routine technique in this procedure.¹ The applicability of this finding to current clinical practice in Britain, however, is unclear since there are considerable differences between the British Society of Gastroenterology's guidelines and the practice in this Danish study.²

In the study group patients did not receive preoxygenation with supplemental oxygen, morphine rather than pethidine was used as the opiate, and larger doses of hyoscine bromide were used than is common in Britain. The use of lignocaine spray in heavily sedated patients is also contrary to the British Society of Gastroenterology's advice. The high incidence of myocardial ischaemia (53%) and the presence of myocardial infarction in the relatively small group of 19 patients treated with placebo suggests that placebo conditions may have been substantially suboptimal. Continuous monitoring of blood pressure during endoscopic cholangiopancreatography is currently not routine in many centres in Britain but would clearly be necessary if premedication with a β blocker was used.

The study highlights the importance of tachycardia and illustrates the risk of myocardial ischaemia, but further data will now be required to determine whether premedication with metoprolol improves safety in patients receiving