and there was no adventitious sound, and no cough or expectoration. The only area of dullness was the one alluded to. This was triangular in shape, with the apex at the fifth rib in the mid-axillary line, becoming broader towards the costal margin, where it was continuous with the swelling felt below. There were no vocal fremitus, *râles*, or crepitation to be detected over the area. The uterus was repeatedly irrigated, and portions of fetid secundines removed. Quinine and opium were administered internally, and enemata to relieve the bowels.

Progress.—June 27th. Condition much improved; there was no sickness, and she was able to take liquid nourishment. Temperature, 102° ; pulse, 108; respirations 32. I was called hurriedly on June 30th, and the patient complained of pain in the left side over the dull area, which I found increased. There was pain at the end of inspiration, and a pleuritic rub was audible over the area. Temperature 104° , pulse 124; respirations 36. On July 2nd, Dr. Tudor Thomas, who visited her for me, found symptoms of effusion into the pleura, and her condition very grave. On July 3rd I made an exploratory examination with a long needle of large bore in the seventh space in the mid-axillary line, and from a depth of $2\frac{1}{2}$ in. withdrew a syringeful of thick, reddish offensive pus, and an operation was advised as the only means of saving the patient.

Operation.—The condition of the patient was extremely grave. The pulse 140, temperature 103° , respirations 40. It was a question whether she would stand the shock of the operation. Dr. Edgar Davies, who gave the anaesthetic, examined the patient and found incidentally a patch of consolidation in the right base. I made a $3\frac{1}{2}$ in. incision over the eighth rib and removed about 2 in. of its length, and on opening the pleural cavity a considerable quantity of thin blood-stained serum escaped from the sac. Portions of costal pleura were excised, and in a window about 2 in. square, the lung presented. The needle was again plunged into abscess and a long sinus forceps was with difficulty forced into the abscess cavity, but so dense and tough was the lung substance that I had to introduce a grooved director and cut it freely with a scalpel before I could introduce the finger. About 10 oz. of thick reddish and most offensive pus was evacuated. The cavity had a craggy uneven wall and extended to the pericardium, the finger feeling in detail the movements of the heart in its sac. Iodoform was introduced into the abscess cavity, two large drainage tubes inserted, the wound around the tubes packed with iodoform gauze, a large padding of alembroth wool applied, and the patient put back to bed. Dr. Davies injected strychnine before and during the administration of the anaesthetic. She bore the operation (which did not last 20 minutes) well.

Progress.—The case was dressed daily, and a great deal of pus continued to be discharged for ten or twelve days. The temperature continued high, 101° to 103° , and though she took nourishment well her condition continued grave through the continuous septic absorption. I now irrigated the abscess cavity with 1 in 40 carbolic, and in this way removed several masses of lymph or necrotic tissue, and improvement was soon evident. Her temperature went down to 99° to 101° , appetite improved, and she began to gain flesh, but the discharge continued in lessening quantities until September 2nd, when the tube was no further used.

She has made a perfect recovery. I examined her chest recently, and the physical signs are normal, excepting a circular area of an inch and a half in diameter over the cicatrix.

REMARKS.

In Case I the signs of pleuritic effusion masked those of pulmonary abscess, and the operation was performed with the idea that the condition was one for empyema. Had I known the case was one of tuberculous abscess of the lung I should not have advised operative measures; but once committed to it, the operation was completed. The result justified the procedure, for the patient's health was improved, and, I have no doubt, her life prolonged. The results of operation on tuberculous cavities have not been satisfactory, and I would not recommend operation excepting in large basic cavities, especially if septic, and then only in the hope of relieving the constant distressing cough and arresting septic absorption.

Case II I have reported in some detail as an example of those that should in my opinion be operated upon. Had this been done in January, in all probability the patient would have recovered. It is a curious pathological fact that most basic abscesses due to pneumonia perforate downwards, and those secondary to abdominal lesions generally open into the bronchus. In my earlier experience three basic abscesses (two following acute lobar pneumonia and one following enteric fever) perforated into the abdominal cavity, and one following typhilitis (or appendicitis) and subphrenic abscess perforated finally into the bronchus, death being due to septic absorption. No attempt was made, or would then have been regarded as justifiable, to operate. In my opinion all were cases that should have been operated upon and with every prospect of success. When non-tuberculous, I regard the aphorism "*Ubi pus ibi incisio*," as being as applicable to the lung as to any other organ, provided Paquelin's cautery and a scalpel are regarded as convertible terms. In Case II every condition favourable to operation were present. The abscess could be definitely

located, the costal and pulmonary pleurae were adherent and yet fatal delay was recommended and adopted by the friends, for the intercession, I suppose, of some special providence.

In Case III the diagnosis was fairly clear when I first saw the patient, and the operation was performed shortly after the appearance of the secondary pleurisy. As to the operation, when the abscess can be accurately located the opening into the pleura need not be large. A long linear incision down to the rib and the removal of 2 in. of its length will be enough. It is stated by Truc that the lung should not be opened if the costal and pulmonary pleurae are not already adherent or sutured together so as to shut off the pleural cavity. This was not done, though the cavity was at any rate partially open, and yet no empyema ensued. The lung tissue was so dense and infiltrated that it was thought safe to open with the knife and not with the cautery. There was scarcely any haemorrhage. It is also laid down as a rule by the same authority that on no account should the abscess cavity be irrigated. For some time after the operation there was evidence of septic absorption, the temperature continued high, and the patient's condition most precarious. There being no communication between the abscess and the bronchus, I irrigated the cavity daily with a 1 in 40 carbolic lotion with immediate benefit, and with no other untoward result than a transient carboloria, which I met by using a weaker solution.

A CASE OF PURULENT PERICARDITIS ASSOCIATED WITH EMPYEMA IN A CHILD AGED TWO AND A-HALF YEARS.

DEATH NEARLY FOUR MONTHS AFTER A DOUBLE OPERATION.

BY

J. A. COUTTS, and
M.B.Cantab., F.R.C.P.,
Physician, East London Hos-
pital for Children.

R. P. ROWLANDS,
M.S.Lond., F.R.C.S.,
Assistant Surgeon, East London
Hospital for Children; Demon-
strator of Anatomy, Guy's
Hospital.

PURULENT pericarditis is a not infrequent disorder in children, but cases of operation for its relief are sufficiently rare to justify the putting each one on record. The present case is a specially noteworthy one, both from the youthful age of the patient and the time he lived after the operation. The notes are abridged and slightly altered from the careful ones taken by Dr. A. Emerson, formerly House-Physician at the East London Hospital for Children.

M. P., a Jewish child, aged $2\frac{1}{2}$ years, was admitted into the East London Hospital for Children on February 11th, 1903, under the care of Dr. Coutts.

Family History.

In the family history there was little to note. The father had been drowned. The mother was healthy, and there were no other children. There was no family history of tuberculosis.

History.

The present illness commenced with pneumonia a month ago, and the child has never been well since. He has had a bad cough ever since pneumonia, and he has suffered from loss of appetite and languor, with occasional vomiting. The bowels were regular. During the past few weeks he has wasted somewhat.

Condition on Admission.

The patient was a fairly well-nourished child, presenting distinct evidence of rickets. There was well-marked beading of the ribs, along with slight enlargement of the epiphyses of the long bones with bowing of the tibiae. The abdomen was not distended, the liver was not enlarged, but the spleen could just be felt below the costal margin. On examination of the thorax, the cardiac dullness was found to be slightly increased, and the heart's apex beat almost impalpable. The heart sounds were weak, but apparently pure. There was no audible pericardial friction. The pulse was 136, regular in force and rhythm, and of fair tension. On the right side posteriorly the percussion note was impaired at the apex, and from the sixth rib downwards absolutely dull. This dull area extended into the axilla and right round to the front of the chest. Over the dull area the breath sounds were extremely weak and faintly bronchial in character. Over the rest of the right chest both percussion and breath sounds were normal. The left lung was normal both to percussion and auscultation, with the exception of a patch at the extreme apex where there was some doubtful bronchial breathing and increased vocal resonance. The respirations were rapid—58 per minute. The child was wakeful and peevish, and seemed in pain; giving a short grunt with each expiration. The tongue was clean, but the appetite poor. There was no diarrhoea or vomiting. On admission the temperature was 99° , but in the evening it rose to 101° .

Progress.

February 13th. The child still looked very ill. Since admission the temperature has varied greatly, reaching on one occasion 104°. The right base was explored twice, but nothing found except a little blood. The needle appeared to touch the lung.

February 16th. The child was much worse and his condition critical in the extreme. The signs over the right lung remained as before, and there were signs of bronchitis over the left. The cardiac dullness had increased and reached to the upper border of the second rib above, and below extended from an inch outside the left nipple to the mid-sternal line. The apex beat could not be felt, and the heart's sounds were almost inaudible. No pericardial rub could be heard. The pulse was very rapid and weak, but regular. This afternoon Dr. Coutts saw the child for the first time and acquiesced in the diagnosis that had already been made of a probable empyema at the right base with a purulent pericarditis. A consultation was held with Mr. Rowlands, who agreed to operate on condition of the mother's consent being obtained.

Operation.

At 9 o'clock at night Dr. W. P. S. Branson gave chloroform, and Mr. Rowlands opened the pericardium through an oblique incision, exposing the fifth left costal cartilage and the neighbouring sternum. The sternal half of this cartilage having been removed with care to avoid perforation of the left pleura, the internal mammary vein was seen greatly distended and lying close the sternal margin. Some fibres of the triangularis sterni were separated, and the thin parietal pleura was seen to extend inwards under the sternal margin; therefore a semi-lunar piece of the still cartilaginous sternum was pared away, the margin of the left pleura discovered and displaced outwards, thus exposing the pericardium in the depth of the inner angle of the wound. An exploring needle was inserted, and withdrew a syringeful of thin, greenish-yellow pus. On incising the pericardium along the needle, the fluid rushed out, reaching to a height of three feet; probably about half a pint escaped, but it was impossible to collect it for measurement. A finger introduced into the pericardial sac failed to discover any masses of lymph, therefore irrigation was not resorted to; but a rubber drainage tube was inserted, and the rest of the wound closed.

The immediate results of the operation were very striking, the internal mammary vein was seen to collapse at once, the cyanosis diminished, the pulse was stronger and better, and its rate and that of respiration diminished. It was not thought well, however, to deal at this juncture with the possible empyema at the right base, and the child was accordingly placed back in bed.

The pathologist's report on the pericardial fluid stated that it contained a coccus which morphologically and from its staining properties and growth on culture media resembled the pneumococcus.

After-History.

February 17th. The improvement after the operation of the night before being maintained, the right pleura was again explored and pus was found. The child was again placed under chloroform and Dr. Branson excised a piece of rib in the post-axillary line, and with some difficulty a small quantity of pus was discovered and evacuated. The pathologist's report on the empyema pus was as follows: "An extremely impure culture; subcultures have not been found to be of much value. A coccus is evident which by culture and morphology resembles the pneumococcus. Other forms of cocci and bacilli are also numerous."

February 22nd. Very slight amount of purulent discharge from the pericardial incision and the empyema opening. The child is holding its own well. The pulse remains good. The cardiac dullness is still much increased, but the heart sounds are normal and clear. The temperature for the last two days has been practically normal.

March 2nd. The tube has been given up in the pericardial wound for the last few days, and a small drain of gauze used instead. There has been very little discharge from either wound to-day, but yesterday there was a fairly large quantity from the pericardial one. For the last week the temperature has been markedly swinging, and has sometimes reached 103° or 104°. The pulse is small and rapid.

March 9th. During the last week the condition of the patient has varied, but to-day he seems a little better. The temperature has fallen to normal. The pericardial wound discharges freely, but the cardiac dullness has diminished and the heart sounds are regular and more distinct. The empyema wound looks dry and the discharge has ceased. The child appears to have still a good deal of pain with cough, and the ensiform cartilage appears to be dislocated upwards and forwards with each inspiration.

April 6th. Since the date of the last note the child's condition has varied from day to day, but on the whole he has improved. The pericardial wound has discharged pus freely, but for the past week the discharge has lessened and to-day has practically stopped. There is still, however, a small sinus in the pericardial wound, but the empyema one has healed. For most of the time the temperature has been about the normal, but for the last few days it has begun occasionally to rise again to 100° or 101°.

April 13th. Dry scales over the two wounds. The cardiac dullness extends from the left nipple line to the left sternal border. Heart sounds clear. Yesterday the temperature, which had been irregular, reached 103°, and the child vomited several times. To-day there are signs of slight bronchial catarrh over both lungs.

April 23rd. The child has been very ill for the last three days. The temperature has been higher and varied less. There has been a purulent discharge from both eyes and nose, and the glands below the jaw are enlarged. To-day he developed a measles rash and was removed to

the isolation block. Yesterday the pericardial wound broke open again and a small quantity of purulent discharge came away.

April 27th. Measles rash almost disappeared; temperature lower. There are signs of bronchitis over the left lung, and in the right axilla are some tubular breathing and increased vocal resonance, considered to be due to a consolidation of the middle lobe of the right lung. The pericardial wound is discharging freely.

May 2nd. There is no improvement in his general condition. The signs of consolidation in the right lung persist. There is a copious discharge from the pericardial wound, and there is well-marked oedema of the feet and ankles. The tongue is clean, and he is taking his food fairly well.

May 7th. The signs of consolidation of the right lung have cleared up, and the oedema of the feet disappeared. The child's condition, however, remains very variable, and he gets very restless at nights and appears to be in a good deal of distress. The pericardial wound discharges freely, and the surrounding skin is red and irritated.

In the rest of the history there is little to note. The child's recovery in the weak condition he was in from a severe attack of measles and broncho-pneumonia was totally unexpected. In the later weeks of his life a permanent recovery never seemed probable. A week before death he developed two bedsores posteriorly, and for the last two months he was emaciated in the extreme, and his skin covered with a profuse branny desquamation. The discharge from the pericardium, after its re-establishment, showed no tendency to diminish, but remained profuse and purulent to the end. The severe precordial pain, too, which had almost gone, reappeared after the measles and persisted until death. In the last few weeks of life tenderness to the gentlest percussion or palpation over the precordium became more marked. The temperature remained very irregular, but scarcely to be called high, until the last few days of life. The child gradually sank, and died on May 22nd, in the sixteenth week after the two operations.

Necropsy.

Notes on the necropsy by Dr. W. P. S. Branson, Pathologist to the Hospital.

On opening the thorax there was found a condition of universal dense adhesion of the contained viscera to each other and to the chest wall, and the whole anterior mediastinum was occupied by a thick fibro-gelatinous deposit. The wound over the pericardial area drained a small pocket on the anterior surface of the heart, but elsewhere the pericardium was inseparably adherent to the heart in front. On attempting to separate the pericardium from the left lung an abscess cavity was opened, with a discharge of 3 oz. of pale thick pus. This cavity lay behind, and to the left of the heart and within the pericardial sac; its walls were very thick from the presence of fibrinous matter similar to that occupying the mediastinum. Between the heart and the right lung there lay another smaller cavity, independent of the first, and full of very inspissated pus. The pus in both cavities was extremely dry and gritty to the feel, as if impregnated with lime salts. The tracheal glands were greatly enlarged, but free from all sign of tubercle. The pleurae were everywhere adherent, but not inseparably except at the right base behind the site of the wound made to drain the empyema. Here the adhesions were old and fibrous and no trace of a pleural membrane could be discovered. Elsewhere the pleurae were much thickened, again with what appeared to be a fibro-gelatinous infiltration. The lung tissue was very dark throughout, but crepitant and not engorged. The liver showed some signs of the "nutmeg" order, and the spleen and kidneys were also engorged. Examination of the head was not allowed.

REMARKS BY DR. COUTTS.

The frequency of purulent pericarditis in childhood is only appreciated by those who have special opportunities of witnessing necropsies in young children. That the complaint is usually first discovered on the *post-mortem* table is due to the large majority of cases being undiagnosable during life, owing to the scantiness of the pericardial effusion and the absence of all symptoms referable to it. Certain cases that might have been diagnosed during life are missed because the condition of the patient has been too urgent to admit of any satisfactory or adequate examination. Other cases, again, go unrecognized owing to the reluctance to harass the child by having the dressings over an empyemic wound removed in order to examine the chest. The complaint, however, is so frequent in infants dying with empyemata that the possibility of its being discovered *post-mortem* should not be lost sight of, although there may never have been any recognizable signs of pericarditis during life.

To calculate roughly the number of cases of purulent pericarditis occurring annually at the Shadwell Hospital at about a dozen would probably be to under-estimate its frequency at that institution. A limited number of such cases occur in the course of a suppurative bone disease, whilst practically all the rest have a causal origin in the presence of the pneumococcus. As might be expected from this, then, purulent pericarditis in children, in the vast majority of instances, is associated with pneumonia or empyema, or a combination of these two last. In brief notes of 15 cases of purulent pericarditis under my care, where a necropsy was made, in 8 cases

it was associated with empyema alone, in 4 with evidence of recent pneumonia alone, and in 2 with a combination of recent pneumonia and empyema. In only one case was there no evidence of either of these last. This association with recent pneumonia or empyema may afford a valuable aid, often an indispensable one, in the diagnosis of the complaint.

The combination of purulent pericarditis and empyema is much more frequent than is commonly supposed. In a valuable paper on empyemata in children,¹ founded on a personal observation of 81 cases, Mr. P. S. Blaker gives statistics which show that purulent pericarditis occurs in 11 per cent. of empyemata in children of all ages, and where infants alone are taken into account this percentage becomes much greater. In 23 cases of fatal empyemata, the same observer found purulent pericarditis *post mortem* in 7; or an average of about 30 per cent. An experience of a much larger number of cases than 81 would not induce me to believe there is anything exceptional in Mr. Blaker's figures, and I believe they may be taken as fairly representative of the frequency of the association of the two conditions, empyema and purulent pericarditis.

There is, unfortunately, a prevalent impression, and one often expressed, that the diagnosis of purulent pericarditis is in every case too difficult, or even impossible, to be made with any degree of precision. In the absence of empyema or suppurative bone disease, or in that of a history of recent pneumonia, there might be justification for this pessimistic view. But given the presence of one of these three last, then I would contend that the diagnosis can at times be made with a reasonable degree of certainty. Under such conditions, in fact, the possibility of the diagnosis mainly rests on that of determining the presence of free fluid in the pericardial sac. That cases arise where it is impossible to distinguish between fluid in the pericardium and a cardiac dilatation has to be admitted. But cases where insuperable difficulties exist in differentiating between the two are, in my opinion, mainly those of rheumatism of some standing, where, in addition to the cardiac dilatation, there is also thickening of the pericardium itself.

The conditions, moreover, associated with purulent pericarditis are not such as to lead frequently to a cardiac dilatation sufficiently great to embarrass the diagnosis when dealing with the question of fluid in the pericardium. The slighter degrees of pericardial effusion are admittedly undiagnosable, but in cases where there is marked pericardial dullness along with diminution of the cardiac sounds the difficulties of diagnosis have certainly been exaggerated. Many cases are missed solely from the want of a thorough examination, either because the urgent state of the patient forbids it or from a natural reluctance to cause discomfort to the child by the removal of the dressings over an empyema wound. A remembrance of the frequency of a combination of the two complaints, and a preliminary hint as to the possibility of a purulent pericarditis being discovered at necropsies on infants dying with empyemata, will often disarm the satire of the pathologist at the expense of the clinician.

Having satisfied oneself as to the presence of fluid in the pericardium in early cases, there need be but little hesitation as to its character; if not already purulent, it will speedily become so. I can see no advantage, then, in a preliminary exploration before deciding on a more radical operation, if such be contemplated. In more than one instance I have known a left-sided pleurisy added to the other gravities of the case simply from the use of the exploratory needle. On six or seven occasions, too, I have known the cardiac wall to be pierced by the needle, and although in no instance was this attended by any subsequent harm, still the possibility of such a contingency is not one to be contemplated with any satisfaction or complacency. The present is the third case I have had operated on without preliminary exploration, and in each one the assumption as to the nature of the fluid was verified at the operation.

In a few other cases where I have ventured on the diagnosis without a preliminary exploration, and where operation was deemed inadvisable or refused, the supposition as to the presence and nature of the fluid in the pericardium has been justified on necropsy. Occasionally, instead of pus, a merely turbid fluid is found, yielding a pure cultivation of the pneumococcus. There is, however, a growing opinion, which I am inclined to share, that such cases should be treated as if the fluid were already pus, whether it be in the pericardium or the pleura. Very often, indeed, flakes of pus are discovered

at the bottom of a cavity that at first sight is apparently filled only with such turbid fluid.

One reason for purulent pericarditis being so frequently missed is the absence, in the vast majority of cases, of any audible pericardial friction. In the rare cases where the pericarditis arises in connexion with bone disease friction may be loud and persistent, but in the far more numerous ones associated with the pneumococcus an audible rub is usually absent, invariably so in my experience. This absence of audible friction is perhaps due to empyemata and pericarditis, when associated with pneumonia, being in no sense sequelae of the last complaint, their origin dating back to that of the pneumonia itself. If this be so, the time for hearing any pericardial friction would be in the very earliest stages of the pneumonia. Unfortunately, in the early stages of that complaint, the attention is so centred on the predominating lung symptoms that, possibly, an adequate examination over the pericardium is seldom made. But, as a rule, no opportunity occurs of seeing these cases of purulent pericarditis until several weeks after the onset of pneumonia or empyema. It may well be, however, that audible friction is absent from the very beginning of the complaint.

Occasional aids to diagnosis are extreme pallor, orthopnoea, irregular pulse, and other signs of cardiac and respiratory distress. Pallor, indeed, is a fairly constant feature in most cases, but the other symptoms are of very uncertain occurrence. In a case I lately published² there were 10 oz. of pus in the pericardium of a child aged 4 years, with a total absence of any respiratory or cardiac embarrassment.

When all things are considered, it has to be admitted that the diagnosis is often extremely difficult, and in the majority of cases impossible. At other times, as I have said, the difficulties are not insuperable, and the diagnosis can be made with a reasonable degree of confidence. And when the frequency of the association of purulent pericarditis with empyema and pneumonia is more generally recognized, I believe that a diagnosis made during life will with increasing frequency replace that which is at present usually only made upon the *post-mortem* table.

In my present case the diagnosis was attended with no special difficulty. Although there was no audible pericardial friction, yet the history of a recent pneumonia, the presence of a probable empyema, the increase of the precordial dullness, along with the diminution of the cardiac sounds, and the absence of a definable apex beat, the extreme pallor, and the precordial pain associated with both cardiac and respiratory distress, all combined to make the diagnosis of pericarditis a fairly certain one. The history of pneumonia, too, with the concomitant empyema, left little room for doubt as to the character of the fluid in the pericardium. When I first saw the child his condition was critical in the extreme, and it was clear that no measure short of operation offered any hope of relief. The case was put fairly before the mother that there was but faint prospect of recovery after an operation; and that there was a great possibility that the child would die under the anaesthetic, and she elected to give the child the outside chance that surgical interference offered.

As the child lived so long after the operation, the ultimate result was disappointing in the extreme. At the time of the unfortunate contraction of measles, the child was improving in many ways and seemed to have a fair chance of recovery. That the measles and the concomitant broncho-pneumonia shortened life will hardly be doubted. But whether ultimate recovery was possible, with the heart clogged as it was with the pockets of inspissated pus, is open to question; it is equally open to question, moreover, whether recovery was desirable with the heart hampered in the manner it must always have been. Life in any case, under such circumstances, could not have been either a pleasant or prolonged one. The distribution of the pus, a not infrequent one in such cases, raises the inquiry whether the usual operation of opening the pericardium from the front is the best one under such circumstances. The question, however, is a purely surgical one, and I leave it to the more capable hands of Mr. Rowlands to deal with.

REMARKS BY MR. ROWLANDS.

It is now generally believed that incision and drainage are indicated in practically all cases of purulent pericarditis; but owing to difficulties of diagnosis, the operation is frequently too long deferred for a permanent success to be expected; and although temporary relief follows the evacuation of the fluid in most cases, recovery is unlikely, when from delay

such complications as myocarditis, mediastinitis, or pyaemia have arisen.

The anatomical relations of the pericardium are not favourable for the safe establishment of efficient drainage, and this may explain the number and the variety of the operations which have been proposed and practised for this condition. It is impossible to open the sac from the back without also opening the pleura. Having noticed that the right extremity of the oblique sinus of the pericardium was opposite and only about 2 in. away from the vertebral end of the seventh right rib, I removed a portion of this bone, and attempted to displace the pleura from the vertebrae, and thus to reach the pericardium through the posterior mediastinum, but in each of four attempts on the cadaver the pleura was perforated; the vena azygos major and thoracic duct are also in danger of being wounded.

Luschka's statement, that no parietal pleura lies between the pericardium and the inner 1.5 cm. of the fifth left costal cartilage, the inner 2.0 cm. of the sixth, and the inner 3.5 cm. of the seventh, is only true of exceptional cases, but it has been so often quoted that it has been almost generally believed. The researches of Sick and others have since established the fact that it is usual for the left pleural reflexion to lie under cover of the sternum at the level of the fifth costal cartilage, and often also at the level of the sixth. My own observations in the dissecting room during some years convince me of the accuracy of Sick's description. Delorme and Mignon have shown that a pericardial effusion has little or no effect in displacing the pleural margin outwards; but protective pleural adhesions sometimes form when that effusion is purulent. Paracentesis, however useful it may be in some cases of serous pericarditis, cannot afford more than a temporary relief of pressure in pyopericardium, for the pus rapidly and persistently reaccumulates; but as a preliminary to a more radical treatment it may be of considerable service in patients too seriously ill to take an anaesthetic safely. A very striking improvement of the circulation generally occurs after allowing the fluid to escape, and then the treatment may be completed with all the advantages of a general anaesthetic.

In a very few cases recovery has followed mere tapping, but only after the lucky establishment of a sinus along the track of the needle. Paracentesis is not quite devoid of danger, for empyema, pleurisy, and perforation of the heart have resulted in rare cases.

From anatomical considerations and experiments on the cadaver it is certain that the safest place to tap or to insert the needle of the exploring syringe is the left costo-xiphoid angle as suggested by Roberts years ago. The instrument should touch the lower end of the gladiolus and pass backwards upwards and a little inwards behind the sternum until it is felt to enter the cavity of the pericardium, at a distance of not more than 2½ in. from the skin. The pleura and the internal mammary vessels are too far out to be injured, and the peritoneum too low, and the heart, unless adherent in front, lies too far back within its distended sac to be reached if the above precautions be taken. The inner extremity of the fifth left space is probably the next best situation, where the needle should be passed backwards and inwards in close contact with the sternal edge; but although the internal mammary vessels are quite safe, the pleura will certainly be pierced occasionally. The sixth left space is to be preferred, if wide enough at its sternal end, which is not often the case.

Exploratory aspiration through the left xiphoid fossa deserves more frequent use in doubtful cases, so that delay in diagnosis and treatment may be avoided if possible.

Operations.

Which is the best operation is a question still to be answered by time and experience.

1. *Simple incision* through the fifth (or fourth) left intercostal space near the sternum has been the method most frequently employed and can be very quickly performed, which is an advantage in these grave cases; but the drainage is far from perfect, especially in children, and the pleura and internal mammary vessels are difficult to define and avoid through the narrow space available.

2. *Excision of the Fifth Left Costal Cartilage.*—Parker was the first to perform this operation on a child at the East London Hospital. It has since been strongly recommended by Porter and others, and has of late become the most popular operation. More room is given for exploration and also for drainage, and the pleura and vessels can generally be seen,

displaced, and avoided; it is, however, not possible to explore the pericardial sac thoroughly and to remove masses of lymph from the auricular regions. Moreover, the wound is not placed low enough on the anterior wall of the sac, so that a pool of fluid remains undrained even when the body is upright. In the case above related, having excised the sternal end of the left fifth costal cartilage, I also removed a semilunar portion of the cartilaginous gladiolus in order to avoid injury of the pleura and to provide better and more permanent drainage. As far as I know, this was the first occasion on which this modification of Parker's operation was employed in actual practice; but Durand and Dwight had each suggested it several years ago. This step, which only takes a few seconds, may be found useful if Parker's operation cannot be completed without endangering the delicate and perhaps extensive pleura of a small child; it gives an excellent view of the pericardium, but in my case the drainage it afforded was not perfect, as proved at the necropsy, and it is possible, or even probable, that Allingham's operation would have been more satisfactory in this important respect.

3. *Flap Operation.*—Roberts,³ in a most instructive and exhaustive paper, proposed a new operation based on experiments on the dead body. A flap containing the fourth and fifth left costal cartilages is lifted upwards and the internal mammary vessels and pleura can then be well seen and displaced outwards, so that the pericardium can be fearlessly opened by a vertical incision. The chief objections to this operation are its unnecessary severity and comparatively long duration in a patient whose circulation is seriously embarrassed; moreover, it is by no means so easy to raise the flap with safety to the pleura and vessels, especially in children. I once opened the pleura in performing the operation on a stillborn but mature infant. It may, however, be of use after pressure symptoms have been relieved by aspiration, as Roberts himself suggests. It certainly gives a better view of and access to the pericardium than any other operation, but it does not provide as efficient drainage as Allingham's operation.

4. *The Epigastric Operation.*—Allingham⁴ in 1900 gave a full account of an operation performed by him on the cadaver. In this operation the left rectus abdominis is incised, and the peritoneum having been avoided, the cellular interval between the sternal and costal fibres of the diaphragm is opened up, and the pericardium thus exposed and incised at the lowest part of its anterior wall. Allingham also claims that a finger can be passed to any part of the sac to remove debris and open loculi, and that the pleura is not in danger.

The following are some of the chief drawbacks of this ingenious operation:

(a) The little room available in most adults, who have wide, firm, or even ossified ensiform cartilages and rigid costal cartilages; the costo-xiphoid space is too narrow in these cases.

(b) The operation is necessarily performed somewhat in the dark, and under cover of the sternum and seventh costal cartilage, and it is not easy to be sure whether the exploring finger is above or below the diaphragm, so that the peritoneum may be opened, especially by an operator not quite familiar with the anatomy of this region.

(c) The superior epigastric artery may be wounded as it comes through the diaphragm, and cause troublesome haemorrhage out of sight in the depth of the wound.

(d) The pericardium may be separated by the finger from the parietes, and pus may then leak into the loose connective tissues and set up a fatal mediastinitis.

When the pericardium is much distended, these dangers and difficulties are diminished. In children, who are the most common subjects of suppurative pericarditis, the operation is much easier owing to the elasticity of the boundaries of the costo-xiphoid angle. In them a large drainage tube can be easily introduced, as I have found in several stillborn children. This operation is without doubt the best one to adopt in children.

In adults I have not been able to pass my fingers through the costo-xiphoid angle into the basal sinuses of the pericardium, nor even satisfactorily into the oblique sinus. In them Allingham's modification of his own operation will be required in most cases; a portion of the seventh and sometimes even of the sixth costal cartilage will have to be removed to provide the necessary room for exploration and drainage; the pleura must be pushed away first if necessary. The patient must be kept upright or even leaning a little forwards before the fluid can drain away; he should also lie on his face for a short time at intervals during the day. A soft rubber tube

may be guided into the lower end of the oblique sinus past the termination of the inferior cava, and another one towards the base of the heart on its right side. The edges of the pericardial wound should be sewn to the parietes whenever possible to try to avoid suppurative mediastinitis. Irrigation at the time of the operation is only necessary if masses of lymph or false membranes can be felt by the finger, and care must be taken to use two tubes side by side to ensure a free exit for the fluid, which may otherwise compress the heart; Mr. Parker believed this was the cause of death in his case. Daily irrigations may be used in the after-treatment. About 60 cases of operations for suppurative pericarditis have been recorded, with a mortality of about 65 per cent.; the real mortality is certain to be higher than this, for it is no doubt true that many unsuccessful operations have not been recorded.

REFERENCES.

¹ BRITISH MEDICAL JOURNAL, 1903, I, pp. 361. ² Ibid., 1903, II, pp. 359.
³ Trans. Amer. Surg. Assoc. ⁴ Lancet, June, 1900.

NOTE ON THE METHOD OF DEMONSTRATING TUBERCLE BACILLI IN THE URINE.

By EDGAR TREVITHICK, M.D. CAMB.,
Physician, Cheltenham General Hospital.

I have heard it remarked by observers that tubercle bacilli, when occurring in urine, are usually only to be found in very scanty numbers, and are to be considered as being somewhat difficult of demonstration. My own experience is opposed to this. And I am led to believe that the secret of finding representative numbers of bacilli in film preparations made from urinary deposits depends upon the thorough washing of such deposits with distilled water before the films are made from them. I have lately compared films made from urinary deposit after being thus washed with films made from the same deposit without this precaution being taken, and I find that in the latter very many fewer bacilli are to be found. The explanation of this appears to be that the urinary salts prevent the bacilli and even the pus cells from adhering sufficiently well to the cover slip to withstand the subsequent treatment necessitated in the process of staining, and that in reality in some parts of this process many, and I believe the majority of them, get dislodged and washed away.

The method which I have found eminently successful is as follows. The lower part of the urine which has been resting in a conical glass is transferred to the receiver of the centrifugal machine. After centrifugalizing, the supernatant fluid is carefully decanted from the minute collection of deposit which will be found adhering to the bottom of the glass. The glass is then filled with distilled water, and the centrifugate shaken up with it. This is once more centrifugalized, and after that the foregoing process is once more repeated. From the final centrifugate the films are prepared. In my experience the number of tubercle bacilli found in such films will be very greatly in excess of those that will be demonstrated from the same urine where this precaution of washing away the urinary salts is not followed.

In my experience the demonstration of tubercle bacilli in tuberculous urine is not more difficult, but on the contrary more simple and certain than their demonstration when occurring in sputum, and I find, moreover, that in the urinary films their numbers are not less, but on the contrary much greater, than is the case with films made from tuberculous sputum.

This is not surprising, seeing that in the case of urine one is able to take advantage of the concentration brought about by the process of centrifugalizing, an advantage which is not easily possible in dealing with sputum.

REACTION WITH PERCHLORIDE OF IRON IN THE URINE OF GRAVE CASES OF HEPATIC CIRRHOSIS.

By F. PARKES WEBER, M.D., F.R.C.P.,
Physician, German Hospital.

I HAVE lately been struck by the presence of a strong perchloride of iron reaction in the urine of several patients with hepatic cirrhosis. On adding a weak solution of perchloride of iron to the urine the colour obtained was sometimes similar to that obtained from the presence of diacetic acid in

cases of diabetes mellitus, but sometimes it appeared to me more like that obtained in the urine of patients taking sodium salicylate, so that I should have expected that some drug like aspirin had been taken. In only one of the cases was there any history of there ever having been any glycosuria. The appearance of the reaction sometimes coincided with very threatening general symptoms, such as drowsiness in Case I. I unfortunately delayed exact analysis, and at present can obtain no urine giving the reaction. I therefore am unable to say that the reaction is due to the presence of diacetic acid in the urine, and must content myself by giving a few clinical details of the cases in which the reaction was present, hoping that it will be detected in other cases of hepatic cirrhosis when its nature can be fully investigated.

CASE I.—Mrs. T. B., aged 42, admitted to the German Hospital June 20th, 1903. A fat woman, with slight jaundice, ascites, and epigastric pains. No cardiac or renal disease. No vomiting. Menses absent last two months. History of free use of beer and whisky during last five years. Ascites commenced apparently about three months before admission. Soon after admission the patient became drowsy and more jaundiced, and her pulse was very weak. She was treated by magnesium sulphate, digitalis, camphor hypodermics, and the temporary use of alcohol, and her general condition soon improved. A strong perchloride of iron reaction was first noted on July 2nd (the first time the urine was tested in that way), and the reaction was present after that date whenever the urine was tested. Paracentesis abdominis was twice performed. The patient left the hospital without permission on July 10th, and died not long afterwards.

CASE II.—Mrs. E. D., aged 53, admitted October 5th, 1903. A very sallow woman, with uniform great enlargement of the liver, but without ascites. History of taking whisky and beer. Enlargement of the abdomen noted two years before admission. Brought up blood three weeks before admission. The urine was free from albumen and sugar, and on many occasions gave a strong perchloride of iron reaction, though it had been rendered alkaline by giving the patient sodium bicarbonate. The patient left the hospital on November 17th. On December 12th, when in her own home, she became unconscious, and died on the following day. No necropsy.

CASE III.—J. A., aged 45, a carter, admitted September 22nd, 1903, for ascites, apparently of recent onset. History of habitual indulgence in beer and whisky. After paracentesis abdominis the liver could be felt enlarged. The urine was free from albumen and sugar, but dark, and at one time showed a distinct urobilin band by spectroscopic examination. On several occasions it gave a marked perchloride of iron reaction, but this has been invariably absent of late, though diligently searched for. The patient has been tapped for ascites six times, and is still in the hospital.

CASE IV.—Mrs. A. B., aged 53, admitted November 13th, 1903, on account of ascites, vomiting, and great frequency of cardiac action. The patient, a fat woman with a bloated face, used to drink much ale and whisky. About seven years before admission she was told she had "diabetes"; this disappeared after some time, and she has generally enjoyed good health. Her abdomen began to swell about six months before admission, and before her admission it had already been tapped twice at another hospital. The urine was free from albumen and sugar, but, although often alkaline, usually gave a marked perchloride of iron reaction. Her general condition has improved, she has had no vomiting lately, and her appetite is better; recently the perchloride of iron reaction has been absent. She has been tapped twice since admission, and is still an in-patient.

A METHOD FOR ANASTOMOSING A SEVERED VAS DEFERENS.

By J. LYNN THOMAS, C.B., F.R.C.S.,

Surgeon, Cardiff Infirmary; Consulting Surgeon, Hamadryad Hospital, Porth Hospital, Bridgend Hospital, and the Cardiff Provident Dispensary.

ABOUT twelve months ago I had under my care a gentleman who had his vasa deferentia tampered with, one being completely divided, and the other one incompletely severed by his wife during a fit of jealousy. I was consulted in the matter about a week after the somewhat tragical episode had taken place, and I found two scars in the scrotum anteriorly, one on each lateral aspect. I was informed that they had been performed by the patient's wife by means of a surgical knife whilst the patient was in a state of advanced semicomatose intoxication. On the side where the vas deferens had been completely divided there was a rounded hard swelling in the spermatic cord about 1 in. above the testis of the size of a big pea; the testis itself being more sensitive than the one on the other side. I was asked if anything could be done in the case, as the participants in the tragedy had become very much concerned as to the result with regard to sexual impotence, for the effect of double vasectomy was