may be guided into the lower end of the oblique sinus past the termination of the inferior cava, and another one towards the base of the heart on its right side. The edges of the pericardial wound should be sewn to the parietes whenever possible to try to avoid suppurative mediastinitis. Irrigation at the time of the operation is only necessary if masses of lymph or false membranes can be felt by the finger, and care must be taken to use two tubes side by side to ensure a free exit for the fluid, which may otherwise compress the heart; Mr. Parker believed this was the cause of death in his case. Daily irrigations may be used in the after-treatment. About 60 cases of operations for suppurative pericarditis have been recorded, with a mortality of about 65 per cent.; the real mortality is certain to be higher than this, for it is no doubt true that many unsuccessful operations have not been recorded.

REFERENCES.

¹ BRITISH MEDICAL JOURNAL, 1903, I, pp. 361. ² Ibid., 1903, II, pp. 359. ³ Trans. Amer. Surg. Assoc. ⁴ Lancet, June, 1900.

NOTE ON THE METHOD OF DEMONSTRATING TUBERCLE BACILLI IN THE URINE.

By EDGAR TREVITHICK, M.D. CAMB.,
Physician, Cheltenham General Hospital.

I have heard it remarked by observers that tubercle bacilli, when occurring in urine, are usually only to be found in very scanty numbers, and are to be considered as being somewhat difficult of demonstration. My own experience is opposed to this. And I am led to believe that the secret of finding representative numbers of bacilli in film preparations made from urinary deposits depends upon the thorough washing of such deposits with distilled water before the films are made from them. I have lately compared films made from urinary deposit after being thus washed with films made from the same deposit without this precaution being taken, and I find that in the latter very many fewer bacilli are to be found. The explanation of this appears to be that the urinary salts prevent the bacilli and even the pus cells from adhering sufficiently well to the cover slip to withstand the subsequent treatment necessitated in the process of staining, and that in reality in some parts of this process many, and I believe the majority of them, get dislodged and washed away.

The method which I have found eminently successful is as follows. The lower part of the urine which has been resting in a conical glass is transferred to the receiver of the centrifugal machine. After centrifugalizing, the supernatant fluid is carefully decanted from the minute collection of deposit which will be found adhering to the bottom of the glass. The glass is then filled with distilled water, and the centrifugate shaken up with it. This is once more centrifugalized, and after that the foregoing process is once more repeated. From the final centrifugate the films are prepared. In my experience the number of tubercle bacilli found in such films will be very greatly in excess of those that will be demonstrated from the same urine where this precaution of washing away the urinary salts is not followed.

In my experience the demonstration of tubercle bacilli in tuberculous urine is not more difficult, but on the contrary more simple and certain than their demonstration when occurring in sputum, and I find, moreover, that in the urinary films their numbers are not less, but on the contrary much greater, than is the case with films made from tuberculous sputum.

This is not surprising, seeing that in the case of urine one is able to take advantage of the concentration brought about by the process of centrifugalizing, an advantage which is not easily possible in dealing with sputum.

REACTION WITH PERCHLORIDE OF IRON IN THE URINE OF GRAVE CASES OF HEPATIC CIRRHOSIS.

By F. PARKES WEBER, M.D., F.R.C.P.,
Physician, German Hospital.

I HAVE lately been struck by the presence of a strong perchloride of iron reaction in the urine of several patients with hepatic cirrhosis. On adding a weak solution of perchloride of iron to the urine the colour obtained was sometimes similar to that obtained from the presence of diacetic acid in

cases of diabetes mellitus, but sometimes it appeared to me more like that obtained in the urine of patients taking sodium salicylate, so that I should have expected that some drug like aspirin had been taken. In only one of the cases was there any history of there ever having been any glycosuria. The appearance of the reaction sometimes coincided with very threatening general symptoms, such as drowsiness in Case I. I unfortunately delayed exact analysis, and at present can obtain no urine giving the reaction. I therefore am unable to say that the reaction is due to the presence of diacetic acid in the urine, and must content myself by giving a few clinical details of the cases in which the reaction was present, hoping that it will be detected in other cases of hepatic cirrhosis when its nature can be fully investigated.

CASE I.—Mrs. T. B., aged 42, admitted to the German Hospital June 20th, 1903. A fat woman, with slight jaundice, ascites, and epigastric pains. No cardiac or renal disease. No vomiting. Menses absent last two months. History of free use of beer and whisky during last five years. Ascites commenced apparently about three months before admission. Soon after admission the patient became drowsy and more jaundiced, and her pulse was very weak. She was treated by magnesium sulphate, digitalis, camphor hypodermics, and the temporary use of alcohol, and her general condition soon improved. A strong perchloride of iron reaction was first noted on July 2nd (the first time the urine was tested in that way), and the reaction was present after that date whenever the urine was tested. Paracentesis abdominis was twice performed. The patient left the hospital without permission on July 10th, and died not long afterwards.

CASE II.—Mrs. E. D., aged 53, admitted October 5th, 1903. A very sallow woman, with uniform great enlargement of the liver, but without ascites. History of taking whisky and beer. Enlargement of the abdomen noted two years before admission. Brought up blood three weeks before admission. The urine was free from albumen and sugar, and on many occasions gave a strong perchloride of iron reaction, though it had been rendered alkaline by giving the patient sodium bicarbonate. The patient left the hospital on November 17th. On December 12th, when in her own home, she became unconscious, and died on the following day. No necropsy.

CASE III.—J. A., aged 45, a carter, admitted September 22nd, 1903, for ascites, apparently of recent onset. History of habitual indulgence in beer and whisky. After paracentesis abdominis the liver could be felt enlarged. The urine was free from albumen and sugar, but dark, and at one time showed a distinct urobilin band by spectroscopic examination. On several occasions it gave a marked perchloride of iron reaction, but this has been invariably absent of late, though diligently searched for. The patient has been tapped for ascites six times, and is still in the hospital.

CASE IV.—Mrs. A. B., aged 53, admitted November 13th, 1903, on account of ascites, vomiting, and great frequency of cardiac action. The patient, a fat woman with a bloated face, used to drink much ale and whisky. About seven years before admission she was told she had "diabetes"; this disappeared after some time, and she has generally enjoyed good health. Her abdomen began to swell about six months before admission, and before her admission it had already been tapped twice at another hospital. The urine was free from albumen and sugar, but, although often alkaline, usually gave a marked perchloride of iron reaction. Her general condition has improved, she has had no vomiting lately, and her appetite is better; recently the perchloride of iron reaction has been absent. She has been tapped twice since admission, and is still an in-patient.

A METHOD FOR ANASTOMOSING A SEVERED VAS DEFERENS.

By J. LYNN THOMAS, C.B., F.R.C.S.,

Surgeon, Cardiff Infirmary; Consulting Surgeon, Hamadryad Hospital, Porth Hospital, Bridgend Hospital, and the Cardiff Provident Dispensary.

ABOUT twelve months ago I had under my care a gentleman who had his vasa deferentia tampered with, one being completely divided, and the other one incompletely severed by his wife during a fit of jealousy. I was consulted in the matter about a week after the somewhat tragical episode had taken place, and I found two scars in the scrotum anteriorly, one on each lateral aspect. I was informed that they had been performed by the patient's wife by means of a surgical knife whilst the patient was in a state of advanced semicomatose intoxication. On the side where the vas deferens had been completely divided there was a rounded hard swelling in the spermatic cord about 1 in. above the testis of the size of a big pea; the testis itself being more sensitive than the one on the other side. I was asked if anything could be done in the case, as the participants in the tragedy had become very much concerned as to the result with regard to sexual impotence, for the effect of double vasectomy was