seemed to be superior to electrocardiography, clinical examination, and the newer creatine kinase MB mass assays.^{1,2} These trials confirm that troponin T is a powerful and independent predictor of an adverse outcome and that this predictive function extends well beyond admission to hospital.² The prognostic importance for the cohort with a final diagnosis of myocardial infarction has been addressed elsewhere.³

Patients were retrospectively given their final diagnostic classification when all investigational data were available. All were initially admitted with a suspected acute coronary syndrome. Patients classified as having unstable angina were required to have evidence of coronary artery disease—that is, either a subsequent cardiac event or results of investigations indicating such disease, as described in the paper. Patients classified as having non-ischaemic chest pain had negative results of exercise tests or normal coronary angiograms or thallium scans and underwent further investigations, including ventilation-perfusion lung scanning and gastro-duodenoscopy, before being given their final diagnostic classification. None of this cohort had any cardiac event on follow up. We therefore believe that this hierarchical diagnostic categorisation protocol allows a clearer definition of unstable angina in our population than the heterogeneous definitions used by other study groups. With the development of a rapid assay and the next generation of assays for troponin T (which are more sensitive and cardiospecific),⁴ this marker can now be measured more widely.

While we hold the view that measurement of troponin T is a useful addition to current practice in stratifying the risk for patients admitted with acute coronary syndromes, we concur with Kennedy's view that it should not be used as the sole discriminant of risk.

PETER STUBBS

Lecturer in cardiovascular medicine

MARK NOBLE

Weston professor of cardiovascular medicine

Charing Cross and Westminster Medical School,
London W6 8RP

PAUL COLLINSON

Consultant chemical pathologist

Mayday University Hospital NHS Trust,
Croydon CR4 7YE

DAVID MOSELEY

Senior medical laboratory scientific officer

TREVOR GREENWOOD

Consultant cardiologist

West Middlesex University NHS Trust,
Isleworth TW7 6AF

1 Ohman EM, Armstrong P, Christenson R, Granger C, Katus H, Hamm C, *et al*. Cardiac troponin T levels for risk stratification in acute myocardial ischemia. *N Engl J Med* 1996;335:1333-41.

2 Lindahl B, Toss H, Venge P, Wallentin L for the FRISC Study Group. Relation between troponin T and the risk of subsequent cardiac events in unstable coronary artery disease. *Circulation* 1996;93:1651-7.

3 Stubbs P, Collinson P, Moseley D, Greenwood T, Noble M. The prognostic significance of admission troponin T concentrations in myocardial infarction. *Circulation* 1996;94:1291-7.

4 Collinson P, Thomas S, Siu L, Vasudeva P, Stubbs P, Canepa-Anson R. Rapid troponin T measurement in whole blood for detection of myocardial damage. *Ann Clin Biochem* 1995;32:454-8.

for the world.² The "revolutionary" ideas of the primary health care strategy included an emphasis on intersectoral action for health, reallocation of health resources from tertiary care to community based health promotion and disease control, participation by the community, and scientifically validated interventions. These are now mainstream concepts and are being increasingly applied in developed and developing countries.

In terms of increases in life expectancy, decreases in infant mortality and in the prevalence of many infectious diseases, and improved access to environmental health and basic maternal and childhood preventive and promotional services, there have been substantial global health gains since the conference in Alma-Ata.³ Nevertheless, along with these gains the WHO has documented numerous obstacles, such as increased inequality in health and wealth in many countries.

As the global health community prepares for the 21st century the WHO is spearheading a global consultation aimed at renewing the concept of health for all. This broad based process aims to update the old policy and add components that address the present and future context of health development. The WHO believes that health for all remains a powerful vision for the 21st century. Yet we are not naive; we realise that certain serious challenges threaten future progress, and in this respect we welcome the authors' identification of macroeconomic factors that influence health. These same factors have been repeatedly emphasised by the WHO's task force on health and development,⁴ of which Mocumbi is a member.

The WHO is realistic, however, about its direct role in alleviating poverty. We will continue to document the root causes of ill health and poverty. We will continue to advocate development policies that reduce inequalities in wealth, build long term human and institutional capacity, use intersectoral approaches, and encourage sustainable development. Beyond advocacy, the WHO supports actions within the health sector that directly reduce the burden of disease and thereby help to improve the scholastic performance of children, increase the productivity of workers, and alleviate human suffering. These direct inputs, when combined with basic education, can accelerate populations' escape from poverty.

F S ANTEZANA

Assistant director general

World Health Organisation,
CH-1211 Geneva,
27 Switzerland

1 Bergström S, Mocumbi P. Health for all by the year 2000? *BMJ* 1996;313:316. (10 August.)

2 World Health Organisation. *Global strategy for health for all by the year 2000*. Geneva: WHO, 1981.

3 World Health Organisation. *The world health report, 1995: bridging the gaps*. Geneva: WHO, 1995.

4 World Health Organisation. *Health in development: prospects for the 21st century. Report of the first meeting of the task force on health in development*. Geneva: WHO, 1994.

Much can still be done

EDITOR,—Staffan Bergström and Pascoal Mocumbi provide a telling description of the health burdens now confronting developing populations.¹ Jordan affords a glaring example of the "debt trap" depicted by the authors: "Pressured by the International Monetary Fund, the Jordanian government raised the price of bread from 12 [US] cents [7p] to 26 cents per kg, and of more than 20 other basic food items."² In Uganda the equivalent of \$3 (£1.86) a head is being spent annually on total health services, compared with \$17 (£10.56) required for the repayment of debt.³ In South Africa the situation is less acute. From figures given in parliament, for every 100 South African (SA) cents (26 US cents, 16p) that the government spends, 18.7 SA cents is required for redemption of debt, while

21.2 SA cents is spent on education, 9.9 SA cents on health, and 9.6 SA cents on welfare.

Although the authors maintain that "health for all is a distant dream," some positive changes are occurring.⁴ Thus in all African countries infant mortality has fallen considerably. In South Africa infant mortality among Africans in big cities is 20-25/100 000. While this is triple that among the white population, it compares favourably with that among Afro-Americans (17/100 000); furthermore, rates in minority populations, such as New Zealand Maoris and Australian Aborigines, are of the same order.

Despite the fearful outlook for developing populations, improvements in health, wellbeing, and community spirit can still be made. The involvement of women must certainly be supported. In this respect, governments' increasing willingness to address women's rights and needs in policy and programmes is encouraging. The previous director general of the World Health Organisation, Dr Halfden Mahler, often emphasised women's important role in health promotion—their role in delivering babies, as volunteers in hospitals, in self help clinics, and in other community organisations and practices. Skilled nurses could be given greater responsibilities in rural hospitals; with training they could conceivably undertake emergency surgery. More mutual help in the community is also effective, as exemplified by the outstanding experience of Kerala, one of the poorest states in India.⁵ Whereas Kerala's infant mortality is about 30/100 000, that of India as a whole is 40-50/100 000. Additionally, greater use must be made not only of health workers based in the community but of traditional healers.

While health for all is a dream for African countries, there is much that is positive and much that can still be done, if not to better then at least to restrain the inevitably worsening economic and health scenarios.

ALEXANDER R P WALKER
Head

Human Biochemistry Research Unit,
Department of Tropical Diseases,
South African Institute for Medical Research,
Johannesburg,
South Africa

LESLEY T BOURNE
Researcher

Urbanisation and Health,
Medical Research Council,
Cape Town,
South Africa

BARBARA J KLUGMAN
Researcher

Women's Health Project,
Department of Community Health,
University of the Witwatersrand,
Johannesburg,
South Africa

1 Bergström S, Mocumbi P. Health for all by the year 2000? *BMJ* 1996;313:316. (10 August.)

2 Talk of the streets. Give us this day our daily bread. *Time* 1996;148:12.

3 Benatar SR. The World Bank, listening and learning. *Lancet* 1996;347:1047.

4 Walker ARP, Walker BF, Glatthaar II. Impoverished Africa: any hope for the future? *J R Soc Med* 1994;87:157-60.

5 Kutty VR. Women's education and its influence on attitude to aspects of child-care in a village community in Kerala. *Soc Sci Med* 1989;29:1299-303.

Health for all by the year 2000

WHO is spearheading global consultation to update policy on health for all

EDITOR,—Staffan Bergström and Pascoal Mocumbi highlight several impediments to the attainment of health for all.¹ In so doing they provide a too pessimistic view of past successes and future opportunities.

At the World Health Organisation/Unicef conference in Alma-Ata in 1978 the concept of health for all was accepted as a motivational vision

Code of practice on managing tuberculosis is satisfactory

EDITOR,—Julia E Clark and Andrew J Cant report three cases of childhood tuberculosis and discuss their management in relation to the 1994 code of practice of the joint tuberculosis committee of the British Thoracic Society.^{1,2} They conclude that all children in contact with tuberculosis should have a chest x ray examination irrespective of the result of the initial tuberculin test. We do not believe that this recommendation is justified for the following