Tertiary cancer services

Psychiatric and psychological liaison should be included

EDITOR,—M A Richards and colleagues report a benchmarking study of activity and facilities at 12 specialist cancer centres.¹ One of the objectives of the study was to help the planning of future cancer services. The authors looked at inpatient and outpatient services and also at radiotherapy and chemotherapy practice. They make no mention, however, of psychological, psychiatric, or social input to patients and their families.

In the same issue of the *BMJ* Sandra Goldbeck-Wood discusses how tomorrow's doctors should be selected and quotes an appeal from patients for doctors who are, among other things, "broad minded," "informative," and "holistic."² The same issue also contains I G McDonald and colleagues' paper looking at how to reassure the worried well.³ So much for the worried well; what about the worried ill or dying?

I have a good friend who is currently a patient in an oncology ward. Watching her become deeply depressed and frightened has reminded me of the importance of considering the psychiatric and psychological aspects of care for a person with cancer. As she acquiesces to her latest course of chemotherapy she struggles with her own grief and that of her family. She feels no pleasure, only guilt and suspicion. Everything is negative in her world: all thoughts, all memories, all contact with others. Gone are the cares about her treatment. They have been superseded by her need to find some peace rather than feeling frightened and eternally damned.

Surely, somewhere in the grand plan for cancer services there should be some mention of psychiatric and psychological liaison. Megavolts and milligrams alone are not enough.

MOLI PAUL

Senior registrar in child and adolescent psychiatry
Shrewsbury Child and Family Service,
Shrewsbury SY3 9BL

- Richards MA, Parrott JC, on behalf of the 12 participating centres. Tertiary cancer services in Britain: benchmarking study of activity and facilities at 12 specialist centres. *BMJ* 1996;313:347-9. (10 August.)
- Goldbeck-Wood S. Choosing tomorrow's doctors. *BMJ* 1996;313:313. (10 August.)
- McDonald IG, Daly J, Jelinek VM, Panetta F, Gutman JM. Opening Pandora's box: the unpredictability of reassurance by a normal test result. *BMJ* 1996;313:329-32. (10 August.)

Use of "population served" distorted findings

EDITOR,—M A Richards and J C Parrott raise important issues about tertiary cancer services.¹ Sadly, their paper does not contribute to our understanding of benchmarking for tertiary cancer centres because the calculation of population denominators is seriously flawed. The standard method of calculating catchment populations for specific services uses the proportions of current activity provided by different providers to allocate a district's resident population to those

providers.² The key principle is that the catchment populations of all providers add up to the total populations of the districts considered.

Instead, Richards and Parrott use the concept of "population served," which involves classifying districts as local or distant and exclusive or shared and then allocating the populations of shared districts by relating activity for them to activity for the host population.

This definition of population served seriously distorts the findings.

Visualise three districts, X, Y, and Z. Each has a population of one million. Cancer centres A and B are located in districts X and Z, respectively. All the activity for the population of X and Z takes place at A and B, respectively, while half of the activity for the population of Y takes place at A and half at B. Under these conditions, whatever the actual levels of activity for the populations of X, Y, and Z, the catchment populations of A and B are each 1.5 million. In contrast, the population served by A and B, as defined in the paper, depends on levels of activity for each population. Table 1 shows this, with four scenarios for activity. Only if the activity for each district is the same will the figure for the population served be the same as that for the catchment population. In scenario 2 the total population served is four million when the total population of the three districts is only three million. Similarly, in scenario 3 (which is possibly more likely) the total population served is only 2.5 million. In scenario 4 the centre that provides most activity looks as though it serves a smaller population.

This is not simply an arithmetical quibble. Multiplying the rate of new referrals contrived from the authors' method by a local population could seem to indicate, out of the blue, a need for more activity. While this may be the situation in many districts, this method will not elucidate which ones to target.

BARRY TENNISON

Director of public health

ALISON FRATER

Assistant director of public health

West Hertfordshire Health Authority,
Welwyn Garden City AL8 6JL

KEITH BAKER

Assistant director of public health

East and North Hertfordshire Health Authority,
Welwyn Garden City AL8 6JL

- Richards MA, Parrott JC, on behalf of the 12 participating centres. Tertiary cancer services in Britain: benchmarking study of activity and facilities at 12 specialist centres. *BMJ* 1996;313:347-9. (10 August.)
- Department of Health. *Health service indicators guidance: dictionary*. London: HMSO, 1989. (Appendix B.)

Authors' reply

EDITOR,—We share Moli Paul's view of the importance of psychosocial care for patients with cancer, which has received specific attention in recent national reports.^{1 2} Our paper, however, was focused on facilities and activity levels at cancer centres. Psychosocial care depends critically on the provision of adequate numbers of appropriately trained staff. As we indicated,

Fig 1—Activity among 12 cancer centres, based on effective population

we are undertaking further work in relation to staffing. This includes benchmarking of staffing levels for nurse specialists, psychiatrists, psychologists, and palliative care specialists. The results will be reported when they have been verified.

Barry Tennison and colleagues seem to have missed a crucial point about our study: it was limited to 12 centres to facilitate detailed discussion and validation of the data. While these centres are among the largest providers of tertiary cancer services in the United Kingdom, many more hospitals provide tertiary services, including radiotherapy. It was therefore impossible to calculate catchment populations in the way that Tennison and colleagues describe because some patients from "distant shared" districts were referred to centres outside those in our analysis. Even if all tertiary providers had participated in this study, different methods of counting cancer activity among centres would have limited the application of Tennison and colleagues' method.

Our methodology was designed to give the best possible estimate of the proportion of a distant shared district's population that was served by the centres in the study. We refute the implication that our methodology "seriously distorts" our findings as suggested by the extreme figures that Tennison and colleagues have chosen to illustrate their argument. Comparison of activity levels for host districts and "distant exclusive" districts for individual centres in our study indicates that wide variations are restricted to remote areas that, for geographical reasons, have poor access to any centre.

At the outset of the study each centre provided an initial estimate of the population served. The total figure for all 12 centres was 25.7 million. After our detailed analysis on a district by district basis the figure was 24.3 million (a 5% change). The benchmarking analyses were undertaken with both sets of population figures. Marginal changes in the rankings of individual centres were observed, but the principal finding of a wide variation in facilities and activity among the 12 centres was unchanged (fig 1). Had we adopted the initial population estimates, the proportion of patients with cancer estimated as being referred to non-surgical oncologists (40-45%) would have been marginally lower. The increase in the administration of chemo-

Table 1—Population served in four possible scenarios of activity at two cancer centres, assuming that all activity for populations of X and Z takes place at A and B, respectively, and that half of activity for population of Y takes place at A and half at B

Scenario	Activity for population of:			Population (million) served by:	
	X	Y	Z	A	B
1	100	100	100	1.5	1.5
2	100	200	100	2	2
3	200	100	200	1.25	1.25
4	400	100	200	1.125	1.25