- 1 Thorne C. *Better medical writing*. London: Pitman Medical, 1970.
- 2 Fulton RA. Goals and methods of audit should be reappraised. *BMJ* 1996;312:1103. (27 April.)
- 3 Miles A, Bentley P, Polychronis A, Price N, Grey J. Goals and methods of audit should be reappraised. *BMJ* 1996;313:497. (24 August.)
- 4 Teasdale S. The future of clinical audit: learning to work together. *BMJ* 1996;313:574. (7 September.)
- 5 Lough JRM. Goals and methods of audit should be reappraised. *BMJ* 1996;313:497. (24 August.)

Evidence of effectiveness of audit in improving care is growing

EDITOR,—Andrew Miles and colleagues' letter about audit exemplifies the confusion in the debate about the achievements of the national audit programme.¹ There are misunderstandings about the true nature of audit among both clinicians and researchers, and this results in opinions that are poorly supported by evidence. In evaluating the impact of audit it is essential to distinguish between audit as a method and the way in which it has been implemented. Although only limited evidence exists about the effectiveness of the audit programme, there is growing evidence of the potential and actual effectiveness of audit in improving standards of care.

Audit consists of a series of steps, beginning with the selection of a topic, followed by the identification of explicit, evidence based, and prioritised criteria against which to judge performance as determined by the collection of valid and reliable data. Discrepancies between performance and criteria are addressed by selected strategies, which may include feedback, facilitation, and reminders, after which data are collected again to check progress.² Activities that comply with this definition bring about improvements in care³; activities that do not comply with this definition often involve aimless collection of data, resulting in the disillusionment and demotivation of participants because they can see no benefit.

Given that audit as a method can be effective, how effective has the audit programme been in applying it? Because concerns have been expressed, it would be wise to investigate the programme in different settings. It cannot be assumed that, because some so called audit meetings for hospital specialists seem to have little impact, audit by general practitioners, physiotherapists, nurses, or hospital specialists who adopt more appropriate methods is also without impact. Participation in audit in general practice is high,⁴ and a growing number of examples are leading to change.⁵ To increase the achievements of the audit programme, clinicians' understanding of audit methods must be improved, and those aspects of the programme that are most effective should replace those that have been less effective.

In common with many colleagues in the NHS in both primary care and beyond, we take a much more positive view of the past achievements and the future potential of audit than do Miles and colleagues.

RICHARD BAKER
Director, Eli Lilly national clinical audit centre
ROBIN C FRASER
Professor of general practice

Department of General Practice and Primary Health Care,
University of Leicester,
Leicester General Hospital,
Leicester LE5 4PW

- 1 Miles A, Bentley P, Polychronis A, Price N, Grey J. Goals and methods of audit should be reappraised. *BMJ* 1996;313:497. (24 August.)
- 2 Lawrence M. What is medical audit? In: Lawrence M, Schofield T, eds. *Medical audit in primary health care*. Oxford: Oxford University Press, 1993:3-13.
- 3 Davis DA, Thomson MA, Oxman AD, Haynes RB. Changing physician performance. A systematic review of the effect of continuing medical education strategies. *JAMA* 1995;274:700-5.

- 4 Baker R, Hearnshaw H, Cooper A, Cheater F, Robertson N. Assessing the work of medical audit advisory groups in promoting audit in general practice. *Quality in Health Care* 1995;4:234-9.
- 5 Fraser RC, Farooqi A, Sorrie R. Use of vitamin B-12 in Leicestershire practices: single topic audit led by medical audit advisory group. *BMJ* 1995;311:28-30.

Blood donated after vaccination with rabies vaccine derived from sheep brain cells might transmit CJD

EDITOR,—Janet Morgan reports that the National Blood Authority in Britain has decided to tighten the donor screening programme to exclude transmission of Creutzfeldt-Jakob disease or its variant through blood donations.¹ Prospective donors will be prevented from donating blood if they have a history of treatment with human growth hormone or if one of their siblings, parents, or grandparents developed the disease. I would point out that similar care should also be taken when immigrants from Asia and Africa offer to donate blood, in case they received rabies vaccine derived from culture of sheep brain cells when they were living in their country of origin.

In many countries in Asia and Africa limited supplies of imported rabies vaccines derived from culture of human cells have been available. Many people continue to be offered indigenously produced sheep brain vaccine after exposure to a rabid animal. Scrapie is known to exist in sheep around many centres where the vaccine is produced. In the mountain sheep of the Kumaon foothills in the Himalayas, for example, scrapie was established more than four decades ago and 1-10% of the flock was reported to have the disease in 1961.² In the Himalayan foothills the Central Research Institute continues to produce four to five million doses of sheep brain vaccine annually. Transmission of abnormal prion protein, PrP^{Sc}, in sheep brain vaccine might have occurred in some of the 30 documented cases of Creutzfeldt-Jakob disease in different regions in India.³ Because Creutzfeldt-Jakob disease has a latency of about 20 years, many recipients of sheep brain rabies vaccine could emigrate to Britain before becoming ill.

Before accepting blood donations from immigrants it would be desirable to ask the potential donors whether they were exposed to a rabid animal and immunised with sheep brain rabies vaccine in their country of origin. Furthermore, indirect assessment should be possible through, for example, assay looking for antibodies specific to rabies.

SUBHASH C ARYA
Clinical microbiologist

Centre for Logistical Research and Innovation,
M-122 (of part 2), Greater Kailash-II,
New Delhi-110048,
India

- 1 Morgan J. Blood to be screened for CJD. *BMJ* 1996;313:441. (24 August.)
- 2 Zlotnik I, Katiyar RD. The occurrence of scrapie disease in sheep of the remote Himalayan foothills. *Vet Rec* 1961;73:543-4.
- 3 Satishchandra P, Shankar SK. Creutzfeldt-Jakob disease in India (1971-1990). *Neuroepidemiology* 1991;10:27-32.

Royal Medical Benevolent Fund's Christmas appeal

EDITOR,—As one of the great festivals of the year, Christmas affects everyone in Britain irrespective of their race or creed. Above all, it means so much to children. Most of us will enjoy this traditional occasion: it is a time for Christmas cards, gifts, and a special dinner. Some people, however, because of poverty, will not be able to

celebrate Christmas as they would like for themselves or for their children.

The beneficiaries of the Royal Medical Benevolent Fund range from very young children to elderly people. Their need for help is carefully assessed, and only those in real need are eligible. At Christmas the fund sends something to each beneficiary. Children receive special gifts. The fund receives many letters of thanks, which show how much the gifts and cards are appreciated.

This will be my last Christmas appeal, as I believe that it is now time for someone younger to take over. I have appreciated the generous response to the appeal over the years and hope that the 1996 appeal will equal or exceed its predecessors. Contributions marked "Christmas appeal" may be sent to the secretary, Royal Medical Benevolent Fund, 24 King's Road, London SW19 8QN or to the treasurer or medical representative of any local guild of the fund.

JOSEPHINE BARNES
President

Royal Medical Benevolent Fund,
London SW19 8QN

Race may be hard to define

EDITOR,—One can classify pretty well anything as long as one has a clear idea of the purpose and axes of the classification: books, plants, race-horses, diseases—even humans.¹ If educational planners were classifying humans they might choose first language as the axis; health planners might choose socioeconomic class; housing authorities might choose immigrant status.

Who would need to classify humans by race, though, and how would they do it, given our propensity to travel the globe and breed where we rest? One of England's greatest journalists summed up the difficulty of defining his own race 300 years ago²:

The Romans first with Julius Caesar came,
Including all the Nations of that Name,
Gauls, Greeks, and Lombards; and by
Computation,
Auxiliaries or Slaves of ev'ry Nation.
With Hengist, Saxons; Danes with Sueno came,
In search of Plunder, not in search of Fame.
Scots, Picts, and Irish from th' Hibernian Shore:
And Conqu'ring William brought the
Normans o're.
All these their Barb'rous Offspring left behind,
The Dregs of Armies, they of all Mankind;
Blended with Britains who before were here,
Of whom the Welsh ha' blest the Character.
From this Amphibious Ill-born Mob began
That vain ill-nature'd thing, an Englishman.

CAROL ORCHARD
Senior researcher

National Casemix Office,
Winchester SO22 5DH

- 1 Describing race, ethnicity, and culture in medical research [letters]. *BMJ* 1996;313:425. (17 August.)
- 2 Defoe D. The English race, from *The True-Born Englishman* (1701). In: *The Oxford book of eighteenth century verse*. Oxford: Oxford University Press, 1926.

Correction

QTc dispersion and risk of cardiac death in peripheral vascular disease

An authors' error occurred in the last letter in this cluster, by Dawood Darbar and Allan D Struthers (26 October, pp 1082-3). The values given for QT dispersion in the penultimate sentence in the second paragraph were incorrect, and the sentence should have read: "QT dispersion and heart rate were not reported due to limitation of space; QT dispersion was significantly prolonged in those dying of cardiac causes compared with those who survived (110 (70) v 50 (20) ms respectively, P<0.001)."