Fig 1—Proportion of editorials devoted to medicine, administration, and other topics in first four months' issues of BMJ at five year intervals, 1958-93

latter figure being the highest in the previous 35 years. The proportion of editorials about medical matters fell from a peak of 77% (86/113) to 49% (47/96) over the same period (fig 1).

This audit shows a shift in the subject matter covered by editorials in the *BMJ* since 1978, from medical to administrative topics. The audit covered only the editorials, and we are aware that the *BMJ* has taken initiatives to increase its coverage of medical topics—for example, it has introduced the ABC and Recent Advances series. Scheckler, however, advised that when one reads a journal one should always read the editorials first.³ It is important that the *BMJ* brings the effects of any changes in NHS administration to readers' attention, but it must surely be a matter for concern when this results in fewer editorials on medical topics. We have not yet assessed whether this trend is occurring in other medical journals.

JONATHAN M GLASS
Registrar in urology
ANURAJ K NIGAM
Senior registrar

St Mary's Hospital,
London W2 1NY

Department of Urology,
St Bartholomew's Hospital,
London EC1A 7BE

- Currie BF. Continuing education from medical periodicals. *J Med Educ* 1976;51:420.
- Krogh CL. A checklist system for critical review of medical literature. *Med Educ* 1985;19:392-5.
- Scheckler WE. A realistic journal reading plan; the cornerstone of continuing medical education. *JAMA* 1982;248:1987-9.

Standardising crash call numbers

A minor issue compared with need to change concept of cardiac arrest team

EDITOR,—Salem A Beshyah and colleagues imply that the adoption of a national telephone number to summon emergency medical help might increase survival from cardiac arrests that occur in hospital.¹ It is disturbing that the authors focus on the side issue of how to summon help and do not address the problem of when to summon help.

At present the only systematic and universal response to serious illness in general wards occurs after the patient has almost died, when the cardiac arrest team is summoned and cardiopulmonary resuscitation started. Despite widespread awareness and teaching of both basic and advanced life support, however, the outcome in patients who have had an arrest has changed little since such teaching was introduced.² More depressing is the knowledge that up to three

fifths of cardiac arrests that occur in hospital have easily recognisable antecedent factors, such as hypotension, tachypnoea, and tachycardia.³ The conclusion must be that serious illness is not managed well in general wards.

If we are to improve survival from cardiac arrest appreciably then we must change the concept of the cardiac arrest team. Instead of having a reactive system that allows the patient to all but die before action is taken we must develop a proactive system that will be initiated according to physiological abnormalities such as tachypnoea, hypotension, and tachycardia; a sudden fall in level of consciousness; and potentially life threatening complications such as airway problems. Such a medical emergency team has been described⁴; it not only may prevent some cardiac arrests but will ensure that appropriate treatment is started while the patient has maximal physiological reserve to respond. This is of great importance because limited physiological reserve has been shown to be an important determinant of mortality.⁵ Additionally, inappropriate admissions to the intensive care unit may be avoided.

It would be also interesting to know which staff make the most telephone calls requesting the cardiac arrest team. From experience, we suspect that it is the nursing staff and not the rotating junior medical staff, who, if present, ought to be beginning resuscitation.

Although we agree with Beshyah and colleagues that a standard emergency telephone number would be appropriate, this is only a minor issue. No matter what telephone number is used, the current system must be changed to one that is activated before the patient is found to be pulseless.

GEORGE P FINDLAY
Research fellow in intensive care medicine
findlay@cardiff.ac.uk
NICK J STALLARD
Registrar in anaesthesia

University Hospital of Wales,
Cardiff CF4 4XW

- Beshyah SA, Schiff R, Al-Jehani FS. National crash call number is needed. *BMJ* 1996;313:429. (17 August.)
- McGrath RB. In-hospital cardiopulmonary resuscitation after a quarter of a century. *Ann Emerg Med* 1987;16:1365-8.
- Schein RMH, Hazday N, Pena M, Ruben BH, Sprung CL. Clinical antecedents to in-hospital cardiopulmonary arrest. *Chest* 1990;98:1388-92.
- Lee A, Bishop G, Hillman KM, Daffurn K. The medical emergency team. *Anaesth Intensive Care* 1995;23:183-6.
- Goldstein RL, Campion EW, Thibault GE, Mulley AG, Skinner E. Functional outcomes following medical intensive care. *Crit Care Med* 1986;14:783-8.

A recommendation already exists

EDITOR,—Salem A Beshyah and colleagues report their survey of switchboards of acute hospitals in the North Thames regions to establish the crash call numbers used.¹ A similar (unpublished) study that I carried out earlier this year in the Oxfordshire region gave similar results. As a result of my study I wrote to ask the chief medical officer to recommend standardisation of crash call numbers along the lines of the 999 emergency system. A reply from the NHS Estates Agency directed me to a publication on telephone services (one of the "health building notes").² In this document a paragraph deals with calls to the emergency services, and a recommendation is given that "the telephone number for emergency services should be easy to remember and easy to dial. Staff moving from one location to another will not be confused if health care premises use the same number, but this may not be practical in every case. The preferred number to call is 222 or 2222 as appropriate."

I understand that this advice is now being revised to recommend that the number 222 or 2222 be reserved for cardiac arrests and 333 or 3333 for all other emergencies (in Northern Ireland the preferred number for all emergencies is 6000 or 6666). It is hoped that the publication will be reissued by the end of this year.

Notwithstanding this advice, I am sorry that the chief medical officer did not take up my invitation to issue a recommendation to the profession; perhaps he believes that this little known health building note is adequate. I hope that all interested clinicians will now draw this recommendation to the attention of their estates managers and telephone switchboard managers so that we can accept Beshyah and colleagues' recommendations without further delay.

MICHAEL WARD
Consultant anaesthetist

Nuffield Department of Anaesthetics,
Oxford Radcliffe Hospital NHS Trust,
John Radcliffe Hospital,
Oxford OX3 9DU

- Beshyah SA, Schiff R, Al-Jehani FH. National crash call number is needed. *BMJ* 1996;313:429. (17 August.)
- NHS Estates, Department of Health, and the Welsh Office. Telephone services. London: DoH, 1989. (Health building note (HBN) 48.)

Goals and methods of audit should be reappraised

Audit as an essential part of summative assessment needs reconsideration

EDITOR,—Over 25 years ago Thorne wrote, "Magpie collection of statistics for no better reason than the accumulation of raw data followed by an aimless feeding of figures into a computer produces nothing of scientific value."¹ The recent critics of some audit activities echo Thorne's statement and express doubts about less than rigorous methods of audit design and analysis.^{2,3} Where an important subject has been audited and a robust method of follow through has been adopted, however, there are good examples of how quality of care can probably be improved.⁴

Summative assessment for registrars in general practice has led to an industry of standardised testing. Audit has been sucked into this and may be inadvertently stifling creativity. The desire for regional audit programmes for training practices⁵ has to be matched with a recognition that audit by itself has only a limited part to play in the overall development of young doctors. Attempts to compress audit into short time scales merely to satisfy regulations often result in the exercise being seen as a chore rather than a learning experience. Audit has its place but would be more acceptable to many young doctors if it was an optional, rather than a compulsory, means of showing competency to be a principal in general practice.

Summative assessment is a good idea in principle but depends on trusted and reliable methods backed up by adequate support systems. Because adequate support systems are in short supply, people at various different levels in my region are finding it increasingly difficult to justify the time and effort required for an ever increasing number of assessment procedures that seem to be more about quantity than quality. I recognise that those of us charged with responsibility in higher education have to try to help people learn new techniques, but the appropriate application of these techniques requires thought, time, adequate resources, and opportunities for feedback, all of which are at a premium. We also have a duty to ensure that expensive systems for collecting data do not expand too quickly before we become engulfed in Thorne's "meaningless collection of statistics."

JOHN BAIN
Professor

Tayside Centre for General Practice,
University of Dundee,
Dundee DD2 4AD