- 1 McKernan RO, Thomas DGT. The clinical study of gliomas. In: *Brain tumours: scientific basis, clinical investigation and current therapy*. In: Thomas DGT, Graham DI, eds. London: Butterworths, 1980:194-230.
- 2 Todd NV, McDonagh T, Miller JD. What follows the diagnosis by computed tomography of solitary brain tumour? Audit of one year's experience in south east Scotland. *Lancet* 1987;i:611-2.
- 3 Thomas DGT, Nouby RM. Experience in 300 cases of CT-guided stereotactic surgery for lesion biopsy and aspiration of haematoma. *Br J Neurosurg* 1984;3:321-6.
- 4 Walker MD, Alexander E, Hunt WE, MacCarty CS, Mahaley MS, Mealey J, et al. Evaluation of BCNU and/or radiotherapy in the treatment of anaplastic gliomas: a co-operative clinical trial. *J Neurosurg* 1978;49:333-43.
- 5 Walker MD, Green SB, Byar DP, Alexander E, Batzdorf U, Brooks WH, et al. Randomised comparisons of radiotherapy and nitrosureas for the treatment of malignant glioma after surgery. *N Engl J Med* 1980;303:1323-9.
- 6 Green SB, Byar DP, Walker MD, Pistenmaa DA, Alexander E, Batzdorf U, et al. Comparisons of carmustine, procarbazine, and high-dose methylprednisolone as additions to surgery and radiotherapy for the treatment of malignant glioma. *Cancer Treat Rep* 1983;67:121-32.
- 7 Gibberd FB, Scott GM. Morbidity in patients with intracranial gliomas. *J Neurol Neurosurg Psychiatry* 1983;46:460.
- 8 Wroe SJ, Foy PM, Shaw MDM, Williams IR, Chadwick DW, West C, et al. Differences between neurological and neurosurgical approaches in the management of malignant brain tumours. *BMJ* 1986;293:1015-8.
- 9 Choucair AK. Proposal for evaluation of toxic effects associated with treatment of gliomas: a call for action. *J Natl Cancer Inst* 1990;82:531-2.
- 10 Pickard JD, Bailey S, Sanderson H, Rees M, Garfield JS. Steps towards cost-benefit analysis of regional neurosurgical care. *BMJ* 1990;301:629-35.
- 11 Robinson R. Economic evaluation and health care: cost-benefit analysis. *BMJ* 1993;307:924-6.
- 12 Davies EA, Clarke CRA, Hopkins AP. Malignant cerebral glioma. II. Patient and relative perspectives on the value of radiotherapy. *BMJ* 1996;313:1512-6.
- 13 Daumas-Duport C, Scheithauer B, O'Fallon J, Kelly P. Grading of astrocytomas: a simple and reproducible method. *Cancer* 1988;62:2152-65.
- 14 Mahoney FI, Barthel DW. Functional evaluation: the Barthel index. *MD Med J* 1965;14:61-5.
- 15 Nouri FM, Lincoln NB. An extended activities of daily living. *Clin Rehabil* 1987;1:301-5.
- 16 Royal College of Physicians. *Standardised assessment scales for the elderly*. London: Royal College of Physicians Publications, 1992.
- 17 World Health Organisation. *WHO handbook for reporting results of cancer treatment*. Geneva: World Health Organisation, 1979.
- 18 Medical Research Council Brain Tumour Working Party. Prognostic factors for high-grade malignant glioma: development of a prognostic index. *J Neurooncol* 1990;9:47-55.
- 19 Bleehen NM, Stenning SP. A Medical Research Council trial of two radiotherapy doses in the treatment of grades 3 and 4 astrocytoma. *Br J Cancer* 1991;64:769-74.
- 20 Sheline GE, Wara WM, Smith V. Therapeutic irradiation and brain injury. *Int J Radiat Oncol Biol Phys* 1980;6:1215-28.
- 21 Leibel SA, Sheline GE. Radiation therapy for neoplasms of the brain. *J Neurosurg* 1987;66:1-22.
- 22 Wara WM, Larson DA. Central nervous system manifestations of radiotherapy. In: Plowman PN, McElwain TJ, Meadows AT, eds. *Complications of cancer management*. London: Butterworth-Heinemann, 1991.
- 23 Freeman JE, Johnston PGB, Voke JM. Somnolence after prophylactic cranial irradiation in children with acute lymphoblastic leukaemia. *BMJ* 1973;iv:523-5.
- 24 Faithfull S. Patients' experiences following cranial radiotherapy: a study of the somnolence syndrome. *J Adv Nurs* 1991;16:936-46.
- 25 Chang CH, Horton J, Schoenfeld D, Salazar O, Perez-Tamayo R, Kramer S, et al. Comparison of postoperative radiotherapy and combined postoperative radiotherapy and chemotherapy in the multidisciplinary management of malignant gliomas. *Cancer* 1983;52:997-1007.
- 26 Burger PC, Green SB. Patient age, histologic features and length of survival in patients with glioblastoma multiforme. *Cancer* 1987;59:1617-25.
- 27 Hutton JL, Smith DF, Sanderman D, Foy PM, Shaw MDW, Williams IR, et al. Development of prognostic index for primary supratentorial intracerebral tumours. *J Neurol Neurosurg Psychiatry* 1992;55:271-4.
- 28 Kristiansen K, Hagen S, Kollevold T, Torvik A, Holme I, Nesbakken R, et al. Combined modality therapy of operated astrocytomas grades III and IV. Confirmation of the value of postoperative irradiation and lack of differentiation of bleomycin on survival time: a prospective multicenter trial of the Scandinavian glioblastoma study group. *Cancer* 1981;47:649-52.
- 29 Whittle IR, Denholm SW, Gregor A. Management of patients aged over 60 years with supratentorial glioma: lessons from an audit. *Surg Neurol* 1991;36:106-11.
- 30 Valk PE, Dillon WP. Radiation injury of the brain. *Am J Neuroradiol* 1991;156:689-706.
- 31 Brada M. Back to the future - radiotherapy in high grade gliomas. *Br J Cancer* 1989;60:1-4.

(Accepted 9 October 1996)

Malignant cerebral glioma—II: Perspectives of patients and relatives on the value of radiotherapy

Elizabeth Davies, Charles Clarke, Anthony Hopkins

Abstract

Objective—To explore the experiences of patients and relatives after the diagnosis and treatment of malignant cerebral glioma.

Design—Two year prospective study with home interviews.

Setting—Six neurosurgery and radiotherapy centres in London.

Subjects—75 patients and 66 close relatives interviewed at diagnosis, 58 patients interviewed after radiotherapy, and 27 interviewed after recurrence.

Main outcome measures—Awareness of likely prognosis, distress, dissatisfaction with radiotherapy, and perception of severe problems in everyday life.

Results—As they began radiotherapy most patients understood that they suffered from a brain tumour (95%; 71/75), but only one quarter (19/75) seemed fully aware of the poor prognosis. Others were unaware (43%; 32/75) or only partly aware (32%; 24/75). The more aware patients were more distressed. Relatives were three times more likely to be aware of the prognosis (67%; 44/66) and were more distressed. Although 39% (29/75) of patients initially made negative comments about radiotherapy, only 17% (13/75) were completely dissatisfied. The decision to accept radiotherapy could be discussed directly with 19 fully aware patients. Twelve found radiotherapy acceptable if it were medically advised or if it improved survival. Assessed by their own reports of problems only 40% of patients achieved a period of stability or remission, yet dissatisfaction with treatment did not increase.

Conclusion—Most patients with malignant glioma initially seemed unaware or only partly aware of the poor prognosis. Relatives were more aware, more distressed, and often concerned to protect patients from full awareness, which made it difficult to explore with patients directly the possible trade off between quality and length of life. Conceptualising the question as a rational choice ignores the social and emotional context of life threatening disease.

Introduction

In some medical settings doctors may perceive treatment to be unwarranted as to prolong a considerably diminished quality of life or to reduce this further by adverse effects may not be justified. Considerations concerning quality of life as opposed to length of life underpin much of the current debate over setting priorities for treatment,^{1,2} and various formulations of how to resolve such conflicts have been published.³⁻⁶ Whether such formulations would gain much favour in the population at large when faced with the disease in themselves or those close to them is uncertain. Some patients with cancer, for instance, view chemotherapy characterised by side effects and a small chance of success as more worth while than do medical staff or a sample from the general population.^{7,8} Moreover, patients with cancer (and those with other serious diseases) may report the quality of their lives in unexpectedly positive terms.⁹⁻¹³

We considered the tension between quality and length of life for patients with malignant cerebral glioma, a disease for which the prognosis is almost uniformly poor.¹⁴⁻¹⁶ In the previous paper in this issue (p 1507) we described a cohort of 105 patients, 92 of

See editorial by
Gregor and Cull
and p 1507

Directorate of
Neurosurgery and
Clinical Neurosciences,
St Bartholomew's
Hospital, London
EC1A 7BE
Elizabeth Davies, clinical
research fellow
Charles Clarke, consultant
neurologist

Research Unit, Royal
College of Physicians,
London NW1 4LE
Anthony Hopkins, director

Correspondence to:
Dr E Davies, Research Unit,
Royal College of Physicians,
London NW1 4LE.

BMJ 1996;313:1512-6

whom were treated with radiotherapy. In common with previous studies, we found a median survival with radiotherapy after diagnosis of 10 months.¹⁶ The median time free from serious disability was only four months. One quarter of those living more than six months after the diagnosis suffered clinical deterioration or disability apparently associated with treatment. A further 42% experienced severe tiredness.¹⁶ This paper explores patients' experiences after the diagnosis and in particular their acceptance of radiotherapy. We also describe the views of close relatives concerned in their care.

Patients and methods

DATA COLLECTION

Recruitment to the study and follow up by home visit is described in the accompanying paper.¹⁶ We did not consider that a questionnaire approach was an appropriate way to broach the sensitive issues we needed to explore. We therefore began the research using open ended interviews with eight patients and relatives. These interviews informed the areas to be explored in the main study, in which we used a semistructured interview format. This method allows respondents to talk at length about any matter that concerns them and yet leaves the interviewer scope to explore difficult issues.^{17 18} A series of four point rating scales on the basis of interviewer interpretation were developed to cover the range of experiences and meanings expressed. Table 1 shows the principal topic areas. An accompanying manual (available from the first author) defined the rating scale points ("marked," "moderate," "some," and "nil") for each topic. It also provided anchoring examples on the basis of verbatim interview comments as standards for the level of detail to be obtained. Our rating of distress, for example, took account of all the concerns expressed and the degree of emotion that was conveyed. Patients or relatives constantly upset and overwhelmed by the situation were rated as "highly" distressed while those who felt upset or anxious a good deal of the time but found pleasure in some aspects of life were rated as "moderately" distressed. Those feeling only occasionally depressed, anxious, or dismayed or remaining generally cheerful were rated as showing "little" or "no" distress.

Data collection was shared by three interviewers with social science training, one of whom (ED) was also medically qualified. All interviews were tape recorded and transcribed in full, with notes of emotional content

Table 2—Demographic characteristics of 105 patients and 91 relatives. Figures are numbers (percentages) of subjects

Characteristic	Patients	Relatives
Age (years):		
21–44	26 (25)	30 (33)
45–59	39 (37)	35 (38)
60–75	40 (38)	26 (29)
Sex:		
Men	72 (69)	29 (31)
Women	33 (31)	62 (69)
Racial origin:		
Afro-Caribbean	2 (2)	2 (2)
Asian	5 (5)	2 (2)
White	98 (93)	87 (96)
Age leaving full time education (years):		
≤16	53 (58)*	49 (56)†
18	18 (20)	26 (30)
≥21	20 (22)	12 (14)
Social class ²⁵ :		
I	12 (11)	12 (13)
II	31 (30)	27 (31)
IIINM	23 (22)	25 (27)
IIIM	25 (23)	17 (19)
IV	6 (6)	4 (4)
V	8 (9)	6 (7)
Marital status:		
Single	11 (10)	9 (10)
Married/cohabiting	82 (78)	78 (86)
Widowed	6 (6)	1 (1)
Divorced	6 (6)	3 (4)

*Missing data in 14 cases.

†Missing data in 4 cases.

such as pauses, false starts, sighs, and weeping. Table 1 shows results of a reliability study, in which 13 transcribed interviews were rated by each of the three raters. Reliability was good or satisfactory¹⁹ particularly when rating scales were dichotomised between "marked/moderate" and "some/nil" (table 1).

ANALYSIS

The themes emerging from the first 30 interviews have been briefly described elsewhere.²⁰ These interviews suggested that we should consider patients' views at three stages in the clinical course rather than at set intervals after the diagnosis. These were, firstly, coming to terms with the diagnosis, secondly, adapting to change during any remission, and, thirdly, a final stage of deterioration and increasing disability. The interviews also suggested that we should explore differences in awareness of the likely prognosis between patients and relatives. In our accompanying paper we showed that the patient's initial disability assessed by the World Health Organisation clinical performance status²¹ was an important prognostic factor.¹⁶ In this paper we compared this measure with patients' own views about their problems. We used the χ^2 test and χ^2 test for trend throughout.

Results

Table 2 summarises some demographic characteristics of the sample. As many as 41% of patients were from professional and managerial classes,²² and just over two thirds were men. Sixty nine percent of carers were women, and most carers were spouses or partners (71/91; 78%).

Because of the severity of their disability, none of the 13 patients who received conservative treatment with steroids could be interviewed in any depth. The main analysis therefore concerns the 92 patients who received radiotherapy. Of these, 75 could be interviewed within three months of the diagnosis, and 59 could be seen for at least one follow up interview. Twenty seven of these were seen again at a time when they had deteriorated after an initial period of stabilisation or improvement.

Table 1—Inter-rater reliability study for 13 interviews*

Detail	κ †	
	4 Point scale	2 Point scale‡
Awareness of likely prognosis		
Belief in possibility of cure	0.58	0.77
Thoughts about possibility of death	0.70	0.55
Problems caused by illness in everyday life		
Hobbies or interests	0.69	0.88
Household tasks	0.55	0.80
Social life	0.59	0.86
Self care	0.60	0.60
Cognitive ability	0.84	1.00
Satisfaction with treatment		
Negative comments about radiotherapy	0.77	0.76
Overall satisfaction with radiotherapy (2 point only)	—	0.86
Social relationships		
Perceived support from relative or carer	0.67	0.69
Overall distress	0.66	0.53

*Thirteen interviews were rated by each of three interviewers and κ s were calculated for 39 pairs of ratings obtained. Pairs of interviewers were compared an equal number of times. Disagreements were settled by consensus when these ratings were entered into main analysis.

† κ Indicates level of agreement between raters. Value of 0.41–0.60 show moderate agreement, those of 0.61–0.80 good agreement, and 0.81–1.00 very good agreement.

‡Two point scales represent dichotomy between "marked" and "moderate" v "some" and "nil."

Death or obvious confusion prevented further patients being seen at this stage of the illness.

AWARENESS OF THE PROGNOSIS

The patients

At our first interview, as they began radiotherapy, we asked patients to recount in detail what they had been told in their medical encounters and what had been said by relatives and friends. From this it was clear that most patients receiving radiotherapy understood they suffered from a brain tumour (95%; 71/75). Awareness of the prognosis, however, varied considerably. We defined three levels of awareness taking into account the degree to which patients believed that radiotherapy would cure them and whether they seemed to be aware that they might die from their disease. From the tape recordings we considered that only a quarter (19/75) were fully aware of the prognosis, conveying that there was little chance of cure and expressing some thoughts or fears about the possibility that they might die. A 56 year old man with a parietal tumour, for example, spoke frankly about radiotherapy:

"Well, fundamentally it just delays the process. It doesn't stop it. Wish to God it did but it doesn't. We talk about it all the time that I'm going to die, that it's going to happen after Christmas or whenever."

We rated just under a third of patients (32%; 24/75) as being partly aware of the likely outcome. One 40 year old man with a frontal tumour, for example, remarked:

"The family are still frightened that something might happen. Do you know what I mean? Because you read about it in the papers everyday don't you—this trouble? Cancer—person dies from brain tumour. So I suppose it's at the back of your mind as well."

Such patients usually conveyed some fear of dying but also that they had a reasonable chance of cure. As many as 43% (32/75) of patients, however, conveyed in the interviews practically no awareness that they might die. One 40 year old man with an occipital tumour, for example, when asked about the chance of cure said,

"I didn't specifically ask the surgeon. He spoke as if it were pretty certain. I can see that people who had only biopsy and growths in awkward places, that there would be a question mark over how complete treatment would be. But in my case, because it seemed to be something that was on the surface...then you should be able to wipe it out completely."

We had initially thought that patient optimism might be explained by high dose steroids²⁰ and obtained data on the patients' doses at the time of interview in 62 cases. A comparison of low dose (0-2 mg) with moder-

ate (3-8 mg) and high doses (9-32 mg) of steroid, however, showed this had no influence on awareness.

The relatives

Sixty six of these 75 patients had relatives who could also be interviewed at diagnosis. These relatives were three times more likely than the patient to be aware of the poor prognosis (67% (44/66) v 21% (14/66); table 3). The wife of the unaware patient quoted above said:

"They told my husband it was a tumour, but they didn't tell him about the prognosis or anything like that.... I caught the surgeon in the corridor and he said that my husband could have been dead in a month if he hadn't had the operation. He said that some people did well—it could be 18 months, or even 5 years but not really—5 years was quite rare. It knocked me for six."

The implications of such news was also often spelt out. A 50 year old husband remarked:

"From now on we're living with a death sentence. The doctors said the tumour could come back quite easily, and I got the feeling they were expecting it to."

Only 9% (6/66) of relatives therefore were unaware of the likely prognosis. The agreement between awareness of patients and relative was poor (κ 0.20). Moreover, there was a consistent tendency for patients to show less awareness (39 out of 66 comparisons between patients and relatives).

On the whole, relatives tended to see their role as one of remaining strong to protect the patient in the hope that they would be among the most fortunate. It was not uncommon for them to convey that to be explicit about the prognosis would be harmful:

"If you say you've got six months or whatever, it does mentally act as a sentence to that person."

As the illness progressed, slightly more patients became fully aware of the prognosis. Most patients interviewed after deterioration were therefore aware that they were dying (52%; 14/27), although 26% (7/27) still conveyed no awareness of this possibility and the 22% (6/27) remaining were partly aware. At all stages of the disease most patients felt highly supported by their relatives (92% (61/66) at diagnosis and 77% (21/27) at deterioration).

DISTRESS EXPERIENCED BY PATIENTS AND THEIR RELATIVES

Despite the seriousness of the condition patients were less distressed than might have been expected. Of the 75 patients who could be interviewed within three months of the diagnosis, we rated only 9% (7/75) as highly distressed. Six patients seemed severely depressed or anxious, and one expressed extreme anger. One fifth (16/75) showed a moderate level of distress, but more than two thirds of the patients (54/75) felt only occasionally depressed, anxious, or dismayed and remained generally cheerful or confident. The level of the patient's distress was moderately related to their awareness of the prognosis (γ 0.53; table 4). Fifty eight percent (11/19) of the fully aware showed marked or moderate distress compared with one quarter of those partly aware (6/24) and 16% of those unaware (5/32) (χ^2 for trend 9.53; 1 df; $P < 0.01$).

In clear contrast with patients, nearly two thirds of relatives experienced a marked or moderate level of distress (64%; 42/66). In most instances, relatives were in fact more distressed than the patients they cared for (58%; 38/66); in only a few instances did the patient seem more distressed (14%; 9/66). Too few relatives remained unaware of the prognosis to explore whether this factor was associated with lower levels of distress.

VIEWS ABOUT RADIOTHERAPY

Initial dissatisfaction with radiotherapy

A substantial number of patients (39%; 42/66) made "marked" or "moderate" negative comments about

Table 3—Comparison of awareness of patients and relatives or carers of likely prognosis shortly after diagnosis

Awareness of patients (n = 66)	Awareness of relatives (n = 66)			
	Fully aware	Some idea	Unaware	Total
Fully aware	13	1	—	14
Some idea	15	7	—	22
Unaware	16	8	6	30
Total	44	16	6	66

Table 4—Awareness of likely prognosis and distress in patients shortly after diagnosis. Figure are numbers (percentages) of patients

Awareness	No of patients	Patient distress		
		Nil	Some	Marked/ moderate
Full awareness	19	2 (11)	6 (32)	11 (58)
Partial awareness	24	5 (21)	13 (54)	6 (25)
No idea	32	14 (44)	13 (41)	5 (16)

Table 5—Initial disability, perception of disability, and proportion of patients reporting
Figures are numbers (percentages) of patients

Initial level of patient disability*	Patients initially reporting at least one severe problem in everyday life†	Patients able to report a period of stability or a remission‡
0-1	7/22 (32)	16/21 (76)
2	20/29 (69)	14/36 (39)
3-4	22/24 (92)	4/28 (14)
Total	75 (100)	85 (100)

* As assessed by WHO clinical performance status.

† "Severe" problems in everyday life refer to ratings made with 2 point scales shown in table 1.

‡ Here we considered all follow up interviews and identified one at which patient reported fewest problems. Period of stability was defined as same number of severe problems at diagnosis interview and at any follow up interview; remission was reduction in number of severe problems reported by patient. Follow up result based on 85 patients as six refused any follow up and data missing for one. Those too confused initially to interview or who died before follow up are classed as not achieving period of stability. Two patients reporting 3-4 severe problems at both points are included as having achieved period of stability as although they continued to have many problems, lack of deterioration may have been attributable to treatment.

radiotherapy at the initial interviews, expressing their dislike of the procedure, adverse effects, or doubt about its value. This compared with 29% (22/74, data missing in one case) of patients who expressed equal dissatisfaction with surgery. Such negative comments about radiotherapy were more common among patients with marked or moderate distress (59% (13/22) *v* 30% (16/53) with little or no distress; χ^2 4.33; $P < 0.05$), but such comments did not relate to awareness of the prognosis or initial clinical performance status. We rated only one fifth of the patient sample (17%; 13/75) as being dissatisfied in an overall sense, however, as they showed considerable reluctance about undergoing radiotherapy. This level of dissatisfaction tended to be more common among patients who were fully aware of the prognosis (32% (6/19) *v* 11% (6/56) of those partly or unaware; χ^2 3.17; 1 df; $P < 0.1$), but this difference was not significant.

We thought it appropriate to explore the decision to accept radiotherapy only with the 19 patients fully aware of the prognosis. Of these, six accepted treatment with reluctance because they saw no alternative but remained dissatisfied. Four accepted treatment as it had been medically advised, and nine believed they could increase their survival by fighting the disease, having accepted radiotherapy for the chance of survival it brought:

"I began to fight it, just began to start the battle.... What I am doing is living for the future. I would like to think that I've got at least a year or so, or more. I'm thinking if you like of a strategy in that I've got to get planned what I can do."

Patients' improvement, deterioration, and subsequent dissatisfaction with radiotherapy

Patients' initial perceptions of problems in everyday life (table 1) correlated, as might be expected, with ratings of disability based on the WHO clinical performance status (table 5). In case dissatisfaction after radiotherapy related to the patient's perception of not improving after treatment, we first determined which patients could be classed, in their own terms, as having achieved a period of stability or a remission. Overall we found that less than a half (40%; 34/85) of those patients treated with radiotherapy on whom we had follow up data achieved either a period of stability or a remission. The most severely disabled patients, whether judged by their own perceived problems or the WHO clinical performance status, were least likely to achieve a period of stable or improved function (table 5). Patients who achieved neither tended to make more negative comments about treatment at their best follow up interview (50% (20/40) *v* 22% (4/18); χ^2 2.89; 1 df; $P < 0.10$), although this was not significant; and patients making such comments were more likely to be markedly

or moderately distressed (58% (15/26) *v* 21% (7/32); χ^2 6.37; 1 df; $P < 0.01$). The proportion of patients dissatisfied in an overall sense with radiotherapy, however, remained stable at one fifth (21%; 12/58). Dissatisfaction at follow up did not relate to awareness of the prognosis, distress over hair loss, or level of distress. At deterioration, fewer patients made highly negative comments about radiotherapy (18%; 5/27) than on completion of radiotherapy (50% (20/40); χ^2 5.55; 1 df; $P < 0.01$). Only one patient entirely regretted having undergone treatment, although in such serious circumstances at this stage of the illness it did not seem appropriate to probe too deeply. Overall we could not therefore find evidence that most patients looked back over the illness feeling dissatisfied that they had undergone radiotherapy.

Discussion

In this study we interviewed 75 patients with malignant glioma and 66 of their relatives to explore the social context of one disease in which treatment can only extend survival. Our findings show some of the difficulties of obtaining clear cut answers to questions about the trade off between quality and length of life. We suggest these difficulties relate to the very nature of such severely threatening situations.

In our study, detailed interviews with patients shortly after the decision to have radiotherapy had been made showed that only one quarter (19/75) seemed to be fully aware of the strong possibility that they would die from their disease and that the best they could expect from radiotherapy was some prolonging of life. Assessed by their own reports of problems, only 40% of patients achieved a period of stability or a remission after radiotherapy. In our accompanying paper¹⁶ we described the extent of adverse effects associated with radiotherapy, yet in terms of this, and the eventual outcome, relatively few patients conveyed that they regretted having undergone treatment. There was a suggestion that patients who had not achieved a remission made more negative comments about radiotherapy, but little more than one fifth of the sample were dissatisfied in an overall sense, and dissatisfaction did not increase with time.

Social classes I and II are disproportionately represented in our sample. This may reflect the distribution of the disease in the population around London or more probably differences in the referral of patients from different social classes to specialist centres. We do not believe this bias reflects selective recruitment as we identified all available patients in a consecutive manner at each centre. Despite this, however, relatively few patients demonstrated the level of awareness that would have allowed us to explore their thoughts about the kind of trade off they might have been making. Other studies have avoided this problem by asking patients about hypothetical situations rather than their own, and this indirect method has shown that cancer patients are prepared to consider more intensive treatment regimens than members of the general population.^{7, 8} That most of the patients whom we could directly question were prepared to undergo radiotherapy for the chance of longer survival is consistent with the results of such studies.

The possibility that the patients' lack of awareness of the prognosis and their acquiescence to radiotherapy was directly due to their brain disease is made unlikely by the fact that severely confused patients were excluded. The rest were interviewed at length and gave informed and interesting answers to many of our questions. Lack of awareness of the prognosis was also unrelated to the patients' dose of steroids at the time of interview. Part of the explanation for limited awareness seemed to lie in the fact that many had been protected by relatives and medical staff. Patients' relatives, for instance, were three times more likely to be aware of the

Key messages

- Most patients with malignant cerebral glioma do not have a clear idea of the prognosis for their disease as they begin treatment. Those who are aware are more likely to be distressed
- In general, relatives are more distressed than patients and more likely to be aware of the poor prognosis but tend to want to protect patients from this knowledge
- Communication with patients needs to take into account their need to make informed decisions about treatment and the implications of the disclosure of such information for communication with families
- Patients who initially have better WHO clinical performance status (0-1) are most likely to report improving or remaining stable after radiotherapy. Overall only 40% of a non-trial population thus achieve a remission

probable outcome and showed more distress in the interviews. Some patients may have had subtle neuropsychological deficits affecting the way they attributed significance to new information, and in some cases such factors may have reinforced the desire of relatives to protect patients from distress. Another recent interview study found that only 11 out of 30 patients with malignant brain tumours associated their tumours with death²³ and described different psychological strategies that patients used to maintain hope. In our study, patients who were aware of the poor prognosis, however, were more distressed. This finding is consistent with the suggestion in some studies that when the prognosis is very poor, some patients may prefer not to know²⁴ or hold on to the belief that they are being treated for a cure.²⁵ It is clear that issues of communication are important in this disease, particularly as it is now generally accepted that communication with cancer patients should be straightforward and open.²⁶⁻²⁸ Our findings underline that the consequences of the communication style adopted by medical staff need to be considered in terms of distress for both the patients and their families, particularly when many families seem to adopt a coping style which is not based on the continuous and explicit discussion of the truth.²⁹⁻³¹

One explanation that would make sense of our findings is that the question of a trade off between quality and length of life is one more often raised by concerned observers than by those facing the possibility of imminent death or bereavement. This observer perspective seems to emphasise the possibility of a rational choice at the cost of largely ignoring associated emotions. In reality many of the patients and their relatives took hope from the apparent element of clinical uncertainty and from the fact that some treatment was being attempted. This strategy may have allowed them to cope with the day to day problems that the disease caused and, by keeping their distress within manageable levels, helped to maintain the quality of their life for a short period. Observers tend to interpret such coping as creating only "false hope", but an alternative perspective is to accept that this need for hope is an objective feature of such situations which cannot be rationalised away. Once accepted, it could be debated whether this need for hope should be incorporated into quality of life measures or into calculations about the cost of interventions to the NHS. In short, the dilemma is how much can health services afford costly interventions which provide principally hope, even though that hope improves the quality of life?

We thank the patients and relatives who agreed to talk to us. Maureen Bannon and Sue Hall helped with the development of the interview and rating scales and with data collection. Rachel Warrick and Valerie Pottinger gave administrative support. We are also grateful for the comments of three anonymous reviewers.

Funding: Cancer Research Campaign grant number CP 1017.

Conflict of interest: None.

- 1 Royal College of Physicians. *Setting priorities in the NHS. A framework for decision-making*. London: Royal College of Physicians, 1995.
- 2 Government Committee. *Choices in health care*. Netherlands: Ministry of Welfare, Health and Cultural Affairs, 1992.
- 3 Rosser RM. From health indicators to quality adjusted life years: technical and ethical issues. In: Hopkins A, Costain D, eds. *Measuring the outcomes of medical care*. London: Royal College of Physicians of London and King's Fund Centre for Health Services Development, 1990:1-17.
- 4 Rosser R, Cottey M, Rabin R, Selai C. Index of health-related quality of life. In: Hopkins A, ed. *Measures of the quality of life and the uses to which they may be put*. London: Royal College of Physicians of London, 1992:81-90.
- 5 Williams A, Kind P. The present state of play about QALYs. In: Hopkins A, ed. *Measures of the quality of life and the uses to which they may be put*. London: Royal College of Physicians of London, 1992:21-39.
- 6 Ebrahim S. Clinical and public health perspectives and applications of health-related quality of life measurement. *Soc Sci Med* 1995;41:1383-93.
- 7 Slevin M, Stubbs L, Plant H, Wilson P, Gregory WM, Armes PJ, et al. Attitudes towards chemotherapy: comparing views of patients with cancer with those of doctors, nurses and the general public. *BMJ* 1990;300:1458-60.
- 8 O'Connor, A. Effects of framing and level of probability on patients' preferences for cancer chemotherapy. *J Clin Epidemiol* 1989;42:119-26.
- 9 Cassileth B, Lusk E, Strouse T, Miller DS, Brown LL, Cross PA, et al. Psychosocial status in chronic illness: a comparative analysis of six diagnostic groups. *N Engl J Med* 1984;311:506.
- 10 Breetvelt IS, Van Dam FSAM. Under reporting by cancer patients: the case of response shift. *Soc Sci Med* 1991;32:981-7.
- 11 McGee H, O'Boyle C, Hickey A, O'Malley K, Joyce CR. Assessing the quality of life of the individual: the SEIQoL with a healthy and a gastroenterology unit population. *Psychol Med* 1991;21:749-59.
- 12 O'Boyle C, McGee H, Hickey A, O'Malley K, Joyce CR. Individual quality of life in patients undergoing hip replacement. *Lancet* 1992;339:1088-91.
- 13 Kreitler S, Chaitnick S, Rapoport Y, Kreitler H, Algor R. Life satisfaction and health in cancer patients, orthopaedic patients and healthy individuals. *Soc Sci Med* 1993;36:547-56.
- 14 Walker MD, Alexander E, Hunt WE, MacCarty CS, Mahaley J, Norrell HA, et al. Evaluation of BCNU and/or radiotherapy in the treatment of anaplastic gliomas. *J Neurosurgery* 1978;49:333-43.
- 15 Bleeher NM, Stenning SP. A Medical Research Council trial of two radiotherapy doses in the treatment of grades 3 and 4 astrocytoma. *Br J Cancer* 1991;64:769-74.
- 16 Davies EA, Hopkins AP, Clarke CRA. Malignant cerebral glioma. I. Survival, disability, and morbidity after radiotherapy. *BMJ* 1996;313:1507-12.
- 17 Brown GW, Rutter M. The measurement of family activities and relationships: a methodological study. *Human Relations* 1966;19:370-4.
- 18 Brown GW, Harris TO. *The social origins of depression*. London: Tavistock, 1978.
- 19 Brennan P, Silman A. Statistical methods for assessing observer variability in clinical measures. *BMJ* 1992;304:1491-4.
- 20 Davies E, Clarke C. Malignant cerebral gliomas: rehabilitation and care. In: Greenwood R, Barnes MP, McMillan TM, Ward CD, eds. *Neurological rehabilitation*. London: Churchill Livingstone, 1993:535-43.
- 21 World Health Organisation. *Handbook for reporting results of cancer treatment*. Geneva: World Health Organisation, 1979.
- 22 Office of Population Censuses and Surveys. *Standard occupational classification*. London: HMSO, 1991.
- 23 Salander P, Bergenheim T, Henrikson R. The creation of protection and hope in patients with malignant brain tumours. *Soc Sci Med* 1996;42:985-96.
- 24 Fallowfield L, Ford S, Lewis S. No news is not good news: information preferences of patients with cancer. *Psycho-oncology* 1995;4:197-202.
- 25 MacKillop WJ, Stewart WE, Ginsberg AD, Stewart SS. Cancer patients' perceptions of their disease and its treatment. *Br J Cancer* 1988;58:355-8.
- 26 Maguire P, Faulkner A. How to do it: communicate with cancer patients. 1. Handling bad news and difficult questions. *BMJ* 1988;297:907-9.
- 27 Maguire P, Faulkner A. How to do it: communicate with cancer patients. 2. Handling uncertainty, collusion and denial. *BMJ* 1988;297:972-4.
- 28 Fallowfield L J. Giving sad and bad news. *Lancet* 1993;341:476-8.
- 29 Hinton J. *Dying*. London: Penguin, 1972.
- 30 Kuebler-Ross E. *On death and dying*. London: Routledge, 1992.
- 31 Glaser BG, Strauss AL. *Awareness of dying*. London: Weidenfeld and Nicolson, 1966.

(Accepted 8 October 1996)

Correction

Causality, menopause, and depression: a critical review of the literature

Three errors occurred in figure 1 of this paper by Louise Nicol-Smith (16 November, pp 1229-32). The perimenopausal area that is crosshatched in Avis *et al*⁹ 1993 should have been plain while that in Avis *et al*⁸ 1994 should have been crosshatched. The dotted box in the age column that straddles Hunter²⁵ 1986 and Ballinger *et al*²⁶ 1985 should have been with Holte¹⁷ 1992. The note on sample size in the legend is wrong owing to an editorial error. It should have read: "Note: Sample size generally was the number available for the relevant data analysis. However, in the cases of Hällström and Samuelsson³³ 1985, Køster and Davidsen⁶ 1993, and Matthews *et al*¹⁴ 1994 and 1995 the number stated is the sample size at study outset."