Observations

ON

SUPRAPUBIC PROSTATECTOMY.

By SIR THORNLEY STOKER, M.D., F.R.C.S.I.,

President of the Royal Academy of Medicine in Ireland; Surgeon,
Richmond Hospital, Dublin.

THE performance of suprapubic prostatectomy is of such comparatively recent origin, and is as yet so much in course of development, that I have an apology for recording the small number of three cases. I do so because every example of an operation of such magnitude is an item of some value. Besides, although I have done but three prostatectomies by this route, I have performed a large number of suprapubic cystotomies for other diseases, and the treatment of the bladder after every suprapubic opening is a matter of importance, and one in which so much is learned by experience, that I wish to describe not only the method I have followed in dealing with the prostate gland itself, but also the points of interest in the treatment of the wound and the bladder during and subsequent to operation. The mere removal of the gland is usually a more easy and simple matter than the after-treatment of the case. I have had the advantage of assisting Sir William Thomson at most of his prostatectomies and have added to my experience in that way. The description of his technique by a surgeon may be interesting to his colleagues, and I offer mine with every respect, knowing how in my own surgical work I have been helped by a practical account of details.

When the bladder and the urine are alkaline and when delay in operating is permissible it is of great advantage to get them into a better condition by the use of a white diet, by washing, and by the use of such other measures or drugs as may appeal to individual surgeons.

I propose to speak first of the removal of the prostate when the bladder has been opened, and afterwards of the method of opening the bladder and the treatment of the parts subsequently. The two latter matters are common to all suprapubic cystotomies and hang naturally together.

When the bladder has been opened, the operator standing on the left of the patient, the prostate should be thoroughly examined with the index finger of the right hand in the bladder, assisted by the left index or the left index and middle fingers if necessary in the rectum; great care should previously have been given to the cleansing of the rectum. Owing to the necessity of having at least one finger in it during most of the time occupied in enucleating the prostate ideal cleanness cannot be achieved without the use of a rub-

ber glove for the rectum, and this lessens the delicacy of manipulation.

During this examination the surgeon must determine whether one or both sides of the gland require removal, and whether some myomatous outgrowth may not have been the cause of obstruction and its removal suffice to open the passage for urine. If the prostate itself is essentially enlarged, or if a nodular condition of the gland due to separate capsulated tumours of an adenomatous or a myomatous nature exist, the entire gland should be removed. If any partial operation be done in such cases, a fresh development of some of these tumours is likely to occur later on and produce fresh obstruction. In one of my cases the obstruction to the urine, which was complete and acute, was not due to the general enlargement of the gland but to a projecting nodule not much larger than a pea; the gland

itself was full of such small tumours, and had it been left behind there is no doubt that sooner or later a fresh obstruction would have taken place either from the growth of such a nodule or from the general gland hypertrophy. When this latter condition exists, I believe it always produces obstruction by lateral approximation of the lobes of the gland towards the middle line so as to press one against the other. The so-called middle lobe I do not believe in, and I think that the outgrowth of a separate adenoma or myoma is what has been so christened.

When the prostate has been thoroughly examined the left index, washed after its removal from the rectum, is placed in the bladder and used as a guide for a pair of blunt-ended scissors, which should be used to snip through the floor of the bladder and capsules of the gland, on one side of the middle line in an antero-posterior direction. This incision should be made over the most projecting part of the larger of the two lobes. It may be found that the whole gland can be delivered through one incision, but if the lobe

Fig. 1.

first incised comes away without its fellow, a second and similar cut can be made on the other side and the remaining lobe disencapsuled. I think it is of the utmost importance to make the incision of sufficient depth, not merely seeking to divide the bladder wall and fascial capsule of the prostate, but to cut for a short distance through its proper capsule into the gland substance. If it be merely sought to expose the gland without injury to its substance, the operator cannot be certain whether he has thoroughly divided its coverings so as to render its enucleation easy. I hold strongly—and have formed the opinion as the result of anatomical as well as surgical observation—that the safest and most easy way to be sure that you are within the capsule of pelvic fascia, and in a region where enucleation can be safely and rapidly effected, is to snip with your scissors into the substance of the prostate.

The incision I have been speaking of need not be more

than 1 in. in length, enough to admit the end of a finger easily; any increase of its size can be effected by tearing with the point of the finger. It will be found after the removal of a large prostate has been effected that the incision or rent in the bladder wall has contracted to a very small size. The scissors are then withdrawn, the left index, with the middle finger if required, again placed in the rectum, and the right in the bladder. The process of enucleation is effected as rapidly as possible, the left fingers being used for counter-pressure and for the protection of the rectal wall from perforation or other injury. In one case I enucleated a prostate weighing $4\frac{1}{2}$ oz. in ten minutes. I have seen much difficulty experienced in enucleating a sclerosed prostate, but in most cases ten to twenty minutes suffice for this stage of the operation.

The question of attempting to conserve the prostatic urethra generally settles itself. The urethra may be so glued to the gland that it separates with it, and comes away attached to the prostatic substance. The prostatic urethra may come away with a soft and easily disencapsuled prostate, and, on the other hand, may remain intact in a case of old and hard hypertrophy. Records seem to show that cases do just as well in which the prostatic urethra has been removed as those in which it has been preserved. Inasmuch as the chief source of hæmorrhage in this operation is from veins between the bladder and fascial capsule or between the fascial capsule and the true capsule of the gland, we find another reason for getting well into the gland so as to avoid extensive injury to them.

When the prostate has been removed, the bladder should be irrigated with hot water to remove clots and check any oozing which may be present. As a rule, hæmorrhage gives no trouble, but if it should occur subsequently to a grave extent a repetition of the irrigating process will usually stop it, and it is in view of this being necessary that one reason is found for leaving the bladder and abdominal wound as open as possible. In this context it may be remarked that the bladder generally contains a good deal of clotted blood for two or three days subsequent to operation, and it is by no means desirable to be too diligent in removing it. Its presence seals the large veins about the prostatic area, and limits both the tendency to bleeding and to septic absorption, so that it is not well to make much effort to remove clots. The urine has a rapid disintegrating effect on them, and they are soon got rid of without effort. I seldom use more than the most gentle syringing of the bladder through the wound for some days after a suprapubic operation. When more energetic washing is called for, it is best effected by injecting boric lotion through a catheter introduced by the urethra, and allowing it to discharge through the abdominal wound.

I now pass to the method of opening the bladder above the pubes. The viscus should be thoroughly washed out, and about 4 to 8 oz. of warm boric lotion allowed, if possible, to remain in it. But even if its contracted state forbids this moderate distension, there should be no difficulty in finding it when the parietal incision has been made. For many years I have abandoned the use of Petersen's bag, as it is both unnecessary and a source of danger by the strain it puts on an

unhealthy bladder. The parietal incision requires to be from 3 to 4 in. long, according to the thickness of the abdominal wall, and should extend to within $\frac{1}{2}$ in. of the symphysis. Once the skin and the musculo-tendinous structures are divided, no more use need be made of the scalpel except to pierce the bladder wall; all the tissues may be got out of the field and the bladder exposed by scratching with a half-blunt instrument, such as the point of Greig Smith's scissors. The peritoneal pouch seldom gives trouble, even when the bladder is contracted, and can readily be held up by the finger of the assistant or by a small retractor. The bladder is best opened by puncturing with a small scalpel, and passing into the opening the blades of Greig Smith's scissors. If these blades are then separated in a vertical direction, they may be used to tear the opening in the bladder to any necessary size, and while doing so to lift it to the surface, so as to enable the operator to insert his finger. If the opening requires further enlargement, or is difficult to tear, owing to hypertrophy of the bladder coats, it can be increased in size by snipping with scissors.

I then take a piece of thick silk which at each end is threaded into the eye of a large fully-curved needle; the point of one needle is passed into the bladder and, guided

by the left index, is made to pass forward through the bladder wall on one side of the incision, at its upper end and about $\frac{1}{2}$ in. from its edge, and then through all the tissues of the abdominal wall anterior to the peritoneum except the skin. This proceeding is then repeated with the second needle on the opposite side; no further use of this suture is made until the intravesical portion of the operation has been completed.

When the operation is ended and the bladder has been irrigated the suture should be tied, not tightly but with just enough force to approximate the opposed surfaces of the subcutaneous portions of the tissues at the upper end of the wound. It should be tied with a bow and the ends left long so that

it may be loosened if necessary. As it does not include the skin it causes no pain. The uses of this suture are as follows:

1. By drawing the bladder and sides of the parietal incision together it effectually prevents any prolapse of the peritoneal sac.
2. It keeps the opening in the bladder accessible, so that a tube can readily be introduced and washing easily and safely performed if called for.
3. It prevents undue contraction of the bladder towards its floor, and provides space for the collection of a small amount of urine, so helping to limit the overflow, which at times wets the patient and causes trouble.
4. By fixing the bladder to the upper end of the wound, and keeping the peritoneal pouch above the area, it prevents any chance of the cicatrix being afterwards subjected to bowel pressure and becoming hernial; and it leaves a cicatrix in such a position and under such conditions that it may, if necessary at any time in future, be safely incised for access to the bladder.

The suture may be removed on the fourth or fifth day, when lymph exudation will have cemented the parts, and made the treatment of the wound a comparatively simple matter.

Fig. 2.

In performing a suprapubic cystotomy the endeavour should be to form a wound with its apex at the bladder and its base at the skin surface. It ought not, as a rule, be closed or sutured in any way, except by the solitary suture I have described. In cases where, owing to the presence of a large amount of fat, the incision through the superficial tissues has been extended much upwards, a single suture of silkworm gut may be introduced towards the superior end of the wound. Attempt to suture the bladder or parietal incision is not only useless but is injurious, and helps to cause infiltration of urine.

Since I have adopted this method I have had entire freedom from the infiltration of urine, which so often gives trouble in the tissues of the abdominal wall or pelvis, producing suppuration and sloughing. Nor have I seen any trouble from the phosphatic concretions which may cause so much suffering and delay in the process of healing.

When the operation of cystotomy has been completed a rubber drainage tube of large size is introduced; it should extend from the surface level into the bladder, but not touch its floor, as if it does it will cause pain resembling that produced by vesical calculus. It should be attached at the surface by a rectangular glass bend to a lighter tube, which is passed over the side of the bed into a receptacle for urine. The wound may be loosely packed with gauze, not for the purpose of occluding it, but in order to act as a conductor of urine and prevent infiltration.

A covering of cellulose wadding and a binder loosely applied, will, after the bladder clots have come away, suffice to keep the patient and the bed dry and prevent much suffering.

The wound when treated in this way heals in from three to four weeks, leaving a narrow cicatrix.

CASE I.

B. O'B., aged 60, a farmer, of medium stature, in spare muscular condition, was admitted to the Richmond Hospital on the recommendation of Dr. Robert Creighton, of Ballyshan-non, on October 6th, 1902.

Two years previously he had suffered from frequency of urination for a few weeks. The attack passed off without any stoppage of water, and he remained in good health until five weeks before his admission to hospital, when he was suddenly seized with complete retention, and up to the date of operation was unable to empty his bladder except with a catheter. When he came under observation his urine was slightly alkaline, contained a little pus and some consequent albumen, and had a specific gravity of 1020. His prostate examined by the rectum revealed a moderate general enlargement of the gland, most marked on the left side; a No. 11 (E) catheter was readily introduced.

Treatment by systematic catheterization, by sodium benzoate, and a white diet was used, and, his urine having become acid, the prostate was removed on October 15th.

The details of the operation and subsequent treatment were in this and the following cases essentially those already detailed.

Recovery was uneventful. A little vesical bleeding took place some sixteen hours after operation, and was checked by the introduction of adrenalin solution. The temperature was never very high, and was practically normal after the ninth day. On November 3rd—the nineteenth day—water was passed per urethram, and from this on the drainage tube was only used at night.

On November 12th (the twenty-eighth day) the man was allowed out of bed into an armchair, and from this the tube was abandoned. The wound was quite healed when he left hospital on December 12th, 1902.

Examination of the parts in this case, both before and after their removal, showed that the general hypertrophy of the prostate had not as yet caused the complete obstruction to urination which existed, but that it was due to a small rounded outgrowth, about the size of the tip of a little finger, which projected into the prostatic urethra. Dr. Earl reports it as being a myoma, and weighing 16 gr. This little tumour explains the unusual circumstance of the sudden complete stoppage of urine in a case where a large catheter was easily passed.

The following is Dr. Earl's report on the parts removed:

"Right lobe, weight 210 gr.; length, 1½ in.; small round mass, 16 gr.; circumference, 4½ in. Two masses of left lobe—one weighs 239 gr.; length, 2 in.; circumference, 4½ in.; second weighs 142 gr.; length, 1½ in.; circumference, 4 in.

"The total weight, therefore, is 11 gr. less than 2 oz. The small round mass is a myoma. There is no glandular tissue contained in it, and only a very small quantity of fibrous connective tissue. The other portions of tissues all show a similar structure. On section they exhibit a number of rounded areas, which are separated from each other by a small quantity of tissue which has not the yellowish tinge the rounded areas have. These rounded areas, on microscopic examination, are found to be composed mainly of tubular glands lined by columnar epithelium with a small quantity of fibrous and muscular tissue lying between them, that is, are adenomata. The tissue lying between the adenomata has the appearance of normal prostate."

Finally, as regards this case, Dr. Creighton, writing on October 10th, 1903, reports the man in his former vigorous health, retaining his urine for normally long periods, passing with a large stream, and having full control of his bladder.

CASE II.

A. B., a gentleman aged 65 years, sent to me by Dr. Goff, of Dundrum, suffering from prostatic obstruction and stone in the bladder, came under my observation in May, 1901. I then removed by litholapaxy a calculus weighing about 90 gr. I hoped that the prostatic enlargement, which seemed to have been provoked by the presence of a stone and which had begun synchronously with the rational evidence of its presence, would, as we sometimes see, diminish when the cause of irritation had been removed. But it was not so, and eventually worn out by the torment of a catheter life, the patient submitted to operation on September 1st, 1903. He was in a soft leucophlegmatic condition, his health being lowered by constant urinary distress. There was always

Fig. 3.

a large quantity of residual urine, and the secretion could only be kept neutral or slightly acid by the use of drugs and by washing the bladder.

The operation was done on September 1st, 1903. It was rapid and easy, not more than ten minutes being occupied in enucleating the prostate. A little urine was passed by the natural channel on the tenth day and gradually increased until the twenty-first day, when the tube was withdrawn during the day and only used on occasional nights. On the twenty-seventh day the patient was allowed up for a short time. From this on he only passed water every fifth hour, and finally left the private hospital on October 20th, forty-nine days after the operation, the wound being healed and his urinary functions in a perfectly normal condition. The convalescence was delayed by a short intercurrent attack of pneumonia. I saw Mr. A. B. on January 15th, 1904, when he visited me at my house in perfect and vigorous health.

Dr. Earl reports: "The right lobe weighs 1,332 gr., and measures 2½ in. in length; the left lobe weighs 631 gr., and measures 2½ in. in length. The total weight, therefore, is 4½ oz., less 27 gr., and the total circumference a little more than 8½ in. Sections were made from each lobe, and portions adhering to their surface were also examined. The lobes have the general structure of the prostate, but the glandular tissue is more abundant than normal. Some small elevations on the surface of the lobes are composed of glands very much dilated. The tissue adhering to the surface of the lobes is fibrous. It contains no glands; one of the pieces examined contains some of the lining

membrane of the urethra." With reference to the urethral membrane, I was not aware, either during the operation or from the subsequent progress of the case, that it had been injured.

CASE III.

C. D., a gentleman aged about 58, was sent to me in January, 1904, by Dr. M. O'Malley Knott of Castlebar, and Dr. Allman of Westport. He had suffered for four or five years from gradually increasing prostatic irritation, which assumed the form of urethral obstruction about two years ago, and has since been increasing in degree. The urine was faintly acid, and contained a little pus and albumen, but no tubercle. The amount of residual urine was never less than 24 oz. The prostate examined by the rectum did not seem to be much increased in size, but was distinctly larger on the left than on the right side. A No. 12 (E) catheter passed without any resistance. The catheter had not been previously used except on one occasion about a year ago for the purpose of examination. The patient could not pass any urine except when lying down on either side; it then dribbled slowly away. From these conditions I expressed the opinion that a pedunculated growth was present without much hypertrophy, and this was found to be the case.

I operated on January 16th, 1904, feeling some anxiety owing to the low condition of the patient and the shabby state of a somewhat exhausted heart, and found a prostate slightly enlarged on the left side, not at all hypertrophied on the right, and with a small flabby growth $\frac{3}{4}$ in. long, with a broad pedicle, growing from the prostate gland at the right side of the neck of the bladder, and movable in such a way that it effected the maximum of obstruction when the patient was erect, and the minimum when lying down; in the former position its action was valvular. I incised, enucleated, and removed this growth, finding, owing to its small size and flabby consistence, more difficulty in doing so than if the whole of an ordinary enlarged prostate were in question. No necessity existed for attempting to remove the gland itself. It was not the seat of any essential enlargement, and was not nodular. Indeed, I doubt if it would be possible, without inflicting grave injury, to enucleate a prostate in a normal or nearly normal condition as regards size and attachment, and which had not by growth freed itself to some extent from its connexions.

The patient is now (on the eighth day), doing well, and has given no cause for anxiety, except for the twenty-four hours following operation, when the weak state of his heart caused some alarm.

Dr. Earl reports that "the part removed weighs 105 gr., and is an adenoma mainly composed of glands lined by columnar epithelium lying close together, and similar to those forming the prostate."

As the nature of prostatic enlargements seems still to be a moot point, I may be allowed to express my views, founded on an extended observation of cases and specimens. There seem to be three conditions of senile enlargement:

1. A true hypertrophy of the gland without any interstitial growths.
2. The existence upon and within the substance of the prostate of one or more encapsuled tumours, distinct from the substance of the gland proper, and either myomatous or adenomatous in structure. Both forms of neoplasm or one only of them may be present.
3. A mixed condition of true hypertrophy and interstitial growths of the second class. This condition is much the most common. The tumours may eventually reduce the prostate to a state of pressure atrophy, or cause it to become a mere capsule for adventitious growths. It is the latter condition which in all probability has given rise to so much dispute and uncertainty as to the true pathology of enlarged prostate. The third of the figures represents a section of one of the lobes seen in Fig. 2, and shows the appearance of the prostate studded with these growths, which are nearly always of a spherical form, and only depart to more irregular outlines when they become of large size or approach the surface of the prostate.

ADULTERATION OF HONEY.—The Laboratory of the Inland Revenue Department of Canada has issued two brief reports by Mr. Thomas Macfarlane, Chief Analyst, on the results of analysis of honey and of flavouring extracts. Ninety-nine samples of honey were examined: eighty-one were genuine, five were doubtful, and thirteen were adulterated (six with glucose sugar and five with cane sugar). Honey may contain cane sugar either owing to artificial addition or because the bees are fed exclusively on it. Invert sugar may also be added and is less easy of detection. This sort of adulteration is known in Europe, where the product is sold as "Turkey honey" or "table honey." Its manufacture has increased greatly during the last decade. Water may also be added to honey above the amount (20 per cent.) normally present. Only eight samples were found to contain water in excess of 25 per cent. No injurious ingredients were met with in this series of analyses. The same may be said for the flavouring agents, many of which however are artificially coloured.

PROSTATIC ENLARGEMENT AND ITS TREATMENT

BY FREYER'S METHOD.*

BY GILBERT BARLING M.B., F.R.C.S.,

Professor of Surgery, University of Birmingham; Surgeon, General Hospital, Birmingham.

ENLARGEMENT of the prostate is a subject of perennial interest from the frequency with which it occurs, the difficulties it offers to the practitioner, the dangers to life which it brings in its train, and last, but by no means least, from the misery and degradation with which it surrounds the later period of life.

The various operative measures for the relief of this condition hitherto at our disposal have on the whole been either disappointing or surrounded with too great risks, hence the readiness with which new proposals are welcomed. The latest of these, namely, the possibility of complete removal of the prostate by the perineal or the suprapubic route with reasonable safety, and with complete relief to the patients' symptoms, is now earnestly receiving the attention of the surgical world. Numerous observations in man and animals show conclusively that the prostate fails to develop if the testicles are removed in early life, further, that the healthy prostate shrinks if the testicles are removed when adult life is attained. We are familiar with the ingenious arguments of White, of Philadelphia, based on these facts, which led him to advocate castration to relieve the symptoms attendant on the enlargement of the prostate associated with advancing age. The proposal was welcomed with enthusiasm, and often indiscreetly practised, but we are now in a position to estimate fairly the true value of the proceeding. As far as my own experience goes, and I believe I am in accord with most other surgeons, the final results have been disappointing, and castration is now rarely done for prostatic enlargement. It was not without risk to life and to the patients' mental condition, but these drawbacks might have been faced if the relief had been thorough and persistent. Unfortunately this has not been realized, and *a priori* argument falls to the ground in the face of clinical experience which is opposed to it. The adenomatous, fibroid, or myomatous overgrowth does not yield to removal of the testicles as does normal prostatic tissue. The lesser operation of vasectomy has given even more disappointing results, and has equally fallen into disrepute.

Ligature of the internal iliac arteries with the purpose of cutting off the prostatic blood supply and inducing atrophy has been resorted to in a few cases, but the results have been disappointing and not commensurate to the severity of the operative proceedings.

Partial removal of the prostate, in this country so honourably associated with the name of McGill of Leeds, has, on the whole, given the best results attained, but its mortality has not been small; whilst in some cases the relief has not been adequate to the risk run, and in others continued growth of the prostate has eventually nullified the advantage at first obtained.

These are the reasons which induce—nay, which require—surgeons to give a fair trial to methods which promise better results, for unless some more satisfactory method is forthcoming than those briefly alluded to the catheter, with all its drawbacks, will still remain, in the large majority of cases, the main remedy by which relief is given.

Complete removal of the prostate, either via the perineum or by the suprapubic route, is now on its trial; each method has its ardent advocates, and if one listened uncritically to either side it would seem as though the question were sure and settled, the risk to life very small, the recovery after operation uneventful, the final results irreproachable; the conclusion is like that of a child's story book, in which the characters "all lived happily ever after."

In bringing the subject before you to-day I do not intend to contrast the two operations—the high and the low—but to place before you the main facts with reference to cases in which I have removed the prostate by the suprapubic route.

It is necessary to make a few preliminary observations on the parts concerned. Without going minutely into the anatomy of the prostate, there are some essential points to which I wish to call attention as bearing directly upon the question before us. We have to deal with an organ which in healthy adult life weighs about 6 drachms, embraces and is intimately connected with that portion of the urethra next to

* Read at a meeting of the Birmingham Branch of the British Medical Association.