occurred the patient should be compelled to resume work. It would be necessary for this purpose to have a certificate from an ophthalmic surgeon—probably from two surgeons, one representing the workman and the other the employer. There should be no difficulty on the part of the two medical men in coming to an agreement on such a matter.

THE POSITION OF THE EXPERT.

In conclusion, I wish to draw your attention to the position of the expert with regard to fees in these cases. If a surgeon examines and reports on a patient on behalf of applicant or respondent, he may be called upon to give evidence at court on subpoena, and at the usual rate of a guinea a day at county court and half a guinea at police courts. To get any more, a private arrangement must be made with the party who desires the evidence. Should they refuse to make such an arrangement, it is in the expert's power to refuse to make the preliminary examination, or to accept the ordinary fees to medical witnesses allowed by the court.

ON REMOVAL OF THE CRYSTALLINE LENS IN HIGH DEGREES OF MYOPIA, AS ILLUSTRATED IN SIXTY CASES.*

By SIMEON SNELL, F.R.C.S. EDIN.,

Ophthalmic Surgeon, Royal Infirmary, Sheffield; and Professor of Ophthalmology, University College, Sheffield.

Four years ago I contributed a paper to the discussion on the operative treatment of high myopia before the Ophthalmological Society, and which is published in the *Transactions* of that Society. I there dealt with twenty operations. Since then I have increased the number in which I have removed the crystalline lens to about sixty, and it is the experience thus gained that I purpose bringing before your notice to-day.

The deplorable results associated not infrequently with high degrees of short sight are well known. The inconveniences attendant on the use of the necessarily powerful glasses are great. Without their aid, those unfortunately afflicted with high myopia are well-nigh blind, and not infrequently having removed and deposited their glasses they are with difficulty able to find them again. To relieve such cases as here depicted the removal of the crystalline lens has been suggested and performed with success in this country and abroad during the last few years.

Donders regarded the cure of myopia as belonging to the *pia vota*. Before his day the removal of the lens had been suggested, but, writing in 1864, Donders treated the proposal with ridicule, and insisted that such a momentous undertaking "would exhibit culpable rashness." The forty years which have elapsed since Donders wrote these words have witnessed many changes in ophthalmic surgery. Asepticism, which has enabled so many operations to be accomplished with safety in general surgery, has extended its benefits, if in a less degree, to the operative surgery of the eye, and the attendant risks in careful and skilful hands are much less now than formerly.

Within defined limits the removal of the crystalline lens for high degrees of myopia may now, I believe, be regarded as a recognized and valuable method of treatment. By its means the myopia will be either entirely obliterated and the eye become emmetropic or a varying degree of low myopia or a low degree of hypermetropia will result, according to the original degree of shortsight.

A consideration of my cases will bring into view several points which should be noticed. The number of patients operated upon has been 40, the number of eyes 59. It may often be well to be content with only treating one eye, and my rule has been to take the worst eye. Frequently, however, my experience is that after having undergone operation on one eye the patient has been so pleased with the result that urgent application is made for the other to be similarly treated. Domestic servants are numbered among those who have returned after one eye has been operated upon and have been desirous to have the second one done. I am always more willing to accede to such wishes in young adults. In one instance, however, a married woman, Mrs. D., aged 48, I treated both eyes four years ago with most gratifying results.

The number of operations reviewed in this article is 59.

* Prepared for a meeting of the Yorkshire Branch of the British Medical Association.

There were 40 patients; in 21 only one eye was treated and in 19 both eyes were operated upon. Of the 40 patients, 11 were males and 29 were females. Of those who underwent operation in both eyes 13 were females and 6 were males.

One patient was aged 48, but this was the only instance over 40. Of the others, one was 37, another 35, and two 32. Those between 20 and 30 numbered eighteen, as follows: three aged 30, one aged 29, four aged 28, two aged 27, three aged 26, one aged 25, one aged 23, two aged 22, and one aged 21. Under 20 there were seventeen, namely: two aged 19, five aged 18, one aged 17, three aged 15, one aged 14, one aged 13, two aged 11, one aged 10, and one aged 9.

Of the instances in which the degree of myopia is accurately recorded I find that 28 operations were performed on eyes with 20 D and above, namely, 15 for 20 D, 1 for 21 D, 2 for 22 D, 1 for 23 D, 1 for 24 D, 2 for 25 D, 3 for 26 D, 1 for 27 D, and 2 for 28 D. The lowest degree for which removal of the lens was performed was 10 D, and this in one instance only. There was 1 eye for 11 D, 2 for 12 D, 2 for 13 D, 3 for 14 D, 1 for 15 D, 3 for 16 D, 10 for 18 D, and 1 for 19 D. The number, therefore, between the eyes of 10 and 19 D inclusive was 24. It will thus be seen that practically all the operations have been performed on eyes having 12 D and more degrees of myopia, the two exceptions being 10 D and 11 D respectively. Moreover, the greatest number of operations have been for 18 D and upwards, 18 D being the degree in 10 eyes and 20 D in 15.

If the operation be limited, as I believe it should be, usually, to cases with a myopia of 14 D and upwards the total number of suitable cases presenting themselves must always be a comparatively small one. In this connexion I have taken from the casebooks of my private practice the degrees of myopia in 3,162 consecutive patients or 6,324 eyes, a number which should give a fair idea of the proportion of the different degrees of myopia. The cases were roughly divided into those above 10 D and those of 10 D and under. If the division had been made at a higher degree of myopia, which, strictly speaking, it should have been, as, save in exceptional instances, the operation would not be performed for less than 13 D or 14 D, the total number of instances suitable for operation as far as the degree of myopia is concerned would be decidedly less.

Of these 3,162 patients I find that in 1,159 the degree of myopia was equal in each eye. In 2,003 the eyes were unequal in refraction; that is to say, that while myopia was present in one eye it was so to a higher degree in one eye than the other, or in some the second eye may have been emmetropic or hypermetropic, but the greater number would be myopic. Of these 2,003 eyes the degree of myopia was higher in the right eye in 1,029, and in the left in 974. Of the total number of eyes under review, namely, 6,324, there were 320 in which the degree was more than 10 D, or a trifle over 5 per cent. The record of my infirmary patients would, I expect, show a larger percentage of the high degrees of myopia, as those with the low degrees less frequently seek advice there than is the case in private practice.

Extraction was deemed unsuitable, and in no instance has it been adopted. In each case the transparent lens has been needled, and at the end of a week, sometimes more or less, when the lens substance has become sufficiently broken up, an incision with a broad needle has been made a little distance inside the periphery of the cornea, and the softened lens has been allowed to escape, or has been coaxed out with the aid of a curette. The pupil has previously been dilated with atropine, and by keeping the situation of the corneal wound inside the limit to which it dilates, the danger of entanglement of the iris in the wound is materially lessened. This operation is similar, of course, to that frequently performed for zonular and other forms of cataract. In all cases a watch must be kept for increased tension, which is liable occasionally to be set up by the swelling of the lens. In a few of my cases glaucoma has in this way supervened. Removal of the broken-up lens was, however, promptly performed, and was followed by immediate subsidence of the pain and sickness, and with excellent visual results.

When operating endeavour is, of course, made to remove as much of the softened lens matter as will escape without undue pressure on the globe. What remains, however, will undergo absorption, but a further needling may be required. Needling may also be necessary for opaque capsule and it is therefore important to bear in mind that treatment in some cases may be somewhat prolonged. In one of my early instances I was perhaps hardly sufficiently alive to the importance of this and operated rather against time, as the patient was leaving England.

If due regard be paid to asepsis and the use of sublimate or some other antiseptic solution which are essential to eye operations, the dangers attendant on this operation are trifling. The introduction of any suction apparatus is altogether unnecessary, for by gentle coaxing the softened lens matter readily comes away through the corneal wound. It is desirable, however, that after the needling operation the patient be kept under close observation, for, as mention has already been made, in some instances the swollen lens is apt to induce increased tension as signified by pain and sickness, and the immediate performance of the second operation for the removal of the broken-up lens will be indicated. A free opening of the capsule is practised, because it is held that by permitting the broken-up lens to escape into the anterior chamber increased tension is less likely to be induced. This is not, however, the only period that it is necessary to be on one's guard against the onset of increased tension.

In a certain number of cases after the usual operative measures have been completed and the pupil is free, or nearly so, of lens matter, increased tension may occur. In these instances atropine has very properly been used to maintain a dilated pupil whilst the remnants of the lens were undergoing absorption. The atropine may be discontinued and eserine substituted, but it will, in my experience, fail frequently to overcome the glaucomatous condition. The best plan is to tap the anterior chamber, and to repeat if necessary the paracentesis. In the few instances that increased tension has arisen in the circumstances just now mentioned, paracentesis has acted well, and the results in those cases have been among my best.

The operation performed in my cases is similar to that practised for lamellar cataract, and the dangers are the same. A bead of vitreous may present at the wound, but this should very rarely occur if care has been taken to limit the tearing of the capsule to the anterior capsule, which should readily be done, and also if too great pressure on the globe is avoided when the lens is coaxed out by the curette.

My practice has been to indicate to a patient the advantages which may be gained by operative measures, but at the same time to point out that to achieve the benefits he or she must be prepared to face the risks which, however, are not great. The decision is left to the patient.

Some authors have alluded to the danger of detachment of the retina resulting from removal of the crystalline lens for high degree of myopia. I have no experience of extraction of the lens in these cases, and cannot therefore say whether it is a danger by that method or not. All my cases have been treated in the manner I have described, and certainly detachment of retina is not a danger of moment in my experience. I recollect only two instances; one was mentioned at some length in my paper before the Ophthalmological Society.

CASE I.

The patient was a young woman, aged 25. The lens was needled, and subsequently evacuated in November, 1897. The result up to May, 1898, was excellent, myopia of 22 D had gone down to 2 D, and vision at this time was $\frac{1}{2}$. Shortly after this, however, owing to the serious illness of a near relative, the patient had very arduous night and day nursing thrown upon her. Vision rapidly deteriorated, and on June 8th in was reduced to $\frac{1}{20}$. Besides some films in the vitreous, no gross changes were noticeable in the eye. Vision remained at this point or a little worse for some months, but she had not been under examination for some time. It should be mentioned that when the lens matter was evacuated a small amount of vitreous escaped.

CASE II.

The second was that of a domestic servant. Both eyes had been operated on with excellent results. A year or more afterwards detachment of retina occurred in one eye. At this time she was in very feeble health. The other eye has remained good as far as I have heard.

These are the only instances of which I have either record or recollection in the 59 eyes which have been operated upon, and many of my cases have been under observation more or less for several years. Myopic eyes—especially the high degrees which are dealt with in this paper—are, it is well known, prone to detached retina. They are cases in which a sword of Damocles is always hanging over their heads. Even, therefore, if the percentage was much larger, as it has been in some published records of instances in which detached retina has occurred after varying periods subsequent to the operation, it could hardly be regarded as discrediting the operative methods here advocated.

The results of a successful operation to the patient will be that vision is as good or better without glasses for distance as it was previously with their assistance. The myopia will have disappeared or be present only in low degree, or there

may be weak hypermetropia, according to the amount of myopia originally existing. It would be very unusual for a patient not to express his or her gratification with the change that has been brought about. There are few subjects in ophthalmic surgery that elicit such expressions of gratitude as do these operations for the relief of myopia.

I give here an extract from a letter received from a patient operated upon some few months since. She was a teacher, aged 28, and the degree of myopia in each eye was 28 D. The right eye only was treated. Before operation, with correction with -28 D, $V = \frac{1}{8}$; after operation, without the aid of glasses, $V = \frac{1}{2}$.

The benefits received from the operations are so numerous that I hardly know how to specify them. The operations, of course, to me appear nothing short of a miracle. The fact that I had worn exceedingly strong glasses for twenty-two or twenty-three years emphasizes this. Prior to the operation objects a few yards distant were quite unrecognizable; clear vision was only obtainable an inch or two from the eye. The uncertainty of locating the outline of objects caused me great trouble, inconvenience, and, at times, even fear. For instance, in crossing the road the edge of the pavement was always uncertain. The same may be said of steps to vehicles, stairs, etc. Nothing of that uncomfortable uncertainty remains, all things are now clear. I see clearly, distinctly, and have quite a long range of vision; not only that, but the continual tired, aching feeling, and fearfully hot sensation have entirely gone; and it is saying a great deal when I say that whereas for many years my eyes have been an ever-present feeling, I now sometimes forget that I have eyes, so very natural and painless are they.

[Since this article was written I have operated on four additional cases, all successful; three females, aged 26, 15, 9, one boy, aged 10; the degree of myopia of each in this order being 18 D, 14 D, 18 D, and 15 D.]

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL, THERAPEUTICAL, PATHOLOGICAL, Etc.

ALLEGED POISONING IN AN INDIGO DYE WORKER. THE following case of poisoning attributable to emanations from the contents of a mixture used in the process of calico-dyeing may be of interest;

A man, aged 37 years, came to my surgery about 6 p.m. on Christmas Eve, complaining of faintness and nausea. He said that about an hour previously, while preparing a dye mixture, he was suddenly seized with nausea, followed by repeated vomiting and prostration. He did not get better when out in the open air, so he came for advice. The man was of sturdy build, but looked haggard; he was much agitated, and in dread of death.

The first thing that struck me was the changed appearance of his skin. His face was covered with a reddish rash, the mucous membrane of his mouth had a brownish discoloration, and there were two rounded black spots on the cheek. The rash extended to the general surface of the body, and scattered over the trunk, upper arms, and thighs were groups of dark spots similar in character to those on the face. They were each a little smaller in size than a shilling piece, quite round, very black in the centre, and shading off to a grey towards the circumference. They were mostly in groups of five, arranged lengthwise, but were each quite distinct—not raised, and not disappearing on pressure.

The pulse was 90 per minute, the temperature 98.8°, heart and respiration were normal. He was perfectly clear in all his statements. His wife who accompanied him told me that he was temperate and careful as to his food and habits. I could not examine the contents of the stomach as the repeated vomiting had emptied it, but was told that he had not taken anything injurious by the mouth in the shape of food or drink. His pupils reacted normally to light, the knee-jerk was slightly increased, otherwise nothing abnormal was to be noticed.

The sample of urine given me was normal. The man said he was sure he was not going to live, as some years ago a similar case happened at their works, which ended fatally.

It was evident that the attack was caused by the inhalation of some poison which emanated from the dye mixture the man prepared, but I could not make sure what particular poison it could have been.

The treatment ordered was rest in bed and an aperient of magnesium sulphate followed by bismuth. The patient improved rapidly. The rash became fainter on the following day, and he was able to retain fluid nourishment without any nausea. On the third day from the time of the onset the