A CASE OF SYPHILITIC CONVULSIONS PRECEDED BY MARKED SOMNOLENCE OF LONG DURATION.

By WILLIAM B. BENNETT, M.R.C.S.ENG., L.R.C.P.LOND.,
Surgeon, St. George's Hospital for Diseases of the Skin, Liverpool.

THE following case of syphilitic convulsions is noteworthy on account of the remarkable tendency on the part of the patient to sleep during the five months preceding the convulsive attack. As will be seen from the history, this somnolence apparently replaced the characteristic headache, which is met with so frequently in these cases as to be almost diagnostic, and which usually produces sleeplessness. Other prominent symptoms were profound stupor and motor aphasia.

HISTORY.

The patient was a married man aged 34. On Saturday, April 26th, 1902, at 7 a.m., the patient according to his usual custom roused his wife, with whom he was sleeping; she got up, dressed, and left the room, having noticed nothing unusual about her husband. At 8 a.m. the man came down to breakfast; but, to his wife's surprise, he was quite unable to speak, although he made great efforts to do so. He then sat down, and with considerable difficulty ate a small bit of toast and drank a cup of tea. He understood what was said to him, but could make no answer. At his wife's request he then returned to bed. At about 9 a.m. his little daughter told her mother that her father was ill; the mother immediately ran upstairs, to find her husband quite unconscious, "making a snoring noise," and with some froth about his lips.

I was immediately sent for, and saw him a few minutes later. He was then lying on his back in a comatose condition, breathing stertorous, teeth tightly clenched. At 10 a.m., whilst I was standing by his bedside, he had an epileptic convulsion, which lasted two or three minutes, and resembled an ordinary epileptic fit, with the exception that the convulsions started in his right arm, leg, and side of face. He was generally convulsed, but most strongly on the right side; as a natural result, he turned over on to his left side, on which he lay throughout the fit. I threw 5 gr. of calomel and 30 of potassium bromide into his mouth, and think they were swallowed. Whilst his wife was out of the room I examined his genitals, but could find no scars or other signs of venereal disease. Between 11 and 2 o'clock he had two more similar attacks, which made three, probably four, altogether. In the interval between two of the fits he once got up, and walked about the room for a few minutes, being in a state of great excitement, but was got back to bed without any difficulty. Whilst in bed he was in much the same state as when I first saw him: eyes tightly closed, teeth firmly clenched, corneal and skin reflexes present, and quite unconscious of his surroundings. It was only with the greatest difficulty that his teeth could be forced apart for the purpose of introducing powders or fluids. In the evening I gave him 5 gr. more of calomel, having previously put him on 15 gr. of potassium iodide, with 1 drachm of liquor hydrargyri perchloridi every three hours. I also gave his relations a favourable prognosis. In the evening he got out of bed and emptied his bladder and rectum quite naturally, getting back again without any assistance, and displaying no bodily weakness. He seemed, however, quite unconscious of his surroundings, and did not attempt to speak, falling at once on his return to bed into his former heavy sleep, which continued throughout the night. At no time had he any incontinence except during the convulsions, when he passed his water unconsciously.

On Sunday morning he was in much the same condition, sleeping heavily and only waking to empty bladder and rectum. His bowels were freely moved both in the night and again in the morning. There was still great difficulty in feeding him, all medicines and nourishments having to be poured into his mouth after forcibly separating his tightly clenched teeth. In the afternoon he was distinctly clearer, and lay for a few minutes with his eyes open: he then noticed a relative whom he had not recently seen, and said to him, "Hullo, Smith," these being his first words since he rose the previous morning. He at once relapsed into his former state of stupor, from which he could, however, be partially roused. Later in the day he said "Yes" and "No" a few times when spoken to, but not intelligently: feeding was still difficult, but less so than formerly. During Sunday night he slept soundly.

On Monday morning he was distinctly clearer and could say a few words. He also endeavoured to dress himself, and seemed very disappointed at not being permitted to do so. The dose of potassium iodide was now reduced to 15 gr. with a drachm of liq. hydrarg. perchlor. four times a day.

On Tuesday he was able to carry on a conversation for a few sentences, but no more, not being able to fix his mind on anything for more than a minute or two.

On Wednesday he was much clearer and daily increased in strength so rapidly in mind and body that he was allowed out of bed on the Saturday for a short time, when he was physically weak but quite rational. Owing to his mouth becoming very inflamed and ulcerated the liq. hydrarg. perchlor. was omitted from his medicine, and for the next few weeks the iodide alone, 1 drachm daily, was given. His mouth soon recovered and in about a week was quite well.

On Tuesday, May 6th, as he seemed quite himself, he went out for a short walk, and on the following Tuesday, the 13th, went away for a week to the seaside, after which he returned to his business. Early in June he was ordered 10 gr. of potassium iodide with 1 drachm of liquor hydrarg. perchlor. thrice daily. This he was able to take without any ill-effects. He told me that he had a distinct recollection of coming down to breakfast and being unable to speak, but that after he returned to bed he remembered no more until he saw and spoke to his relative the following day, after which for a day or two only a few incidents appear to have left any impression on his mind.

The history previous to his attack, which I have since obtained, is as follows: He had always been healthy and well, never having been seriously ill before; he had a chancre in the usual position eight years ago, for which he was treated by a doctor for one month; he was not aware that it was necessary to undergo any further treatment, and had had no rash, sore throat, or secondaries of any description.

About five months before the present illness he began to feel drowsy at all hours, falling off to sleep with great ease. During the last two months this tendency to sleep had become most pronounced. He would return at 7 in the evening, eat his dinner, and then sleep before the fire till 10, when he went to bed and at once fell sound asleep, remaining so until he got up at 8 next morning. Whilst riding to his place of business after breakfast on an omnibus he felt sleepy, and during the day was often found fast asleep in his office. He never had any pain in his head or elsewhere, neither did he feel any discomfort of any kind besides this one desire to sleep. These facts were corroborated by his wife, who also told me that she noticed no change whatever in his disposition or behaviour.

The subsequent history has been quite uneventful. The drowsiness, which completely left him a few days after his convulsive attack, has never returned, and he only sleeps a normal number of hours at the proper time.

For some months his memory was a little defective. He would forget some incident, either trivial or important, which had happened the previous day or less recently, until reminded, when he at once remembered. This gave him some trouble, as it happened frequently, but it passed off completely, and in July, 1903, he was perfectly well in all respects.

I think this well-marked prodromal somnolence must be very rare, as I have not been able to find the report of any similar case. The symptom is, of course, known, since Gowers¹ writes of "considerable mental dullness or a somnolent condition lasting for weeks," as being an occasional premonitory symptom in cerebral thrombosis.

Aldren Turner² also writes of a "tendency to somnolency, so that the person drops off to sleep while at work," as being sometimes associated with a change in the character and disposition of a patient suffering from cerebral syphilis.

Buzzard³ has recorded several cases of prolonged somnolence associated with hemiplegia; but in his cases the somnolence did not precede, but accompanied or followed the acute attack, and, moreover, produced great mental dullness.

DIAGNOSIS.

Quite apart from the history of chancre, the gradual onset, together with the complete and rapid recovery under anti-syphilitic treatment, clearly proves that the lesion was of a syphilitic nature.

NATURE OF LESION.

The most common syphilitic disease of the brain is that known as obliterative endarteritis of the cerebral arteries—a disease which slowly narrows the cavity of the vessel to a

greater extent than does atheroma, but produces very similar pathological effects.⁴

Heubner is quoted by Cornil⁵ as attributing the somnolency to the narrowing of a number of arteries, in consequence of which there is congestion and defective oxidation of the brain substance. Motor aphasia would be produced by occlusion of the first cortical branch of the middle cerebral artery.

According to Osler,⁶ transient aphasia with perfect recovery may occur in advanced arterio-sclerosis, but the explanation of this fact is not easy.

Convulsions are very frequently caused by cerebral syphilis, and both Fournier and H. C. Wood⁷ have laid great stress on the value of this symptom in diagnosis when occurring in persons over 30. They may be caused, like aphasia, by advanced arterio-sclerosis,⁸ but are decidedly rare in atheroma, being more often due to meningeal growth or chronic inflammation.⁹ This may, of course, have been present in the above case, in addition to the more obvious endarteritis, but had it been I do not think the patient would have been so completely free from headache of some kind or another.

I am therefore inclined to think, from the foregoing considerations, that the lesion in the above case was extensive syphilitic endarteritis of the cerebral arteries, without thrombosis or embolism, but possibly with spasm of the arteries.

The most interesting feature about this case is, of course, the preceding somnolence, but the entire absence of the more common symptoms—for example, headache and hemiplegia—together with the apparently complete recovery of the patient, is not without interest.

REFERENCES.

- ¹ Gowers's *Diseases of the Nervous System*, 2nd edition, vol. ii, p. 432.
² Clifford Allbutt's *System of Medicine*, vol. vii, p. 682. ³ *Lancet*, June 7th, 1879. ⁴ Gowers, vol. ii, p. 429. ⁵ Cornil's *Syphilis*, p. 317. ⁶ Osler's *Principles and Practice of Medicine*, 4th edition, p. 775. ⁷ *Ibid.*, p. 246.
⁸ Gowers, vol. ii, p. 734.

BILATERAL NEURITIS OF FIFTH AND SIXTH CERVICAL NERVES.

By C. H. CATTLE, M.D.LOND., M.R.C.P.,

Honorary Physician, Sherwood Forest Sanatorium for Consumption;
 Assistant Physician, Nottingham General Hospital.

THE following case, while presenting some interesting points in etiology and localization, also shows that the results of brachial neuritis are not always so favourable as in the cases recently reported by Dr. Simon.¹ For the notes I am indebted to Mr. Edwards, House-Physician at the Nottingham General Hospital.

E. S., a coal miner, aged 29, was sent to me by Dr. Nesbitt, of Sutton-in-Ashfield, and was admitted into hospital on July 16th, 1903, having previously attended for about a month as an out-patient. He was a steady, industrious man, non-smoker, and almost a total abstainer. No evidence or history of syphilis. His previous health had been good and the family history was unimportant. For a few weeks before the illness began he had been extra hard worked, and worried about business and family matters. His appetite failed and he slept badly. He persisted in working, although his wife begged him to take rest.

Since May 25th he had complained of very severe pain in the root of neck on both sides, spreading over the shoulders and down the arms. This still continued when I first saw him, about a fortnight later. There was much tenderness on pressure over the cervical nerves on each side of the spine. Power of abduction of the left arm was lost; the other movements of the shoulder and elbow on both sides could be performed, but feebly. Flexion of the elbow was impaired, but not lost. The loss of power was greatest on the left side, and this has been the case throughout. (He is left-handed.) Tenderness was not marked over the main nerves in the limbs. Sensation was distinctly blunted over the cutaneous supply of the left circumflex nerve (both to pain and tactile sense), and later on there was a slighter amount of impairment on the right side. The pupils were equal and reacted normally. At no time have there been symptoms pointing to disease of the brain, cord, or membranes.

After admission on July 16th pain rapidly diminished in the right arm, and power of movement returned. In the left, also, the pains gradually diminished, and sensation on the outer side of both arms became normal. Both deltoids were wasted, the left being the worst. This ultimately became extremely atrophic. The left supra-spinati and infra-spinati were somewhat atrophied. At this time all the upper arm muscles, except the left deltoid, were stated to react to faradism. In the left deltoid there was absolutely no voluntary power, and with galvanism the muscle gave reaction of degeneration. He remained in hospital about a month, and spent about a fortnight at the Cedars Convalescent Home, galvanism being persevered with the whole time.

On September 5th the right deltoid had quite recovered, while the left remained powerless, and the horizontal measurement round the body

of the muscle was $1\frac{1}{2}$ in. less than on the right side. The left spinati muscles were also atrophied.

On December 16th the deltoid had recovered a certain amount of voluntary power, and the difference in measurement was less than before—namely, $\frac{1}{2}$ in. The muscular wasting, however, was now more extensive than had hitherto been noted. Besides the two spinati on left side, the right serratus magnus and infra-spinatus were completely atrophied. None of these reacted with either current. The left deltoid, notwithstanding improved nutrition and return of voluntary power, still gave the degenerative reaction, and did not react to faradism.

The etiology of the case appears to resemble that of Dr. Simon's first case, the only ascertainable causes being hard work, mental worry, and a low state of health. Possibly continued work with the arms raised above the head might cause injurious pressure on the nerves by exciting contractions in the scalenus medius muscle which they pierce. An interesting feature of the case is the bilateral distribution of the muscular paresis and wasting. In two of Dr. Simon's cases, and one of those reported by Drs. Harris and Low,² all like the present, spontaneous or non-traumatic cases, one limb only was involved. Falls on the shoulder, sprains, or other mechanical force are the commonest causes of neuritis of the cervical nerves. Carrying weights on the shoulder is a common cause of this affection, the nerve to the serratus being particularly exposed to pressure. Thus I lately had an out-patient under treatment with the interesting condition of bilateral wasting and paralysis of the serratus magnus, due to pressure on the nerve produced as above mentioned.

The muscular weakness at the commencement of the illness corresponded pretty closely with Erb's type of upper-arm palsy. This is dependent on lesion of the fifth cervical nerve, the muscles most commonly affected being the deltoid, supra-spinatus and infra-spinatus, biceps, brachialis anticus, and sometimes the supinator, longus and brevis.

In the case under notice the flexors of the elbow, though distinctly weakened in the first instance, rapidly regained their strength and tone, while the chief stress of the disease fell on some of the shoulder muscles. The sixth cervical nerve may also have been inflamed to a slighter extent, especially on the right side, on which the serratus magnus is atrophied. The nerve to the serratus comes from the fifth and sixth cervical before they unite to form a common trunk, and sometimes also in part from the seventh nerve. The distribution of the fifth cervical nerve has been pretty fully worked out by different observers, and a short recapitulation of what is known or surmised may prove useful. After receiving a branch from the fourth, the fifth cervical supplies the nerve to the rhomboids, probably deriving its filaments from the fourth, as these muscles were not affected on either side in this case. Then comes a branch which unites with a branch from the sixth cervical to form Bell's nerve (for the serratus). Lower down the nerve, according to Drs. Harris and Low,³ can be divided into an upper and lower fasciculus. The former, after giving off the supra-scapular nerve to the supra-spinati and infra-spinati, joins the posterior cord of the brachial plexus, and furnishes the circumflex nerve, which supplies the deltoid, the teres minor, and the skin on the outer side of the upper arm.

The lower fasciculus forms the greater part of the musculocutaneous nerve, from the outer cord of the plexus, supplying the biceps, brachialis anticus, and skin of the outer side of the forearm. It will be seen that the main incidence of the paralysis in this case corresponds on the left side with the assumption that the upper fasciculus of the fifth cervical nerve was the portion chiefly affected. On the right side the only portion of this bundle of fibres which has not recovered is the nerve to the infra-spinatus, while the paralysis of the serratus magnus shows a picking out of some of the fibres of the nerve immediately after its emergence from the spinal foramen.

Paralysis of the serratus magnus gives rise to a characteristic deformity, for when the arm is moved forward the lower angle of the scapula is rotated inwards, and its posterior edge projects like a wing. As the scapula is rotated and fixed by the serratus when the arm is raised above the horizontal position, its paralysis considerably weakens this movement. This man, notwithstanding wasting of the serratus on one side and only partial recovery of the deltoid on the other, has no difficulty in elevating the arms above the head. The trapezius normally co-operates with the other elevators of the arm, and can to some extent supply their place when paralysed. The spine of the left scapula in the patient under consideration is prominent, and the supra-spinous and infra-spinous fossae unduly hollowed from wasting of the spinati