fective is at once destroyed. Attempts have been made to obtain artificial sponges which could be sterilized by heat. But those which have been tried have been hard and harsh and deficient in absorptive properties.

NOTES AND REFERENCES.

¹ Carl S. Haegler, Basel, 1900. *Handreinigung, Handdesinfection und Handeschutz*, gives a complete bibliography up to date. ² See Haegler, p. 27. Also a valuable essay by Kronig and Blumberg, *Beiträge zur Handdesinfection*, Leipzig, 1900. Also Poter, *Die chirurgische Asepsis der Hände*, Berlin, 1907. ³ Sarwey and Paul, *Arch. f. klin. Chir.*, 1900, p. 463, etc. ⁴ This excellent paper is in the BRITISH MEDICAL JOURNAL, October 17th, 1895, p. 1109. See also Lauerstein, *Munch. med. Woch.*, July 20th, 1902. ⁵ ABST. EPITOME OF BRITISH MEDICAL JOURNAL, October 25th, 1902, par. 253. ⁶ See Stewey, p. 465. ⁷ The house-surgeons were Messrs. Jennings, Pinker, Scott, Farncombe, Rose, and Wenham. I am much indebted to these gentlemen for their help. ⁸ See reports in BRITISH MEDICAL JOURNAL. ⁹ *Lessons in Disinfection and Sterilization*, p. 93. ¹⁰ As given in *Aseptic Surgery* 2nd ed., p. 177. ¹¹ J. Lynn Thomas, On the Value of Gloves in Operative Work, BRITISH MEDICAL JOURNAL, January 21st, 1899, p. 140. ¹² BRITISH MEDICAL JOURNAL, May 28th, 1892. ¹³ From March, 1900, to December, 1903. ¹⁴ Lucas-Championnière, *Cure Radicale des Hernies*, Paris, 1892. ¹⁵ *Wien. klin. Woch.*, October 1st, 1896. ¹⁶ ABST., see EPITOME OF BRITISH MEDICAL JOURNAL, December 5th, 1896, par. 460. ¹⁷ An Address on the Organization of Aseptic Operations and some of the Causes of Failure, BRITISH MEDICAL JOURNAL, February 24th, 1900, p. 429. ¹⁸ *Lancet*, January 9th, 1904, case iv, p. 93. ¹⁹ BRITISH MEDICAL JOURNAL, September 17th, 1893, p. 802. ²⁰ *Johns Hopkins Hospital Reports*, vol. vii, Nos. 5-9, Baltimore, 1899. ²¹ *Loc. cit.*, p. 225. ²² *The Radical Cure of Hernia, Hydrocele, and Varicocele*, by C. B. Lockwood, Edinburgh and London, 1893, p. 153. ²³ New sponges are (1) shaken and beaten to get rid of sand; (2) soaked in solution of hydrochloric acid and water to remove shell and coral (1 drachm of strong hydrochloric acid to 1 pint of water); (3) washed out in alkaline water, and subsequently in sterile warm water (1 drachm of washing soda to 1 pint of water); (4) twelve hours in cold solution of sulphurous acid (1 in 5), which is washed out with sterile water.

THE TREATMENT OF APPENDIX-ABSCESS IN THE RECTO-VESICAL POUCH.

By EDMUND OWEN, F.R.C.S.,

Consulting Surgeon, St. Mary's Hospital and Hospital for Sick Children, Great Ormond Street, London.

IN the autumn of last year some interesting and valuable lectures were published upon inflammatory disease of the appendix, in the course of which special reference was made to the treatment of those cases which were complicated with intra-pelvic suppuration.

It was affirmed in one of the lectures that it was always better to drain a localized pelvic abscess from the rectum or vagina. That by either of these methods the patients were well in a fortnight, and never in the hospital more than three weeks; and, moreover, that they thus ran no risk of ventral hernia. But that if, on the other hand, the abscess were drained by a suprapubic incision the wound took six weeks at least to heal, and often much longer.

The object of this paper is to call into question the propriety of this teaching, and to affirm that the safest way of dealing with an intrapelvic appendicular abscess is by abdominal section. I will briefly cite an illustrative case which was lately under my care:

A gentleman, aged 45 years, was sent to me on August 17th, 1903, with a history of intermittent trouble with defaecation and of a feeling of fullness and weight about the neck of the bladder. His doctor, who is a most competent practitioner, wrote that last May the patient had been treated by some one else (he did not say where, or by whom) for obscure abdominal pains which were centred in the hypogastric region. At that time, he said, no rectal examination had been made, but when the practitioner had introduced his finger into the rectum he had found a tumour which felt like an enormously enlarged prostate gland. The patient told me that he had constant pain on micturition and defaecation, and that occasionally there was a "discharge" from the urethra. He denied all possibility of venereal infection. For several years he had had obscure abdominal pains, and he had become dyspeptic to such an extent that he was afraid to eat meat. Three months previously he had had severe pains in the hypogastric region, for which he was kept in bed. He told the doctor who was attending him that he thought he must have got "appendicitis," so great was his distress. But the doctor, finding that there was neither swelling nor tenderness in the right iliac fossa, put down the distress to the after-effects of influenza. He applied fomentations over the hypogastric region. In due course the acute attack passed off, and the patient was able to resume work. But he was very miserable and depressed, and was steadily losing weight. He also told me that he felt as if he had a cricket ball near the rectum, and that he could never obtain a proper faecal evacuation.

The finger introduced into the bowel at once came upon a hard, tender swelling at the base of the bladder. It was about the size of a small hen's egg. It was not prostatic, but it was so close to the prostate gland, and so intimately attached to it, that there was little wonder that it had recently been likened to an enlarged prostate. No lymphatic glandular invasion was anywhere discoverable, but I was afraid lest the tumour might prove to be malignant. In the course of a consultation

a surgeon hazarded the speculation that the tumour might be hydatid, but we agreed that it was impossible to be sure of its nature without operation, and that the case most certainly needed operation.

I therefore opened the abdomen in the middle line, and, working cautiously amongst coils of closely-adherent intestine, found at the very bottom of the recto-vesical pouch a tightly-filled abscess, the walls of which were composed of the rectum behind, the prostate below, and the bladder in front, whilst a kinked piece of the colon formed its roof. To the right side of the abscess, and firmly glued to the base of the bladder, was the end of the appendix, inflamed and ulcerated. Having removed the appendix, and having thoroughly wiped out the abscess cavity with dry swabs of absorbent wool, I passed a very large drainage tube down into the recto-vesical pouch and brought the other end out a little below the umbilicus. Though the peritoneum showed itself proof against infection by the bacilli coli, the surface-wound became infected about a fortnight after the operation, and so severe was the attack of cellulitis caused thereby that some large branches of artery in the epigastric region burst out bleeding on two occasions, and the patient was near dying from secondary haemorrhage. As it is, however, he is now in the North of Africa completing his convalescence.

He was fully apprised of the fact that we had been throughout in uncertainty as to the diagnosis, and was entirely in agreement with us that he had better be operated on. After the operation, when we told him that the outlook was quite cheerful as the tumour had proved to be an abscess in connexion with a diseased appendix, which we had also removed, he said, "Ah, when I was so bad I told the doctor that I believed I had appendicitis!" But, inasmuch as the present time everybody—lay and professional—talks familiarly about "appendicitis," I am afraid that I cannot award this patient full credit for the correctness of his diagnosis. Nor would I blame his doctor who apparently had declined to accept it, for, so far as I can ascertain, the pain and tenderness had never been in the right iliac fossa, but always in the pelvic and hypogastric region. My explanation of the case is that, some indefinite time ago, he had probably had a milk attack of inflammation, which fixed the tip of a long and wandering appendix in the depths of the pelvis, and that a few months ago there was recrudescence of the inflammation with severe hypogastric pains, and with the formation of the abscess. Had he been operated on just after his first attack (whenever that was), he would, I feel sure, have been spared a large amount of suffering, and would have avoided the very serious risks which were inseparable from the more serious operation.

In the case of suppuration in connexion with an inflamed appendix there is no telling in what direction the exigencies of the operation may not lead the surgeon. He may explore the iliac fossa and be disappointed at finding that it contains neither appendix nor caecum, and it may take him some time to discover any part of the large intestine. Eventually, maybe, he comes across a piece of large intestine which he takes to be the right colon, and, guided by one of its muscular bands, he looks onwards for the appendix, which he may discover in due course but after much delay. Sometimes he finds the appendix—as in this case—in the depths of the pelvis. Not infrequently he has to continue the abdominal wound up to the costal region, and may then discover it hidden away behind the liver with which it had formed an illicit connexion. There is no organ in the body which is so regardless of anatomical proprieties as the appendix. Man ought to be for ever grateful that it is not one of a pair!

There are two grave objections to opening an appendicular abscess through the vagina or rectum. The first is that the operator cannot tell exactly what he is doing. He is operating in the dark, as it were, and for aught he knows the instrument which he is using in his attempt to reach the pus may be traversing a piece of small intestine which has been emptied by pressure, softened by inflammation, and glued by adhesions in a faulty position. The second objection is that his operation may be followed by a faecal fistula which absolutely refuses to close. It would be a real calamity if by chance a wounded piece of small intestine persisted in draining its contents off into the rectum or vagina, or if, the large intestine being implicated, faecal evacuations continued to take place through the vagina. Of course there is a risk of faecal fistula following a straightforward suprapubic opening of a pelvic abscess. But in this case the surgeon would probably see exactly what was amiss when he had the abdomen open and the piece of bowel between his fingers, and he could deal with the condition then or later on. Supposing that he thought it advisable to defer attempting to close the intestinal perforation, he could at least bring it up to the abdominal wound and fix it in a position where it would at any time be readily accessible.

Another strong argument in favour of the abdominal route

is that, by adopting it, the surgeon may be able not only to clear out the abscess, but to remove the diseased appendix. It would by no means follow that because an abscess had been evacuated by a blind, vaginal puncture there would be no further suppuration; and it would be quite easy to believe in the probability of the existence of oft-recurring suppuration in connexion with the continued presence of the diseased appendix; and no operator, I should imagine, would propose resection of the appendix by the vaginal route. Then if, as often happens, the chief purulent focus is associated with secondary abscesses amongst the adherent coils of intestines they can all be thoroughly dealt with by abdominal section, whilst a rectal or vaginal operation would leave them untouched. The question of the length of stay in hospital as an argument in favour of the vaginal or rectal operation is not worthy of consideration by the side of the great questions of thoroughness and safety. Indeed, it appears to me that the recommendation to treat an appendicular intrapelvic abscess by the rectal or vaginal route is as unsound as it would be to advise that a pyosalpinx should be dealt with in that manner.

Having thus made a very definite recommendation in connexion with the treatment of appendicular intrapelvic abscess, I now venture to proceed to draw a moral in connexion with the subject of inflamed appendix, which certainly is not foreign to the subject of this paper. It is this: that if an individual has had one unmistakable attack of appendicular inflammation the appendix had better be resected as soon as the inflammation has quieted down. This recommendation comes under the head of the prophylactic treatment of intrapelvic abscess. If the appendix had not been in the pelvis there would have been no abscess there. The appendix was not there originally. It was probably dragged there by the adhesions which formed about it at the time of its first inflammatory attack, and, if it had been resected as soon as that attack had passed off, there would have been no intrapelvic abscess, and the man would have been saved an enormous amount of risk and worry.

I am quite prepared to find that a good many surgeons are ready to maintain that it is rather precipitate to operate after a first attack, which may have been only a transient catarrhal affair. If one could but look into the abdomen and ascertain the exact state of the appendix, it would be an easy matter to say that one should not advise operation after a first attack. This is, of course, equally desirable and impossible; without making an opening into the abdomen, it is absolutely impossible to know in what condition the attack of inflammation has left the appendix. It may have left it none the worse, all pain, tenderness, and swelling having passed entirely away, in which case its removal would, of course, be superfluous. But, in my experience, this happy state of things is not of frequent occurrence, one attack being generally the precursor of other attacks.

One definite result of the inflammation may be that when the attack passes off it leaves the appendix permanently glued to adjacent or neighbouring viscera—the caecum, colon, bladder, or ovary. This may, by chance, give rise to no inconvenience whatever, but if the tip of the appendix happen to be fixed whilst the rest of it up to the base is free, it is apt at any moment to occasion an acute intestinal strangulation. The presence of an appendix which is fixed only at the extremities is a most dangerous object in the abdomen, and it needs but a slight attack of inflammation to cause the tip of it to acquire firm adhesions. If the appendix contracts adhesions in its entire length this element of danger does not arise. Constantly, when operating in the abdomen for conditions other than those which are due to an inflamed appendix, one has come across the results of an antecedent, and perhaps long-forgotten, attack of inflammation, the appendix being closely adherent to some serous surface. But as this is doing no harm the surgeon passes by it without further heed. But neither is this, in my experience the way in which the clinical history of an attack of inflammation of the appendix ends. As I have said, one attack is generally the precursor of other attacks, and if only the appendix could have been taken away after the first of them, how much suffering and risk would have been saved the patient and how much anxiety the friends.

Supposing that a first attack of inflammation quiets down, and the appendix is not then resected, there may be no further trouble from it whatever, and the individual may be as useful and vigorous a member of society as he was before. Or, after a certain time, there may be a recrudescence of the disease, on the happy subsidence of which, few surgeons, I

should think, would willingly fall in with the desire expressed by the patient or his friends that operation should be still further postponed. If, however, the resection is not only deferred but altogether declined, attack may follow attack; and if, by good fortune, the patient is not carried off by some unlooked for complication, at least he may have his activity and comfort materially interfered with. Then, at last, when there is obviously no chance of further escape from operation, the surgeon's work is possibly rendered extremely difficult, whilst the risks of the operation, the length of time needed for convalescence from it, and the chance of a weakened abdominal wall in the future are all increased.

Many of us, I suppose, have had friends, patients, or acquaintances whose lives have been sacrificed, or whose happiness has been clouded, or whose prospects in life have been seriously damaged, by the dyspepsia, symptoms, by the aches and pains, or by the downright distress which have been caused by the continued presence of an appendix which has been the subject of recurring attacks of inflammation.

The more that I see of appendicular disease, of the danger of delay in operating, of the comparative slight amount of risk which the early operation involves, and of the great sense of relief and freedom from anxiety which it affords, the more fully am I convinced that the patient had best take his courage in both hands and submit to a prompt resection even after a first attack.

THE DIMINISHING BIRTH-RATE AND WHAT IS INVOLVED BY IT,

BEING AN ABSTRACT OF THE INAUGURAL ADDRESS DELIVERED BEFORE THE BRITISH GYNAECOLOGICAL SOCIETY.

By J. W. TAYLOR, F.R.C.S. ENG.

Professor of Gynaecology, University of Birmingham; President of the Society.

The address was divided into four parts.

I.

The first was devoted to a critical study of the statistics of the subject during the last twenty-five years. The birth-rate of the United Kingdom was compared with that of other countries both east and west, and it was shown that during the last twenty years no other nation had sustained so great a loss as we have in this definite period of time.

The marriage-rate in the United Kingdom was then considered, and found to be slowly rising, so that the birth-loss must be due to causes operating in the married life of our population.

The value of the population of to-day was then compared with that of twenty years ago, and it was found that in juvenile criminality, in mental disease and brain weakness and in alcoholism, the restricted population of the present day compared unfavourably with the freer population of a former generation.

A tentative effort (over a limited field) was then made to compare the general culture of the more intellectual classes of the two generations, and this (so far as it went) was found to work out disadvantageously to the present generation.

The work of Professor Karl Pearson on heredity was then considered, and found to point to a similar conclusion.

All this, it was urged, tended to show that the marriages of to-day were not only relatively infertile, but either:

1. That the children born of such marriages were weak, neurotic, specially liable to alcoholism, criminality, and insanity, and so far unfit for the battle of life; or
2. That marriages of the middle and better classes were now so sterile that quite an undue and dangerous proportion of the rising generation was recruited from the lower, the more ignorant, the more vicious, and semi-criminal population.

II.

The second part dealt with the cause and life-history of these relatively sterile marriages. The cause was traced to the deliberate prevention of conception, and reasons were brought forward supporting the view that this was not only destructive of the future life of the nation, but seriously injurious to those who practised it, and to such children as were "arranged" for and produced. The prevalence of neurasthenia was increased by the practice, and the absence of sexual restraint encouraged by it had an evil effect on the moral character which was liable to be transmitted to the children. In addition to this, the child of the relatively sterile marriage suffered from a debased and stunted educa-