Fig. 18 shows the same incidents as Fig. 15. Mrs. T., November 3rd, 1903.

Fig. 19 shows a perfectly normal pulse, with all the events in the cardiac cycle following one another in a normal manner. Mrs. T., November 4th, 1903.

Fig. 20.—Inferences drawn from the radial pulse would assuredly mislead in recognizing the cause of the irregularity, as here there seem normal pulse beats followed by extra systoles. The jugular tracing demonstrates the real nature of the irregular rhythm, which shows the occurrence of auricular systole (a) and diastole (x) during the ventricular systole (E). Mrs. T., November 14th, 1903.

Fig. 21 shows the frequent occurrence of extra systoles of ventricular origin, and gives the clue to the explanation of the whole series of irregularities from which the patient suffered. The explanation of the extra systoles and regular appearance of the auricular waves is the same as that given from Fig. 2. Mrs. T., November 19th, 1903.

ON SOME HAEMOGLOBIN INVESTIGATIONS IN CASES OF CHRONIC HEART DISEASE.

By PROFESSOR THEODOR SCHOTT, M.D.,
Bad-Nauheim.

THE physical and dietetic treatment of chronic heart complaints in general, as well as the special treatment by baths and gymnastics, are so widespread nowadays that in the most recent literature the question is much less frequently asked whether these methods have a good effect on heart diseases, but rather in what way they act on the heart and body generally. It would lead us too far to repeat here the various opinions of the subject. The readers of the *BRITISH MEDICAL JOURNAL* would gain more information from the works of Bezly Thorne, Grainger Stewart, Lauder Brunton, Bowles, William Broadbent, Douglas Powell, and others.

In this short essay I only desire to set forth the results of a series of investigations which I undertook last summer with my assistant, Dr. Wolfson, for the special purpose of giving some support to my brother's opinion and my own that the combined treatment by baths and gymnastics has a tonic action not only on the heart but on the whole system. If also the patients should show when the cure is finished a more robust appearance, increased strength of muscle, especially displayed in greater ease in walking and climbing, better sleeping and improved appetite, etc., in short, greater bodily strength, yet it is not superfluous, but rather desirable to become acquainted with those clinical methods of investigation which make it possible for medical men living at a distance to get a clear idea whether such treatment is really a tonic. Besides the weighing of the patient during and after the period of treatment, my former researches, which were principally directed to the state of the heart and vessels, had included heart diagrams, sphygmograms, and blood-pressure tracings, while some other investigations, especially those relating to the composition of the blood, had not been carried out. The reason for not having made the last named was that for general purposes former methods of investigation left much to be desired; either too much time was needed for each separate investigation, or the mass of blood to be removed was so large that it must be taken direct from a vessel; in other words, such fuller investigations as would have extended over several weeks were scarcely applicable. Besides this, with most apparatus the blood had to be diluted or modified.

In the autumn of 1902 I obtained Dare's haemoglobinometer. I must not allow myself to go more fully into a description of this instrument here, I only desire to describe its most essential merits; the first is that one drop of undiluted blood is sufficient for the investigation. For this purpose Dare has constructed a pipette so evenly that in an area of $\frac{1}{2}$ square centimetre one drop of blood is sufficient to fill the capillary space completely. In the apparatus there is a standard solution, the red colour of which corresponds to that of a solution of blood containing 100 per cent. haemoglobin. The drop of blood in the apparatus is compared with this standard solution. Candle light is used, so that the investigator is independent of daylight. The scale is on a revolving graduated wheel, and in that way, as the observer looks through a camera on to a white surface, the proportion of leucocytes in the blood need not be considered.

By following these directions every practitioner can in a short time learn how to make similar haemoglobin investigations within two minutes, and further so to diminish the sources of error that the differences between two observations and the readings of two observers amount eventually to only 1 to 2 per cent.

After sufficient practice on healthy people we began our observations on various forms of heart disease. Naturally care was taken that—besides the cautious manner of living which those suffering from heart disease must observe, with respect to the kind of nourishment, especially as to the nourishing power of the food taken—on the whole no essential change was made. In order to obtain correct comparisons the same hour was always chosen for the same patient for observation, but above all things care was taken that an equal time always elapsed after the last meal. Also it was made the rule that no examination of blood should take place during or after menstruation, or if it did take place it was not to be looked on as of any value as a comparison. The usual investigations of the heart with regard to heart diagrams and heart sounds, the sphygmogram, and above all successive observations with Gaertner's tonometer² always went hand in hand. In this way last summer 120 patients were examined under successive observations, and I permit myself to report here some of them very shortly and cursorily, reserving a more detailed description for a later time. In all cases except where otherwise stated the patients were subjected to combined treatment by baths and gymnastics.

CASE I. *Weak Heart, Dilatation of the Left Ventricle; Anaemia*.—Miss M., aged 18.

May 26th. Before the first bath, haemoglobin 69 per cent. (blood pressure 85 mm. Hg.).

June 3rd. After the sixth bath, haemoglobin 78 per cent.

June 23rd. After the eighteenth bath, haemoglobin 79 per cent.

July 9th. After the twenty-seventh bath, haemoglobin 81 per cent. (blood pressure 110 mm. Hg.).

The patient appeared much more robust after the cure, could walk for hours on level ground, and could climb hills without much difficulty. The treatment consisted of baths only.

CASE II. *Weakness of the Cardiac Muscle; Dilatation of both Ventricles; Anaemia*.—Countess St., aged 52 years.

May 25th. After the fourth bath, haemoglobin 60 per cent. (blood pressure 85 mm. Hg.).

July 13th. After the thirty-third bath, haemoglobin 75 per cent. (blood pressure 100 mm. Hg.).

This very feeble patient had taken iron at home for several weeks without result; during the first three weeks of her stay in Nauheim three bottles of haematicum (Glausch) were taken. The face was still pale at the end of the cure, though rather less so than at first, but the patient felt much stronger.

CASE III. *Mitral Insufficiency after Acute Rheumatic Polyarthritis; Anaemia*.—Miss M. D., aged 15.

June 5th. Before the first bath, haemoglobin 53 per cent. (blood pressure 80 mm. Hg.).

June 14th. After the sixth bath, CO₂ (Spring 7.), haemoglobin 65 per cent.

June 23rd. After the twelfth bath, CO₂ (Spring 7.), haemoglobin 74 per cent.

July 3rd. After the nineteenth bath, CO₂ (Spring 7.), haemoglobin 79 per cent.

July 15th. After the twenty-second bath (third effervescent bath), haemoglobin 81 per cent.

July 30th. After the twenty-seventh bath (eighth effervescent bath), haemoglobin 79 per cent. (blood pressure 105 mm. Hg.).

The patient took no kind of medicine, either for the heart or for anaemia. On leaving she felt completely recovered, and only complained of slight palpitation after climbing stairs and hills. Within the first fortnight she gained in weight 3 lb., and at the end of her stay 9 lb.

CASE IV. *Aortic Insufficiency after Acute Rheumatic Polyarthritis*.—Mr. R. F., aged 14.

June 2nd. Before the first bath, haemoglobin 55 per cent. (blood pressure 105 mm. Hg.).

July 7th. After the twenty-fourth bath (of which nine were sprudel baths), haemoglobin 78 per cent. (blood pressure 118 mm. Hg.).

The young man, who looked anaemic before, looked blooming after the treatment.

CASE V. *Mitral Stenosis*.—Miss H., aged 18.

June 9th. Before the first bath, haemoglobin 68 per cent. (blood pressure 100 mm. Hg.).

June 15th. After the sixth bath, haemoglobin 80 per cent.

June 28th. After the fourteenth bath, haemoglobin 80 per cent.

July 4th. After the sixteenth bath (one sprudel bath, Spring 7), haemoglobin 83 to 84 per cent. (blood pressure 120 mm. Hg.).

The remaining observations of haemoglobin and blood pressure all gave the last-mentioned result. Altogether twenty-four baths were taken.

CASE VI. *Mitral Stenosis and Insufficiency*.—Miss A., aged 23. Whether the condition was congenital or acquired in the first year of life could not be determined.

July 2nd. Before the first bath, haemoglobin 64 per cent. (blood pressure 85 mm. Hg.).

August 3rd. Before the first sprudel bath (Spring 7), haemoglobin 76 per cent.

September 15th. After the thirtieth bath (of which sixteen were sprudel baths), haemoglobin 81 per cent. (blood pressure 120 mm. Hg.).

The patient who had suffered much in youth from heart disease, and

in later years could hardly walk at all, when first examined was of a dark-blue colour, both in face and fingers. On departure the dark-blue colour had in a great measure disappeared, and the swelling of the liver had distinctly diminished. At the end of the cure the patient was able to take short walks of fifteen to twenty minutes, and even to make without difficulty the slight ascent of ten minutes from my house to her dwelling, whereas at first she had to be wheeled up in a bath chair.

CASE VII. *Graves's Disease; Myocarditis; Severe Anaemia.*—Mrs. A., aged 45. The patient had taken iron and digitalis before her stay at Nauheim. A short time after her arrival she had right-sided pulmonary infarct (embolism). She continued the digitalis for three days, and later for a short time took haematicum (Glausch) three bottles.

May 26th. After the fifth bath (first measurement), haemoglobin 58 per cent (blood pressure 95 mm. Hg.).

July 8th. After the thirty-second bath (of which ten sprudel baths), haemoglobin 69-70 per cent. (blood pressure 115 mm. Hg.).

CASE VIII. *Myocarditis Due to Abuse of Alcohol and Tobacco; Dilatation of both Ventricles, chiefly of the Left; Paroxysmal Tachycardia.*—Mr. F., aged 54.

July 8th. Before the first bath, haemoglobin 65 per cent. (blood pressure 115 mm. Hg.).

August 28th. After the thirty-second bath, haemoglobin 85 per cent. (blood pressure 135 mm. Hg.).

CASE IX. *Mitral Insufficiency and Stenosis; Chronic Nephritis.*—Mr. O., aged 22.

May 20th. Before the first bath, haemoglobin 68 per cent. (blood pressure 85 mm. Hg.).

June 4th. After the ninth bath, haemoglobin 75 per cent.

June 15th. After the sixteenth bath, haemoglobin 76 per cent.

June 24th. After the twenty-first bath, haemoglobin 81 per cent.

July 4th. After the twenty-seventh bath (ten of which were thermal-sprudel baths), haemoglobin 82 per cent. (blood pressure 120 mm. Hg.).

The patient, who took no medicine, and at first was very pale, looked blooming in the end, and could walk without the slightest difficulty for hours.

These few cases described here show that the treatment by baths and gymnastics is capable of increasing the formerly depreciated haemoglobin in cases of heart affection, be they cases of weakness of the heart muscle or valvular diseases (simple or complicated), Graves's disease, or chronic myocarditis, with or without kidney disease. Naturally there were many cases in which this increase was only very slight. According to my observations it seems far easier to raise the amount of haemoglobin in young and middle-aged persons than in those of riper years. Even if these latter feel much better after following the treatment, there is frequently no change to be seen in the amount of haemoglobin. The following is an example of this:

CASE X. *Chronic Myocarditis; Severe Arteriosclerosis; Angina Pectoris; Bradycardia* (pulse 30-36).—Mr. L., aged 68.

May 20th. Before the first bath, haemoglobin 79-80 per cent. (blood pressure 160 mm. Hg.).

May 29th. After the seventh bath, haemoglobin 80 per cent.

June 17th. After the twenty-first bath, haemoglobin 79 per cent. (blood pressure 150 mm. Hg.).

During the last part of the cure duplicated heart beats were sometimes noticed, but only on auscultation; not in the radial pulse. Very often there were 54 heart beats, with a radial pulse of 30 to 36.

Very instructive for medical men are these cases in which, during the treatment, a diminution of the haemoglobin is observed; and let it here be noted that severe physical over-exertion and violent emotional disturbances, especially in febrile diseases, in a short time rapidly produce a diminution in the haemoglobin. Of this, in conclusion, one example:

CASE XI. *Weak Heart; Dilatation of Both Ventricles.*—Mr. S., aged 44. He had severe anaemia, for which for many years iron had been taken at home.

May 19th. After the first bath, haemoglobin 70 per cent. (blood pressure 130 mm. Hg.).

June 2nd. After the tenth bath, haemoglobin 85 per cent.

June 22nd. After the twenty-fifth bath, haemoglobin 80 per cent. The patient suffered from feverish bronchitis and had undergone severe physical exertion during a several days' journey.

June 23rd. After the twenty-fifth bath, haemoglobin 79 per cent. (blood pressure 110 mm. Hg.).

During the following eight days the haemoglobin rose gradually again to 85 per cent. and the blood pressure to 130 mm. Hg.

I am well convinced that in an investigation of the blood which in some measure lays claim to thoroughness it is necessary to count the blood corpuscles, both red and white, and to fix the amount of ash of the blood. Such experiments, however, take so much time that in practice, at any rate, investigations on a large scale are very difficult to carry out. I propose at some other time to supply what I have here omitted.

NOTES.

¹ For description see *Philadelphia Medical Journal*, September 22nd, 1900. ² Of these tonometer measurements I give only the first and last in parentheses.

IMPROVEMENTS IN DIETETICS OF DIABETICS.

By SIR JAMES SAWYER, M.D.LOND., F.R.C.P.,
Senior Consulting Physician Queen's Hospital, Birmingham.

COMPARATIVELY successful as is the modern medicinal treatment of diabetes mellitus (glycosuria) under medical prescription and guidance, in many cases—perhaps in all—the appropriate regulation of the patient's food is a chief therapeutic indication. In the management of this difficult malady, it has been recognized by the profession for many years that abstinence from sugar and from other articles of food which are convertible after ingestion into glucose causes a great lessening in the quantity of glucose in a diabetic patient's urine, and also a marked diminution towards normal of the heightened density and quantity of the urine, and that these changes for the better are coincident with arrest of bodily wasting, even with some gain of bodily weight, and also with general improvement, in a very large proportion of those persons who are suffering from diabetes.

To meet these rational and practical indications as to treatment much care has been taken by many physicians to devise an extended, varied, and agreeable and practicable dietary. Until the researches of Mossé were published two years ago,¹ it was our therapeutic rule to withhold potatoes in saccharine diabetes. The potato contains from 16 to 24 per cent. of starch, and it used to be classed without qualification amongst food inadmissible in glycosuria in a case treated upon the dietetic principle I have referred to. His researches have led Mossé to conclude that potatoes, far from being harmful, form a useful and beneficial food in glycosuria, and that they are capable of being substituted for ordinary wheaten bread in daily proportions sufficient to maintain the alimentary ratio, that is to say, in the proportion of two and a half to three of potatoes for one of bread. Mossé found that a daily ingestion of potatoes in quantities of from 1,000 to 1,500 grams, roughly of from 2 lb. to 3 lb. avoirdupois, brought about in nineteen out of twenty cases of diabetes speedy diminution of the glycosuria, quick relief of thirst, and general improvement in the patient, and all this in all forms of diabetes. The salts contained in the potato are chiefly those of potash. Potash is contained in potatoes in much larger proportion than in wheaten bread, and Mossé attributes the superiority of potatoes in a diabetic dietary to the increased ingestion of potash.² My own experience in practice during the last two years is confirmatory of Mossé's conclusions.

I think it will be found in practice that the permission of potatoes in the food of diabetics is one of the greatest dietetic advances of our times. Potatoes cooked in any of the ordinary ways are a grateful food. I would point out that for the retention of the salts of the potato the vegetable should be cooked in a peculiar way. It should be cooked by steaming with its "skin" on. If potatoes be peeled and then cooked by steaming, they lose in the cooking large proportions of their potash and of their phosphoric acid, and they lose more of each of these their constituents if they be peeled and then cooked by boiling.

It has occurred to me that the beneficial results of Mossé's discovery and teaching as to the use of potatoes as a food in diabetes might well be carried much further in the dietetics of diabetics than in merely the free allowance of properly-cooked potatoes in a dietary. I propose that the therapeutic difficulty as to the prohibition of ordinary bread for a diabetic may be met advantageously by making bread, cakes, and biscuits for diabetics by using the "flour" of properly-cooked potatoes, instead of the flour of grain. I find that excellent and delicious baked cakes can be made from paste composed of a kind of flour prepared by rubbing down potatoes cooked with their coats on by steaming, blended with cream and butter. With the aid of some good cooks in my kitchen I have made many experiments as to the formation of bread and biscuits from bran and the flour of potatoes. As the best results at present reached from these inquiries, I have decided upon the following details for the making of these new biscuits and bread for diabetics:

Bran and Potato Bread.—Take half a pound of flour of steamed potatoes, quarter of a pound of bran, half an ounce of German yeast, half an ounce of butter, one egg. Twenty-four hours before making the dough cook the potatoes by steaming them in their "jackets," then peel and break up into flour with the fingers. Mix all the ingredients together, and let the paste stand near the fire for an hour to "rise." Bake in greased tins for an hour and a-half.

In common with many other physicians, I have arranged and employed in practice for many years a diabetic dietary which is sufficiently strict for practical purposes, and which