where it had been previously ligatured the tube was completely divided, and the two ends separated by a distance of about $\frac{1}{4}$ in. No trace of a ligature could be seen. The right tube was then drawn into the wound; it appeared in every way natural, there was no solution of continuity and no trace of the ligature. A loop of each tube in turn was drawn up into the wound, a ligature passed through the mesosalpinx between the two sides of the loop, and the whole tied as a pedicle with a Staffordshire knot. The knuckle of tube on the distal side of the ligature was cut away. The abdominal wound was then closed. The recovery was uninterrupted.

Subsequent Pregnancy.

In January, 1901, the patient was again admitted suffering from an incomplete abortion. The cervix was dilated and the uterine cavity explored with the finger. Portions of placenta were removed.

Cases of pregnancy after ligature of the Fallopian tubes have been from time to time reported, and the following instances throw some light upon the method of passage of the ova along tubes whose lumina have been apparently occluded:

A case is reported by Horrocks¹¹ "in which a patient had been sterilized after Caesarean section by ligaturing the Fallopian tubes. In spite of this she became pregnant again, and the uterus ruptured along the line of incision when near full term, the child and a portion of the placenta escaping into the abdomen. Dr. Galabin removed the uterus, and the specimen is now in the Guy's Hospital Museum. The ligature on one side had cut through the tube, and the severed ends lay about half an inch apart. On the other side the tube and ligature looked as if the tying had only just been done, but on experimenting it was impossible to force a coloured liquid through the tube. Still, it was thought that in all probability the ovum, which had become impregnated, had got past the constriction produced by the ligature, and had so entered the uterus."

Galabin¹² reports a case in which, after performing Caesarean section, he tied the tubes with kangaroo tendon, and the patient became pregnant again within a year or so.

Taught by the experience of these cases, he continues, I have since then adopted the plan of cutting a piece out of the tube between the two ligatures, then pulling out the stump on each side as far as possible and cutting it off, so that the open lumen was left at the bottom of an inverted case of cellular tissue. None of the patients so treated have become pregnant again, and I am inclined to regard this as a reliable method.

That removal of portions of the tube wall alone, however, is not a reliable method is shown by the case we have just recorded.

Fig. 1.

Doran has suggested that the ligature may perhaps ulcerate through the tube, which then heals behind it without causing stricture of its canal. It is well known that sometimes a ligature which has been applied to the cut end of the ureter after nephrectomy, may subsequently be passed from the bladder, and the following case shows that the same thing is possible in a Fallopian tube.

Fig. 2.

In 1898 Blaud-Sutton removed an ovarian cyst, its slender pedicle was tied with thin silk; the recovery was uneventful, but for many weeks afterwards the patient complained of cramp-like pains on the side from which the cyst had been removed. The pains gradually

passed away and ten months later, during menstruation, the patient accidentally noticed on her napkin, a tiny loop of silk. This proved to be the loop of silk used to secure the Fallopian tube; it had ulcerated into the tube, and had been very slowly conducted into the uterus and so escaped.

The occurrence of pregnancy after removal of portions of the tubes is more difficult to understand. It can hardly be that the severed ends of the tube have become reunited, and the canal again patent; but it is possible that a process of absorption may have occurred in the walls of those parts of the tube which were in contact with one another. The process ultimately leading to the formation of a fistula between the two portions, so that the continuity of the canal once again becomes established.

There is another point of both clinical and pathological importance to be noticed; in all the cases quoted the lumen was either completely obstructed or narrowed—in Horrocks's case coloured fluid could not be made to pass the obstruction—yet in none of them was the subsequent pregnancy a tubal one.

These two cases taken together may perhaps add something to our knowledge of the causation of tubal gestation. In Case I we have an instance of tubal gestation with external migration of the ovum, but with no occlusion of the lumen of the tube; in Case II the lumen of the tube on one side was completely obliterated, and on the other must have been considerably narrowed, but the pregnancy was uterine.

We wish to tender our thanks to Dr. Champneys for permission to mention these cases.

REFERENCES.

- ¹ Baldwin, *Amer. Jour. Obstet.*, December, 1902. ² Ueber zählige Eierstöcke *Arch. f. Gyn.*, 1898. ³ *Obs. Soc. Trans.*, 1902, page 231 et. seq. ⁴ *Wien. klin. Woch.*, 1891. ⁵ *Obstetrics*, page 79. ⁶ "Ueber Extrauterinschwangerschaft" *Volkman's Sammlung klin. Vorträge*, N. F., 1899, Nos. 244, 245. ⁷ *Operative Gynaecology*, vol. ii, 189. ⁸ *Obstetrics*, page 80. ⁹ "Ueber die Ursachen der Ansidlung des Eies im Eileiter," *Zeit. f. Geb. und Gyn.*, 1890. ¹⁰ *Zur Kenntnis und Behandlung der Jubaasschwangerschaften*, *Monats. f. Geb. und Gyn.*, 1897. ¹¹ *Obs. Soc. Trans.*, 1900, page 243. ¹² *Ibid*, 1902, page 246.

ON VENTRIFIXATION OF THE UTERUS.

By W. J. SINCLAIR, M.D. ABERD.,

Professor of Obstetrics and Gynaecology, Victoria University of Manchester.

In a large proportion of the cases of old standing chronic backward displacement of the uterus our ordinary methods of treatment do not give satisfactory results. When the body of the uterus has become considerably enlarged after backward displacement, or when even the slightest tubal and ovarian complications exist, what could be expected from more or less forcible turning forward of the body, and the insertion of a pessary, in the consulting room? There is an element of levity in this method of dealing with a flexion producing symptoms.

When the patient is persuaded to take to bed, and remain as long as if she had to undergo an important operation, some good can usually be done. The congestion of the body of the uterus can be relieved by tampons and douches, the amount of relief being measured by the diminution of the tenderness to touch and certain changes in the visible portion of the uterus. When the uterus has been restored by tampons almost or altogether to its normal position, then is the time for the introduction of a pessary. In some uncertain proportion of even old chronic cases this treatment gives sufficient relief to satisfy the patient, and a small number of cases are actually cured in time. As a general rule it may be stated that the older the patient the more suitable the palliative treatment.

But in married women at the child-bearing time of life, and in many single women who cannot, or will not, lead restful lives, the treatment by mechanical support is unsatisfactory in many ways; and something more must be done.

Schultze's process may be dismissed with a word or two. It has never done positive harm to any patient of mine as far as I know, but upon the whole it has given no satisfaction.

Schücking's operation and similar timid and dangerous proceedings I have never tried, and never will.

Alexander's operation of shortening the round ligaments obtained great favour for many years, and still holds its place as a valuable method of treatment in a certain limited class of cases. The eagerness with which it was adopted by gynaecologists throughout the world was a fair measure of the universal dissatisfaction with the mechanical treatment of displacements of the uterus. We may also now consider

the number and variety of the recent modifications of Alexander's operation as a fair indication of the disappointment of operators who have applied it without sufficient discrimination as to its limitations. For my own part, after operating on 45 cases I abandoned the method of treatment altogether, as the result of certain experiences, to detail which would be too much of digression considering my present object. In one case I could not find the round ligament at all on one side; that of course would be explained by my ignorance of anatomy. In another case the ligaments were pulled out of their sockets, as it were, at six months' pregnancy, and the patient had complete prolapse after her confinement. My troubles suggested to my mind serious objections to the operation in private practice, owing to an element of uncertainty as to the possibility of its satisfactory completion.

The modifications of Alexander's operation which have been most pressed upon the attention of gynaecologists appear to have all the weaknesses of the original operation, or to be more extensive and presumably more dangerous than ventrifixation. This objection attaches, for example, to the proceeding of opening the vagina behind the uterus in order to break down adhesions before dealing with the round ligaments, themselves also probably involved in the inflammation which produced the adhesions. And the objection on account of its formidable character applies to Goldspohn's modification which involves all the details of abdominal section with two wounds instead of one as in ventrifixation. The formidable array of modifications involving hooking and turning and twisting of the ligaments from a wound in the vagina either in front of or behind the uterus, or by means of more or less formidable wounds in the inguinal regions, seem hardly worthy of serious attention, at least from the practical therapeutic point of view.

The operation of ventrifixation appears to me to be the best for the treatment of chronic retroflexion of the uterus in women at the child-bearing time of life. It was suggested, I believe, by Olshausen in 1886, when he fixed the uterus very slightly as an incident in an operation undertaken owing to other indications than displacement. The operation has been much practised with a defective technique in Germany, and what implies much in its favour, it has been taken up in France, on its merits, although originating in Germany.

My first operation was done in September, 1890, and I have gradually evolved a method of operating which has fulfilled all my hopes and expectations. As an operation of choice it had to comply with certain requirements or to stand condemned on its demerits. It does not seem to me to deserve condemnation. The tests which such an operation must stand appeared to me to be (1) it must be devoid of danger when the ordinary care is exercised; (2) it must be easy to perform and the operator must know in recommending it to a patient that he can undertake it with absolute certainty of being able to finish it as planned; (3) it must relieve the symptoms which it was undertaken in order to relieve; (4) it must not produce any new symptoms that will even remind the patient for any considerable time that she has undergone an operation; (5) it must leave the patient free from any danger and permanently in a condition for active exertion and for all the duties of life indistinguishable from her original normal state of health; (6) it must stand the strain of pregnancy and parturition and the puerperium, as if the operation had never been done; and (7) the uterus must afterwards continue to maintain the position it was placed in by the operation. An operation which complies with all these conditions has many merits, whatever *a priori* objections may be raised by armchair gynaecologists, or whatever may be the terms employed to justly condemn an operation labelled "ventrifixation" by those who have met with the unfortunate experiences which were only to be expected from a defective method of operating.

In order to obtain the desired results certain principles must be kept in mind and complied with in every case. In the first place, *the fundus uteri and the round ligaments must be strictly safeguarded*, so that if pregnancy occurs it will run a perfectly normal course. Some time ago, in a discussion on Alexander's operation at a meeting of a gynaecological society, I expressed this opinion, and was hardly surprised at the shout of derisive laughter with which the statement was received. However, the confidence derived from long experience of the operation was not likely to be shaken by the adverse opinion of men who obviously did not know the essentials of an efficient operation, or who accepted and applied de-

fective surgical methods without criticism and without endeavouring to account to themselves for their failures.

Another important point is to effect complete and permanent adhesion of a sufficient area of the anterior surface of the uterus to the abdominal wall. The ideal ventrifixation is that which results from Caesarean section, when the wound is longitudinal, involving about the middle third of the uterus, and complete adhesion takes place between the uterine and abdominal wounds. In these cases when pregnancy occurs again—as I have known it to do repeatedly—there are absolutely no abnormal symptoms, either functional or mechanical, and the operation may be repeated almost or altogether without opening the peritoneum. I have only once had occasion, apart from Caesarean section, to open the abdomen in a patient who had undergone ventrifixation in the manner which I am advocating, and I took the opportunity of examining the relation of the uterus to the abdominal wall. The uterus appeared to be completely incorporated with the anterior parietes in its middle third, and yet the patient had never experienced any unpleasant symptom referable to the uterus or bladder during the years that had elapsed since the operation. If this can be true in one case, why not in all? Firm attachment is not in itself troublesome or dangerous.

Before going into the details of the operation, let us consider what we should avoid. We can, perhaps, best do this by looking at the descriptions of the operation in some of the manuals and textbooks to which beginners in abdominal surgery naturally look for guidance.

Dr. Howard A. Kelly's method of operating has been much followed by gynaecologists abroad, and to some extent, I believe, in this country; but it is extremely faulty in several respects. The introduction of sutures "through the posterior surface of the uterus" should be in itself sufficient to condemn it. In what respect is it better than that worst of all important operations practised on women at the child-bearing time of life, vagino-fixation, to wit? The operation is suspension of the uterus, not fixation; and to the extent to which it fails, owing to stretching of the adhesions, it benefits the patient only by relieving the fundus from its misplaced attachment, while perhaps it adds to her danger from ileus owing to the presence of adhesion strings or bands between the fundus and the abdominal wound. No wonder that a large proportion of Dr. Kelly's patients who became pregnant experienced suffering from abnormalities. But suspension injuriously affects all cases, whether pregnancy occurs or not. It causes dragging upwards of the uterus instead of supporting it, prevents it from firmly contracting, and makes it into an atonic bag, with resulting menstrual anomalies and troubles.

This is the operation of which Dr. Clarence Webster of Chicago says, in his textbook, it "may be recommended." I have looked through most of the American manuals of gynaecology, and it may be stated generally that the authors follow Kelly, or do worse. Dr. E. C. Dudley of Chicago, whose work is deservedly held in high esteem in the United States, "uses two formaldehyde catgut sutures, one on each side of the posterior wall of the corpus uteri."

Among the books which are read for guidance in England is *Diseases of Women*, by Dr. G. E. Herman. If I understand the author's meaning in the paragraph headed "How to Perform Ventral Fixation," I must say, with deep regret, that I regard his method as defective and faulty in almost every detail, even down to the "blunt" volsella. The first suture is made to enter close to the Fallopian tube; the muscle, not the peritoneum, is depended upon for adhesion and suspension of the uterus; catgut is the material employed to fix the fundus in the wound, and a silk thread is inserted still further back. Silk sutures passing through the skin and uterus are allowed to remain *in situ* for eight days. The patient is kept in bed only a fortnight.

The defects of the operation, described by Sutton and Giles, are in the material "fishing gut," which is referred to as if it stood on the same level of suitability with silk, and in the fewness of sutures. The operation may be completed with one suture through the uterus and peritoneum only. "If desirable two or more sutures may be introduced."

Exercising such judgement as I am capable of, I cannot see why such operations should be designated ventral fixation, or why they should ever be expected to give more than temporary relief from the symptoms due to retroflexion.

The operation, which was performed by Laroyenne, of Lyons, complied with all the requirements necessary to

confer permanent benefit. Instead of using the volsella in the fundus to steady and manipulate the uterus, Laroyenne employed a stem in the uterine canal, which was manipulated by an assistant, a detail of the technique, with obvious disadvantages. But the operation produced permanent adhesion of the uterus to the parietes above the bladder.

Some of the German operations appear to be worthy of imitation with modifications. Olshausen's method is defective in as far as it fixes in the wound the cornua of the uterus. The method associated with the names of Leopold and Sippel appears to be a good operation, but it is not even mentioned by Küstner, who writes the chapters on malpositions of the uterus in Veit's *Handbuch der Gynäkologie*. Küstner makes much of the "*Beweglichkeit*" of the uterus as a condition necessary for success, but the need for mobility is a pure *a priori* assumption. He cannot find a shred of support for his objections in the practice of any one who does an efficient operation, producing firm "fixation."

In performing the operation of ventrifixation I have discarded all suture material except catgut and the finest silk thread which can be conveniently manipulated. At first I used to try to close completely the vesico-uterine fold, but now for years I have introduced for this purpose only two catgut sutures, one at each side, well below the lowest point of the abdominal wound, while the uterus is firmly pulled upwards by means of an ordinary volsella inserted in the centre of the fundus and manipulated by an assistant. This is quite sufficient practically to close the fold without interfering with the distension of the bladder, and it is more of a preventive to any possible insinuation of intestine than shortening of the round ligaments, to the extent practised by most operators, can be. The next step is to suture the peritoneum at the edge of the wound on each side. The fine silk sutures are introduced from the outside surface of the parietal peritoneum, passed through the peritoneum, and after taking a good hold of the uterus low down and near its margin, they are brought back through the peritoneum and firmly tied. The knot is thus always extraperitoneal. Two of such sutures are, as a rule, introduced on each side. The same process is now continued, with the addition that the sutures include the fascia as well as the peritoneum on each side. The transverse sutures are next introduced. These are two or three in number, consist of fine silk, and include fascia and peritoneum at the sides, and each takes a fairly extensive hold of the anterior surface of the uterus. The highest of the sutures is placed about half-way up from the isthmus to the fundus. In complicated cases, and when the prospect of pregnancy is very remote, the sutures have been placed occasionally a little higher, though not often. All the sutures that penetrate to any part of the uterine tissue are buried sutures at the finish; or if any suture includes skin as well, it is never left for more than three days, usually only forty-eight hours. There must never be suppuration in the track of this suture. The highest transverse suture should always be buried. The abdominal wound is closed in the usual way, and it is an advantage to suture separately the edges of the fascia with catgut. The last step in every case is the introduction of a glycerine-pad pessary into the vagina to press forward and support the uterus during the formation of adhesions. The patients are kept in bed for a month or six weeks, and they are advised to lead invalid lives for several months.

There is hardly ever the slightest reaction after operation in simple cases. The pulse and temperature remain normal. No case has given rise to even momentary anxiety.

This operation has stood the tests of efficiency already mentioned in nearly one hundred cases. The details of these cases I hope to find an opportunity of publishing before long. Many of the operations have been done on account of prolapse of the uterus in comparatively young women, but I do not propose to complicate matters by considering the treatment of prolapse on the present occasion.

In only one case has the treatment of ventrifixation been alleged to have failed. The medical practitioner, who sent the case to the Southern Hospital, when applied to a year or so after for information returned a discourteous reply, alleging that the operation was a failure; but he has given no information, and he has never complied with the request for the address of the patient, so that the details of the after-history could be obtained. The patient was perfectly well when she left the hospital.

A considerable proportion of the patients have borne one or more children since the operation. There have been some abortions and one or two cases of malpresentation, but not, I

believe, to an extent greater than might be expected among the same class of women, apart altogether from operations. The after-history of every private patient has left nothing to be desired.

Sometimes there has been some bladder discomfort for a time, but it is remarkable how seldom any complaint is made even temporarily.

Efforts have been made to get at the present state of health of the patients, but it is often difficult or impossible to trace them and obtain information. As far as I can learn, there has been no case of relapse, even after labour. In one woman of the poorest apathetic class, ventral hernia has occurred owing to premature hard work and leaving off her abdominal belt.

I have entire confidence in the future of ventrifixation if an efficient method of operating is generally adopted, while joining in the condemnation of ventral suspension of the uterus, designated improperly by so many operators as ventral fixation.

A WARNING SIGN OF DANGER TO THE CHILD DURING LABOUR.*

By FREDERIC E. DANIEL, M.R.C.S., L.R.C.P.,

Honorary Secretary, Furness Division of the North Lancashire and South Westmorland Branch of the British Medical Association.

In an ordinary breech presentation, after the birth of the body, there is generally some delay, especially in a primipara, before the head follows. During this time a certain amount of compression of the cord is inevitable, and, if this is prolonged too much, the child's life is threatened from deficient aeration of the blood. While the child is in this position the practitioner is generally cognizant of any attempts it makes at respiration. In fact, the movements of the child are the only indications that we have that the cord is being compressed.

Now, I believe it is not very generally known that during the second stage of labour in an instrumental case with the head presenting, when from any cause the placental circulation is interrupted, the same movement can be felt by the surgeon, communicated through the handles of the forceps. This movement is an indication of a pressing desire on the part of the child to breathe, and can only be brought about by interruption to the free circulation through the umbilical cord. It is quite different to an ordinary kick, and is distinctly characteristic of an inspiratory effort.

CASES.

Before describing it further I will briefly narrate four illustrative cases that have occurred in my practice since February, 1902. I had during that year a run of bad luck, and was unfortunate enough to lose four firstborn children, all of fairly well-to-do people, who had looked forward with lively satisfaction to becoming parents. The mothers were all primiparae, and the four children were fine, healthy, well-developed babies, and at the commencement of labour were undoubtedly alive and, literally, kicking, yet were all from an unavoidable cause stillborn.

CASE I.

The patient was about 34 years of age. The presentation was normal, and everything promised well, and the mother was told that everything was straightforward. All went well until the full dilatation of the os, when progress stopped, though pains continued regularly. After waiting some time, I administered chloroform and applied the forceps. Dilatation was slow, and a good deal of traction during the pains was necessary. Before the head reached the perineum, during the intervals between several pains, I noticed that I could feel a fetal movement through the handles of the forceps. It was quite different to the ordinary kick, and I am sure it was not the first time I had felt it, but it occurred to me for the first time that it might be caused by the child attempting to take an inspiration. This was repeated several times, and eventually ceased.

It was an hour or more before I was able to deliver the child, as the perineum was very rigid, and required a great amount of patience; but delivery was effected without any rupture. Unfortunately, however, the child was dead. The umbilical cord was round the neck, and in a state of great tension, and I was unable to get it over the head, and was obliged to ligature it where it was, and divide, before extracting the body. I, of course, adopted the usual resuscitating methods, but they were of no avail. I had not, however, much expectation of being able to resuscitate, as the pulsation of the heart had clearly stopped. Now the mother had felt the child strongly up to the time of administering chloroform, so that there was no doubt that it had died during the second stage of labour. On thinking the matter over, I came to the con-

* Read at a meeting of the Division.