looked upon as a mutilation from tuberculosis, the true magnitude of the evil would, I think, strike home more forcibly. We should hear far more of efforts at prevention than of academic discussions upon cardiac *bruits*, and of hopeless attempts at treating scar tissue. It is with this aim in view that I have again brought forward a study of the infective nature of the disease, so clearly foreshadowed in this country by Dr. A. Mantle nearly twenty years ago, yet then so little regarded. The logical step from a practical interpretation of these facts is the prevention of rheumatic fever. It matters not whether, as some hold, there is a specific micro-organism, or as others, that there are several different infective agents, the essential fact remains unaltered. Rheumatic fever is infective—much heart disease results from infection—the first step in treatment must be preventive.

The time, I believe, has come to study once again by the light of this view the predisposing causes of rheumatic fever. To trace once more the influence of climate, of cold and damp, and of sudden changes in temperature. To inquire into the incidence of the disease on those who live in new and jerry-built houses, into the effects of overcrowding and malsanitation. To study epidemic outbreaks and cases of apparent contagion. To investigate the part taken by school life among the poor, with its outbreaks of sore throats and its strain upon the system of the ill-fed and ill-clothed. Surely there is hope from such investigations that the prevention of some of the rheumatic fever which causes such pitiable suffering among the children of the poor will be thus obtained.

ABDOMINAL PAIN IN ACUTE RHEUMATISM.

By S. VERE PEARSON, M.B.CANTAB., M.R.C.P.LOND.,

Medical Registrar, St. George's Hospital; Assistant-Physician, East London Hospital for Children, Shadwell.

AMONGST the minor symptoms of acute rheumatism of frequent occurrence in childhood still pointed out some years ago¹ that pains in the epigastrium and about the bases of the axillae were of the first importance. Poynton,² too, refers to the frequency of "obscure abdominal pains" in the rheumatism of childhood, and gives it as his impression that they "are generally paroxysmal, and usually referred above the umbilicus to the epigastric region." Donkin, in his *Diseases of Childhood* (1893), p. 212, states that "the painful onset of rheumatic attacks is sometimes accompanied by rigors, occasionally by vomiting, and not very rarely, especially when there is cardiac involvement, by marked precordial or epigastric pain." Much might be said about each of these features, but it is to the pain in the upper abdomen that I wish to draw attention now; for having seen a good deal of acute rheumatism recently, especially in children, I have been much struck not only by the frequency of this symptom, but also by some characteristic features not previously noted, so far as I know, which are present in a certain number of cases.

The manifestations of acute rheumatism in childhood are so manifold, and occasionally their advent is so insidious and ambiguous, that it is rather surprising how little some of the "minor symptoms of frequent occurrence" have been studied and emphasized. The epigastric pain is not peculiar amongst these minor symptoms in repaying a more thorough investigation than has hitherto been granted them, but in this paper I shall confine my attention to this symptom alone. Perhaps the frequency of aches and pains about the abdomen in children has interfered with the detailed examination of the "epigastric pain" of rheumatism, and the common occurrence of "pains all over" in many a febrile infection accompanied by pains in the limbs has led to a callousness on the part of clinical observers. In my experience few pay attention to the abdominal symptoms occurring in the acute rheumatism of either children or adults, yet often it is these symptoms that are the important ones in diagnosis and treatment. In children pain is the most frequent of the abdominal symptoms, and it is often of help diagnostically. In its most typical form it conforms to the following description.

SITE AND CHARACTER OF THE PAIN.

The abdominal pain is nearly always situated about the upper half of the abdomen. The frequency with which children are liable to refer all intra-abdominal pain to the umbilicus makes it striking that this pain is seldom referred solely

to that spot, for though occasionally it is said to run across the middle line a little above this landmark it is rare for the child to point there only. It usually seems to start from a point a little below the costal margin, close to the mid-clavicular line on either side, and to run from there either towards the umbilicus or straight across the epigastrium. At first I thought it was rather commoner on the left side than on the right, but recently I have seen a number of cases where it has been confined to the right side, and until I have collected a rather larger number of cases I am not prepared to say upon which side it is commoner for it to begin. It often occurs on both sides, and then may be either equally common on both sides, or more frequent on one than the other. Occasionally it is more purely epigastric, and about the middle line.

It is not a superficial pain, but is invariably said to be "inside," yet it does not go through to the back nor travel round the trunk. It is always described as being of a sharp nature, and in many cases that I could relate has been a persistently recurring symptom giving rise to considerable suffering. It is more acute than most of the ordinary stomachaches, but by those children who have experienced both this pain and a stitch in the side it is described as being less severe as a rule than the stitch. It is "quite different" from the pain that may be produced as the result of the administration of aperient drugs, but most children are unable to put into words the features upon which this difference depends. It is of short duration but recurs. Each paroxysm lasts from a minute to four or five minutes, though occasionally its duration is somewhat more prolonged than this. "Running," "crossing the street," and "going to bed," are the three commonest exciting causes of paroxysms. It appears to be brought on, in other words, most frequently by muscular exertion, by excitement, and by other factors the exact nature of which are somewhat obscure.

The pain is not associated with nausea or vomiting, nor with any of the ordinary symptoms of gastric derangement, except accidentally. It is not the result of any irregularity or indiscretion in the diet, nor does it appear to be in any way causally connected with either constipation or diarrhoea. There is no abdominal distension accompanying it, and as far as I can make out it is not associated with any gurglings or ordinary colics of the gut. Loss of appetite is not necessarily, nor usually, an accompaniment of it. The pain comes on independently of food, and the nature of the diet the child is taking does not generally influence its occurrence. It does not as a rule double the patient up, but should it come on during exercise it will most likely cause the patient to keep quite still, and often to sit down also. It does not appear to be associated with soreness of the abdominal walls nor with much tenderness, either superficial or deep. The attacks vary very much as to the frequency of their occurrence. I am inclined to think that a typical case would be one in which six or seven paroxysms occur daily for a few days, then three or four days come with fewer attacks, or perhaps none, followed by another period of increasing frequency culminating at about the end of a total of ten days in an illness that manifests itself as acute rheumatism by the occurrence of some of the well-recognized phenomena of that malady. But it is perhaps futile to talk of a typical case when the variability of the frequency of occurrence of these attacks is so marked and when the number of cases illustrating the feature is not yet very large.

This symptom is, I think, commoner in girls than in boys. The cases conforming most accurately to the above description have been in fairly well-nourished girls of 9 to 16. When much younger than this the patients are generally incapable of describing their symptoms with precision, and amongst patients older than this the symptom is not so common. It does occur more or less typically, however, in older children, and probably occasionally in young adults. The phenomena of rheumatism that are likely to be associated with this symptom follow no rule. As far as I can make out, any of the better recognized manifestations of acute rheumatism may occur in association with these paroxysms of abdominal pain, and the latter are not more likely to be associated with cardiac manifestations than with some of the other phenomena, except, of course, in so far as is explained by the great frequency of cardiac affection at this age. The pain does not appear to have any direct connexion with the severity of the rheumatism, either as it affects the heart or as it affects other structures. The peculiar paroxysms moreover appear to be rare in relapses of rheumatic arthritis.

ETIOLOGY.

The question as to the cause of this pain about the upper abdomen is of great interest. Still³ says, "the 'pains in the stomach' may sometimes be in the abdominal muscles, but I suspect that it is usually gastric in origin, and is due to a catarrhal condition in some way dependent on the rheumatism." Any explanation of its causation must at present rest upon mere speculation, and I do not propose to enter now into the discussion of the various theories that could be propounded. But from what I have already said it is clear that I do not agree with the suggestion that these peculiar paroxysms of pain are dependent upon gastric catarrh.

TREATMENT.

The treatment of this symptom is the treatment of acute rheumatism, often in children, especially when symptoms of but slight severity are present, a matter of considerable difficulty. Ordinary remedies directed to cure other kinds of recurring abdominal pains and to alleviate stomach-aches of more usual occurrence appear to have no influence upon the symptom, whereas if other indications of rheumatism arise and are such as to necessitate thorough treatment of the malady in bed, the pains will almost immediately disappear. This therapeutic test may be of diagnostic value. Like so many of the symptoms of acute rheumatism in childhood sodium salicylate, when given in moderate-sized or small doses to a patient who is allowed to go about and live on a more or less ordinary diet, will not infrequently alleviate this symptom and help to ward off its recurrence. But as a rule when the symptom is at all prominent, or when the urgency of other rheumatic manifestations demand, its cure is only affected by putting the patient to bed and insisting on full antirheumatic treatment. I may add that I have found aspirin very useful in one or two cases for which I have tried it.

DIAGNOSIS.

This is not the only kind of abdominal pain that may be associated with the outburst of acute rheumatism. And even should pain having these peculiarities be present, it is hardly to be expected, considering the age of the patients and the variety and variability of the circumstances under which they are likely to be situated, that it is always possible to get a full and exact account of them. I have already indicated that other pains about the belly arise not infrequently in connexion with this malady. Sometimes they are the result of treatment, for large doses of alkalies and sodium salicylates are administered as well as active aperient drugs. But I am quite sure that the symptom I have described is not the result of treatment. Moreover, stomach-aches of an ordinary nature, usually dependent upon some gastric derangement arising from one cause or another, are occasionally prominent symptoms of the onset, or more rarely during the course, of an attack of rheumatism. Then there is rheumatic appendicitis, and a rare condition thought to be due to rheumatic affection of the fibro-muscular abdominal parietes which arises suddenly and is associated with "acute abdominal pain and tenderness so severe as to simulate peritonitis" (Dawson Williams).⁴ But the point I wish to emphasize is that these pains that I have described are of fairly frequent occurrence and are more or less characteristic of acute rheumatism, for they have peculiarities that distinguish them from many of the other pains occurring about children's bellies. They are not absolutely pathognomonic of acute rheumatism, but when a definite account of them can be obtained and most of their typical features are present, they should arouse strong suspicions of the disease, and often their presence leads to a more careful search for indications of rheumatism than might otherwise be undertaken, and is a helpful feature in clearing up the nature of an otherwise doubtful case.

Much patience is required in investigating such symptoms in a child; one has to make friends with the patient first of all. Leading questions cannot be avoided altogether, but by careful and judicious questioning details as to special symptoms can be obtained. It is all too easy to get an account of a symptom out of children that conforms to the preconceived ideas concerning that symptom, the patient acquiescing too readily in any suggestions offered to him. I have tried to avoid such a source of error, and one reason why I have not delayed publishing these remarks for a longer time, during which I might have gathered more material, is that I might incite others to investigate this symptom and thereby remove the personal element from the investigation. If the peculiar abdominal pains I have described are not met with by other observers, I shall at all events have drawn further

attention to the frequency of pain in the upper abdomen as a symptom of acute rheumatism in children, a frequency not yet, I think, sufficiently well recognized. But that there is some foundation for my description and that I have not drawn an altogether fanciful account of this feature can be seen by referring to the few cases I detail below. They are the most typical that I have seen during the past two months, and I have chosen them out of many that I have collected because they are fresh in my memory, and well illustrate many of the points emphasized above. For permission to make use of four of the following cases I am much indebted to the physicians of St. George's Hospital.

CASE I.

G. A. was a well-nourished girl of 6, who was seen by Mr. Fairley Clarke on January 6th on account of stomach-ache. He administered a purgative and gave her an ordinary mixture and instructions as to diet, but did not relieve the symptoms. A week later on examining her very thoroughly he discovered two small subcutaneous nodules close to the spinous processes of the upper lumbar vertebrae on the left side of the back. Not being sure as to whether these were rheumatic nodules or not, and being somewhat puzzled about the nature and treatment of the abdominal pain he referred the case to me. I saw the case on January 13th. I made the following notes at the time of my examination:

Family History.—Mother had acute rheumatism six years ago; in bed three weeks. Father well. One sister, aged 5, well. No other children.

Personal History.—Enjoyed good health up to about a fortnight ago, except for measles and whooping-cough in the past. About a fortnight ago started wasting, and having pains in her inside coming on suddenly when she walked quickly or when she stooped. Usually on starting to walk. They last a few minutes at a time, but were frequently repeated. No particular time of day. Not associated with any indiscretion in diet, and bowels always regular, no nausea or vomiting. Appetite poor for a fortnight. Three days ago had transient pain and slight swelling in the knees and lower parts of thighs.

Present State.—Looks rather pale. Tongue: Moist; yellowish fur, not thick. Rheumatic nodules on the back, none elsewhere. Joints: Nothing abnormal. Heart: No dilatation, no murmurs, first sound rather soft.

Course.—The child was put to bed and treated in the ordinary way for acute rheumatism. Well-marked crops of nodules made their appearance, some arising about the spines of the scapulae and about the posterior parietal regions of the scalp, as well as in the commoner situations for them. There was no fever. The abdominal pains did not recur. Recovery was eventually satisfactory.

CASE II.

C. A. was a girl, aged 10, who was admitted into St. George's Hospital on February 23rd under Dr. Ewart complaining of pains in the stomach, joints, throat, and head. Her temperature was 101°, pulse 112; slight dilatation of the heart; coated tongue. Abdomen: Slight tenderness and resistance to deep pressure over the upper part of the abdomen, especially about the middle of both hypogastric regions. No superficial tenderness, no distension, and movement good. Joints: Slight pain on movement of both elbows and knees. About the middle of both antecubital fossae there is a little fullness, pinkness, and tenderness. Tonsils appeared fairly normal, perhaps a little hyperaemic. Her history was as follows:

Family History.—Negative.

Past Personal History.—No previous rheumatism. Except for measles when 3 years old quite healthy.

History of Present Illness.—On February 19th she went to school complaining of pains about the abdomen; at midday was sent home on account of these becoming worse, and because slight headache and pain in the elbows were also complained of. Headache, pain on movement of the elbows, and "pains in the stomach" continued up to the time of admission into the hospital. On February 22nd pain in both knees and some soreness of the throat was added to the other symptoms. February 21st was the only day she was kept in bed.

She was a particularly bright and intelligent child, and though only 10, could define all her symptoms very clearly. I made the following notes about her abdominal pain on February 24th. "They are sharp pains of a severe nature situated about the upper abdomen on each side, lasting only a few minutes, but recurring several times during the day. These pains are brought out especially on walking fast, or when she goes to bed at night. They were the first symptoms present; they have been the worst symptom, the most distressing trouble throughout. They are not made worse on coughing or breathing deeply. They are situated in the right upper and left upper abdominal segments, more frequently on the right side than on the left. They have no association with the taking of food; they are not accompanied by any nausea. They 'seem to be inside,' and not superficial in the least. They have not been produced by the taking of any purgative, nor do they accord well with the pains due to intestinal colic. They run from the middle of the hypochondriac region from a point just below the costal margin to a point just below and outside the umbilicus on each side. The pain is often present on both sides at the same time. There appears to have been been some soreness to the touch, but this was not a prominent symptom; rubbing the stomach was not thought of." Then follows an exact account of the action of the bowels and the administration of

some liquorice powder eliminating these as causes of pain, and a note, which I need not add, setting forth the mother's account of the symptoms corroborating the above details.

The patient was treated in the ordinary way. All her pains left her soon, and her recovery was satisfactory.

CASE III.

N. H. was a girl of 12, whom I saw first of all on February 5th as an out-patient at St. George's Hospital. The symptoms complained of were recurring "fainting fits," and recurring paroxysms of abdominal pain. There was no family history of rheumatism, but a fortnight before the patient had had sore throat and pain all over. She was a well-nourished, well-developed, intelligent child. The abdomen was normal. Temperature 100.2°. Heart's impulse diffuse, action rapid, and distinct left-side dilatation. There were no vacant beds in the hospital that day, so she was given some sodium salicylate and sent home; but her symptoms were not relieved, and on February 19th a vacancy was found for her and she was admitted into the hospital under the care of Dr. Rolleston. Her condition then was much the same as upon February 5th, the temperature being 100°, and the pulse 120, low tension, rather irritable and jerky. The left heart was dilated, the area of cardiac dullness extending to just outside the left nipple. At the apex, the sounds were embryocardial and rather abrupt in character, and the rhythm tended to alter on movement. There was a haemic *bruit*, best heard in the third left space, and the pulmonary second sound was a little accentuated.

A few days after her admission I wrote the following note about her abdominal pain: "Nearly two years ago she saw a doctor in the country on account of recurrent sharp pains about the upper abdomen. In other respects her health was good. She was told that it was colic, and prescribing was probably in accordance with this diagnosis. But she was not relieved, for from time to time since then attacks of this pain have recurred. They have sometimes been associated with growing pains in the legs, but these have not been sufficient to make her limp. The pain in the upper abdomen has occurred in short paroxysms lasting but a few minutes, but recurring frequently throughout the day. They come on especially when she runs or walks fast, and when they come on she stays still, and doubles herself up somewhat; she does not rub her stomach, nor sit down. The pain comes in the right hypochondrium and runs across the epigastrium just above the umbilicus. 'It is inside.' It is not associated with tenderness or soreness of the abdomen. It has nothing at all to do with the ingestion of food; it is not accompanied by vomiting or nausea. She distinguishes it absolutely from a 'stitch in the side,' which she has had on one or two occasions on running, and from colicky pain, such as that she has had since admission to the hospital, probably due to the administration of an aperient. Though these pains may occur in exactly the same situation as the kind first described, they are both 'quite different.' She is unable, however, to explain in what way they are different. She says that they are worse than the colicky pain she has had since admission, but not so bad as a stitch in the side. The pain does not go through to the back. Since admission none of the peculiar attacks of abdominal pain have occurred."

CASE IV.

E. T., aged 16, was admitted into St. George's Hospital on February 25th, 1904, under the care of Sir Isambard Owen. She was complaining of joint pains and swellings, with sweating and headache.

Family History.—Negative.

Past Personal History.—Scarlet fever several years ago. Since the age of 14 occasional slight growing pain in the legs; never bad enough to make her limp.

Present Illness.—Began on February 18th with pain in the left ankle and right knee. She tried to do her work, that of a general servant, up to February 23rd. About this time joint pains and headache got worse, and in addition some shooting pains occurred about the upper abdomen. They were not associated with nausea or vomiting nor with the taking of any medicine, for none had been given her. The bowels were regular. The pain came on independently of food. It seemed to be inside and went across the epigastrium, but seemed to have started in the hypochondriac region, more commonly on the right side than on the left. It did not go round or through to the back at all. Though constantly recurring it did not last very long at a time. The paroxysms came on during the night, waking her out of sleep, as well as during the day. They occurred five or six times during the night and three or four times during the day. She appeared never to have had a "stitch in the side," and so cannot compare them to this pain. She had never experienced pains like them before.

Course.—Under the usual treatment the pains as well as the other symptoms of acute rheumatism subsided satisfactorily. The patient is still in the hospital, however, for on March 21st she had to be readmitted, three days after being discharged, on account of a relapse. The abdominal pain did not return during the relapse. She is now feeling well again and will soon be able to leave the hospital, for it seems that her heart has not been permanently damaged.

CASE V.

A. S. was a boy of 16, who was admitted into St. George's Hospital on March 4th under the care of Dr. Penrose. He was complaining of joint pains flying about. These, slight at first, affected chiefly the ankles and knees, and had come on originally three weeks before his admission. Shortly before admission they were associated with pain in the neck and across the shoulders, and on February 29th he got "pains in the stomach." These were not associated with anorexia, or vomiting, nor with the taking of food. Action of the bowels did not influence them,

they were not connected with the taking of a purgative, and they were unlike the pains that may occur after the administration of an aperient, though they were occasionally associated with gurgling sensations of the bowels, and once with slight nausea. The bowels had been regular. They came on in short attacks lasting a minute or two, situated in the epigastrium, not sufficiently severe to double him up, but quite sharp. They were not below the level of the umbilicus, and were not associated with soreness of the abdominal wall, but seemed to be inside. They were in the hypochondrium sometimes, more commonly on the left than on the right side. They occurred chiefly in the night while he was warm in bed. They woke him up, recurring four or five times during the night.

On admission the patient's temperature was 101°, he had a moist skin and smelt rheumatic. There was arthritis of both knees. The first sound of the heart was muffled, but the heart was not dilated. Under full doses of salicylate and the other usual remedies his pain went away. An eruption of exudative erythema on the trunk occurred on March 8th to 11th. On March 30th he was able to go home, somewhat anaemic, but fairly well.

REFERENCES.

¹ *Practitioner*, 1901; ² *Med. Soc. Trans.*, vol. xxiv, p. 24; ³ *Practitioner*, 1901, p. 61; ⁴ *Medical Diseases of Infancy and Childhood*.

ON THE BACTERIOLOGY OF SO-CALLED
"STERILIZED MILK,"

WITH SPECIAL REFERENCE TO THE MUNICIPAL SUPPLY OF
MILK FOR THE FEEDING OF INFANTS.

BY

WM. ROBERTSON, and WM. MAIR, M.B.,
M.D.Glas., D.P.H., Ch.B., B.Sc.Ed., D.P.H.Camb.,
Medical Officer of Health, Lecturer on Bacteriology, Queen's
Leith. College, Belfast.

It is now nearly ten years since Professor Flügge, of Breslau, called attention to the dangers of imperfectly sterilized milk in the feeding of infants. The subject does not appear, in this country at least, to have attracted the attention which its importance deserves. And in view of the activity of many municipalities in the direction of establishing infant milk dépôts, the present appears an opportune time to rediscuss the question. The chief results of a bacteriological examination of the milk supplied by the Leith Corporation's infant milk dépôt, given below, form the basis of the present article.

It is now, we suppose, universally admitted that the great majority of intestinal disorders of infancy occur in artificially-fed babies, the large proportion of these affections being caused by the bacterial contamination of cows' milk used as a substitute for mothers' milk. We are unfortunately still in the dark as to which group or groups of the infinite variety of bacteria occurring in milk are to be regarded as the chief offenders. Our ignorance on this point would matter less if by any convenient method we could kill all the bacteria in milk. But, as will be presently shown, this is almost, if not actually impossible, as long as we continue to adopt our present methods.

The bacteria in milk may be roughly divided into three classes, according to their powers of resisting heat.

1. Non-spore-bearing bacteria. These include the cocci, the lactic acid, coli and proteus groups. The members of this group are very easily killed by exposure to temperatures of 100° C., or as an alternative by heating for a short time at a temperature ranging between 90° C. and 95° C. This latter temperature is reached when in domestic parlance, articles of food are "brought to the boil." Pasteurization—subjecting the milk to a temperature of 70° C. for half-an-hour—will also kill these bacteria. The bacteria which are the normal inhabitants of the mammary ducts of the cow, and which cannot be avoided even with attention to the utmost care and cleanliness, belong to this group.

2. The butyric acid bacilli. These are strictly anaerobic spore-bearing bacilli. They rapidly decompose milk when stored in closely-stoppered bottles, with strongly acid reaction, coagulation and butyric acid and gas formation. Their spores are killed by an exposure of one to one hour and a-half at a temperature of 100° C. To this group belongs Klein's bacillus enteritidis sporogenes.

3. The bacillus subtilis group. These are aerobic spore-bearing bacilli, but are capable of growing under anaerobic conditions. They decompose milk with faintly acid or unchanged reaction, coagulation of the casein and subsequent liquefaction of the coagulum.

From the latter fact Flügge has called them the "peptonizing" bacilli. They are characterized by an extraordinarily rapid rate of growth at high temperatures (25° C. and up-