interest, for it depends on the production of referred pain. The appendix is supplied by nerves corresponding to the tenth dorsal segment. The same level controls the kidney, and since testis and kidney develop together their spinal segmentations are identical. The pain, therefore, in the appendix was referred to two other viscera of similar innervation, and shot from one to the other, making both tender.

In this connexion let me warn you of the manner in which the pain of an early pneumonia will sometimes simulate appendicitis. In these cases the earliest and most prominent symptom is sudden severe pain—referred pain—in the right iliac fossa; the abdomen is often rigid, and the diagnosis of appendicitis seems probable, yet if the patient be watched the abdominal symptoms mollify, and the signs of pneumonia develop at the right base.

Still more difficult diagnoses are those where the patient complains of pain which is undoubtedly referred and yet corresponds to no spinal segmentation. Such cases show the pain transgressing the limits of the spinal areas with perplexing freedom, and are illustrations of what has been called the "generalization" of referred pain. By this is meant the tendency that exists in some patients, especially those in indifferent health, for the pain gradually to spread to more and more remote areas until the original referred pain forms but a fraction of the whole. This spread is seen—I select common examples—from so simple a primary lesion as dental caries. The referred pain probably takes at first the form of a neuralgia limited to one branch of the trigeminal. This continues for a time, and then the greater part of the same side of the head is affected, and later the neuralgia attacks the opposite side. As a result the patient grows exhausted and run down, and pains begin to show themselves all over the body, especially in such places as have at previous times been painful—the joints feel rheumatic, dyspepsia accompanies digestion, or dysmenorrhoea becomes accentuated. If a patient has reached this stage when first seen, you can readily understand the difficulty of tracking the trouble to its lair. Yet the pursuit is worth undertaking, for to discover the prime cause is to convert your patient from a state of misery and dejection to one of relief and happiness.

As the most frequent causes of this generalization you should remember, in addition to hunger and fatigue, anaemia in its various forms, and hypermetropia—both fertile provocations—while in women menstruation and pregnancy fan the spread. Remember, too, that in women the spinal areas which are earliest to fall before the attack are the sixth and tenth dorsal—in other words, pain and tenderness in the breast and in the ovary.

No lecture on referred pain would be complete without reference to that interesting condition known as *allochiria*. With this uncommon symptom the patient refers stimuli applied to one part of his body to a corresponding position on the opposite side. It occurs in certain nervous diseases, and has been described in connexion with diphtheritic paralysis. Gowers has recorded a case where a stimulus was referred to both sides at once.

A last condition that bears on our subject is true neuralgia. This depends on no nerve lesion, neither is it referred from any remote condition. It is a distinct entity as a disease, and as such must be carefully distinguished from referred pain. The diagnosis may be difficult, and the points on which you must rely are the area of distribution and the presence or absence of superficial tenderness. As we saw earlier, referred pain is associated with hyperalgesia; with neuralgia this is unusual. Referred pain (and its superficial tenderness) is limited by boundaries set in the central nervous system; the distribution in neuralgia corresponds to some peripheral nerve. These distinctions, however, will sometimes fail to settle the question, and a definite diagnosis may have to be withheld until the effects of treatment have been watched.

HOLIDAY COURSES IN BERLIN.—The next "cycus" of the Berlin Dozenten-Verein for medical holiday courses will begin on September 30th and continue till October 26th. A list of the courses, etc., will be sent free, and any information required will be supplied on application to Herr Melzer, Ziegelstrasse, 10-11, Berlin.

THE TREATMENT OF LOCOMOTOR ATAXY BY FIBROLYSIN.

By F. M. POPE, M.D.,
LEICESTER.

In the following case of locomotor ataxy treatment by fibrolysin was followed by much improvement and by return of the knee-jerks.

J. B., aged 32, was admitted to Leicester Infirmary on December 20th, 1906, complaining of shooting pains in the legs and numbness of feet. He had noticed the symptoms for the past two years; had had great difficulty in walking in the dark; for the previous four weeks he had been confined to bed unable to walk at all.

Past History.

He had an "abscess on the chest" fourteen years ago, urethral discharge ten years ago; no clear history of syphilis.

Condition on Admission.

His nutrition is fair, general health fair, sleeps poorly; complains of legs feeling cold. Ill-marked areas of paraesthesia in both legs; tactile sense impaired; thermal sensation impaired but not lost; no complete analgesia; tactile conduction retarded. He is very ataxic. Romberg's symptom marked. Inco-ordination of both upper and lower extremities marked. Is quite unable either to stand or walk.

Argyll-Robertson pupil phenomena marked; no nystagmus. Left pupil smaller than right. Knee-jerks absent. Plantar reflex flexor. Has slight difficulty in micturition, none in defaecation. Facial muscles of right side slightly atrophied. Right side of tongue slightly atrophied. No morbid changes to be detected in other organs.

Treatment.

He was kept under observation for nearly a fortnight without treatment, and his condition underwent no change.

On January 2nd, 1907, treatment by intramuscular injections of fibrolysin was commenced. He received 2.3 c.cm. every alternate day.

February 12th. He has had nineteen injections. He can stand better, and walk a little with assistance. Says legs are warmer. Still much inco-ordination.

February 20th. Has had twenty-one injections. Knee-jerks are present on both sides.

March 6th. Had in all twenty-four injections. No injurious local effects. Immediate effects of each injection were a feeling of warmth and diaphoresis. He goes out to-day walking with assistance.

May 1st. He walked up to the hospital to-day. He can walk with two sticks. The knee-jerks are still present. He has no shooting pains. Pupil reaction unchanged. Achilles reflex faint but perceptible.

I have little to add to the above facts. A second case treated at the same time shows little or no improvement, but this does not, in my opinion, outweigh the positive results obtained in the first case. Ormerod, in *Allochiria's System of Medicine*, says: "We think that when ataxia has developed steadily and become thoroughly established it is likely to be permanent"; and most people will agree with him. In this case advanced ataxia has retrogressed so that the patient can get about and the knee-jerks have returned.

I regard this communication as merely a preliminary note, hoping that it may lead to a farther trial of this remedy.

ON THE CAUSES AND SIGNIFICANCE OF CERTAIN AMBIGUOUS REACTIONS OBTAINED IN TESTING URINE FOR SUGAR.

By HUGH MACLEAN, M.D.,

LECTURER ON CHEMICAL PHYSIOLOGY IN THE UNIVERSITY OF ABERDEEN.

(From the Physiological Laboratory.)

Fehling's Test.

For the detection of sugar in the urine, the test most in vogue in this country is Fehling's alkaline cupric tartrate solution. This test depends on the fact that dextrose in hot alkaline solution is capable of reducing cupric salts (which are soluble in Fehling's solution) to the cuprous form in which they are insoluble, and so separate out as a red or yellow precipitate. When the amount of sugar present is not too small this test answers admirably, and a distinct reaction with Fehling's solution indicates in almost every case that sugar is present in excess. With regard to the detection of small amounts of sugar in urine,

Fehling's test is, unfortunately, not so definite. If a normal urine, giving no reaction with this test, be taken and a small percentage of sugar added, it will generally be found on subsequent testing with Fehling's solution that no evidence of the presence of the added sugar can be obtained. It might be argued that Fehling's solution, though a good general test for sugar, is lacking in delicacy, and is therefore incapable of detecting very small quantities of this substance. This solution, however, is really one of the most delicate indicators of sugar we possess, and if some sugar be dissolved in distilled water, such minute traces as 1 in 120,000 can be distinctly demonstrated. The following table shows the relative delicacy of the more common tests for sugar.¹

	Sugar in Aqueous Solution.
Fehling's test ...	0.0008 per cent.
Trommer's test ...	0.0025 "
Nylander's test ...	0.025 "
Fermentation test ...	0.1-0.5 per cent.
Phenyl-hydrazine test ...	0.025-0.05 "
Polarimetric test ...	0.025-0.05 "

Normal urine, to which many times 0.0008 per cent. of sugar has been added, gives no result with Fehling's test, and this at once leads to the conclusion that urine contains something which prevents the sugar, when present in small amounts, giving a reaction with Fehling's solution. Pavy² explained this on the assumption that the action of the alkaline part of Fehling's solution on the nitrogenous constituents of the urine might generate free ammonia in quantity sufficient to hold the reduced suboxide in solution and so form no precipitate. The writer finds, however, that this explanation is not by any means adequate, as the amount of ammonia given off when a urine is heated with an alkali of the strength usually present in Fehling's solution is not nearly sufficient to exert any influence in inhibiting the reaction.³ In short, this effect is dependent on the fact that the kreatinin normally present in urine is capable of holding the reduced suboxide in solution, so that no apparent result is obtained. The sugar added to normal urine reduces its equivalent amount of copper, just as it does in aqueous solution; but since the kreatinin normally present is capable of holding this reduced suboxide in solution, we have no indication that any change has taken place. Thus, it is obvious that urine may contain some sugar and yet not give any indication of its presence when tested by Fehling's solution.

Kreatinin is generally mentioned in the textbooks as a substance interfering with Fehling's test. Other so-called "interfering" substances are uric acid, hippuric acid, glycuronic acid, homogentisinic acid (alkaptonuria), lactose, pentoses, and certain drug products. It is generally stated that kreatinin, like many of the other substances mentioned, interferes with the reaction in virtue of its direct reducing action, thus apparently producing a positive result when sugar is not really present. This general statement needs revision, and gives no explanation of many ambiguous reactions which from time to time make it exceedingly difficult to say whether or not a particular urine should be held to contain sugar in slight excess, or merely gives a doubtful reaction as the result of the presence of some other substance. To clear up some of these obscure points, it will be most appropriate to consider these "interfering" substances individually, in order to ascertain how and to what extent they really do interfere with the reaction. In this way it will be possible to explain many points which are at present either unexplained altogether, or, if any explanation is forthcoming, it is unsatisfactory and insufficient.

Kreatinin.

In February, 1906, I published the results of an investigation undertaken to determine the effect of kreatinin in modifying Fehling's test for sugar.⁴ In the following October an article by Kellas and Wethered⁵ appeared, in which much of this work had been repeated and the general conclusions advanced in my paper confirmed. That kreatinin constitutes the most important interfering substance constantly present in urine is undoubtedly true, but the manner in which it acts is very different from that generally described in the textbooks. Here it is considered in view of its direct reducing power on Fehling's solution, the general impres-

sion conveyed being that kreatinin, if present in excess in urine, may cause a precipitation of the reduced cuprous oxide, and so in this way simulate sugar. That this is quite erroneous is at once obvious from a consideration of the results obtained on heating solutions of pure kreatinin with Fehling's solution. Kreatinin is capable of directly reducing Fehling's solution, but this action occurs very slowly, so that an artificial solution of kreatinin equivalent in strength to what may be regarded as the maximum amount normally present in urine requires to be boiled for three to six minutes or more before it gives any appreciable precipitation. Strong solutions of kreatinin do decolorize Fehling's solution after moderate boiling, but it is almost certain that the maximum amount of kreatinin ever present in urine is of itself quite insufficient to cause any very appreciable reaction in this direction. In an adult the quantity of kreatinin passed per day is variously estimated at from 0.6 gram to somewhat over 2 grams. Neubauer found 0.6 to 1.3 grams, while Johnson gave a daily average of from 1.7 to 2.1 grams.⁶ Generally it may be taken that the average normal quantity of kreatinin daily excreted by an adult on ordinary mixed diet is, roughly, between 1 and 2 grams. Taking the daily output of urine as 1,500 c.cm., this gives an amount of kreatinin equivalent to about 0.7 mg. to 1.33 mg. per c.cm. Voit states that it may sometimes reach as high as 3 mg., or somewhat more, per c.cm. (4.9 grams per diem), but this would seem to be quite exceptional, and probably represents the maximum value. In highly concentrated urine the amount per c.cm. would, of course, be somewhat increased, but it is doubtful whether in any but very exceptional cases it is ever present in greater amount than 3 to 4 mg. per c.cm., and it is certain that this strength is only reached very rarely indeed. A small quantity of kreatinin such as might be present in normal urine gives no apparent reaction with Fehling's solution even if boiling is fairly prolonged, while such an amount as 15 mg. per c.cm., though it gives a yellow solution, gives no precipitate whatever on moderate boiling. The following table indicates the results obtained on boiling solutions of kreatinin with an equal volume of Fehling's solution for a moderate time.⁷

1 mg. kreatinin in 1 c.cm. H ₂ O...	No change even on fairly prolonged boiling; no change on cooling.
2 mg. " "	No apparent change.
3 mg. " "	No change on boiling; blue colour seemed to be very slightly less intense after cooling, but hardly apparent.
5 mg. " "	Somewhat decolorized on boiling; on cooling, reduction in intensity of colour but still fairly blue.
9 mg. " "	Decolorized on boiling; on cooling gave clear solution.
15 mg. " "	Decolorized on boiling; gave yellow colour; no precipitate, even after long standing.

This direct reducing power possessed by kreatinin is therefore of little importance in regard to its interference with Fehling's test; it is possible that in exceptional cases it might be present in amount sufficient to cause some change in the blue colour of the solution, but as it never seems to give a precipitate unless boiling is continued for a comparatively long time, this direct reducing power is unimportant, and should never give any trouble. This aspect of kreatinin may therefore be entirely neglected in urine-testing with Fehling's solution.

Kreatinin, however, exerts a very important influence in other ways. Not only, as above stated, does it prevent a small amount of sugar giving any apparent reaction with Fehling's solution, but, what is much more important from the clinical point of view, it constitutes the chief factor in the genesis of many of those ambiguous reactions which are so puzzling to the clinician. This interfering action may be considered from—

1. Its power of preventing any reaction with Fehling's solution in the presence of a small amount of sugar.
2. Its power of modifying and masking many sugar reactions when the latter is present in somewhat larger amount.

With regard to its effect in preventing a reaction with Fehling's solution, experiment shows that sugar may be

present in urine to the extent of about 0.1 per cent. or somewhat beyond this without giving any indication of its presence, the kreatinin being able to inhibit the formation of the ordinary precipitate. It is probably to this action of kreatinin that the statement to the effect that normal urine contained no sugar owed its origin. It is now held as a proved fact that all urines contain a certain amount of sugar, and in view of the above statements it is at once obvious that Fehling's test for sugar in urine is but a relative reaction, depending within certain limits on the amount of kreatinin present in relation to the amount of sugar or reducing substance. Thus, a urine very poor in kreatinin might give a fairly distinct reaction with Fehling's test, while another specimen containing more sugar, but at the same time an increased amount of kreatinin, might give no reaction. As the result of the examination of a very large number of specimens giving results more or less definite with Fehling's solution, it was generally found that a reaction really indicated slight excess of reducing substances, though, on the other hand, it sometimes happened that a urine which contained but the normal amount of reducing constituents gave a somewhat ambiguous reaction. This I interpreted as indicating a rather low percentage of kreatinin, and this seems the only satisfactory explanation; in a few cases colorimetric experiments were carried out in order to determine the amount of kreatinin actually present; this view seemed to be corroborated by the results.

More important still is the power of kreatinin in modifying certain sugar reactions with Fehling's solution. If the amount of sugar in urine is just too large to be prevented from reacting in the ordinary way, the reaction which results is much modified. After boiling for some time with Fehling's solution no result is obtained, or perhaps the mixture may be somewhat changed in colour, presenting a somewhat greenish or greenish-blue appearance; the test tube is put aside, and, after standing for some time, it is noticed that there is now a distinct greenish-yellow opalescent fluid containing a finely-divided precipitate. If the amount of sugar present is rather greater, the gradual formation of a precipitate may be evident after boiling for a fairly long time; it does not, however, form quickly, the first indication of a change being a greenish tint in the solution, which gradually becomes intensified, and passes on to give a definite dirty, milky, greenish, opalescent liquid. In some of these cases it is difficult to detect a distinct precipitate with the eye, but the centrifuge indicates that the opalescence is due to the suspension of many fine particles. In other cases where the sugar is present in fair amount (say 0.3 per cent. to 0.4 per cent. or more), no precipitate may be obtained on boiling for the usual time, but on removal from the flame or shortly afterwards, a fairly dense precipitate separates out.

On the Manner in which Kreatinin gives rise to these Ambiguous Reactions.

If a liquid containing a small amount of sugar and kreatinin be boiled with Fehling's solution, the sugar present almost immediately acts on the copper and reduces it to the suboxide. The kreatinin, however, is capable of preventing any indication of this reaction in virtue of its power of preventing the reduced copper suboxide from separating out as a precipitate. Therefore, after boiling for a few seconds we have a liquid containing kreatinin and some cuprous oxide, which is held in solution. The power of holding this lower oxide in solution is directly dependent on the amount of kreatinin present at any given moment, so that if by any process the amount of kreatinin be lessened, there is an increased tendency on the part of the suboxide to separate out as a precipitate. Now, kreatinin itself is gradually changed when acted on by Fehling's solution, part of it being probably changed into kreatin and part destroyed; if the amount of kreatinin present is just sufficient to prevent the sugar giving a precipitate, it follows that boiling for a comparatively short time will disturb the balance sufficiently to enable a small amount of the suboxide to separate out as a precipitate. This takes place gradually, and is naturally more liable to happen after the solution has cooled, the presumption being that a hot solution of kreatinin has a greater solvent power on cuprous oxide than a cold one. The suboxide at first separates out in the form of exceedingly minute particles which are held in suspension

in the blue liquid, constituting the dirty-greenish opalescent fluids so often encountered in testing certain urines with Fehling's solution. If this liquid be centrifuged for an hour or two a blue solution remains, and a greenish precipitate is found at the bottom of the tube, indicating that the opalescence was really due to a suspension of fine particles; of course there may be many modifications of this reaction depending on the amount of sugar actually present; in the presence of a fair amount the resulting precipitate may be quite easily detected as such by the naked eye. From the above it is obvious that the time for which a urine must be boiled in order to give a precipitate depends essentially (within certain limits) on the relative proportions of reducing substances and kreatinin present; the more of the latter there is in relation to reducing substances, such as sugar, the more boiling will be necessary in order to obtain a precipitate. In all normal urines, as described later, a very distinct turbid precipitate can always be obtained with Fehling's solution if boiling be continued long enough.

These ambiguous reactions are often mentioned in clinical textbooks, but there seems to be no explanation of their cause or significance. Simon,⁸ in his book on clinical diagnosis, states that not infrequently when applying Fehling's test it will be observed that upon standing, when no precipitate has occurred previously, the blue colour of the mixture changes to emerald green, while the solution at the same time becomes turbid, but that such a phenomenon should not be referred to sugar, as it is in all probability due to the action of other reducing substances. Allen,⁹ on the other hand, while admitting that a urine giving a reaction similar to that described above may contain sugar, explains the phenomenon on the assumption that some of the reducing substances of urine, such as uric acid, hippuric acid, hypoxanthine, glycuronic acid, or kreatinin, may be preventing the normal reactions of sugar. The substance, however, which causes this is really kreatinin, this effect having no relation to its reducing action. To give this modified result with moderate boiling two substances are necessary:

(a) Kreatinin.

(b) Substances reducing cupric oxide directly and capable of giving a precipitate in the ordinary way with Fehling's solution.

All the so-called interfering substances in urine, except kreatinin, act in virtue of their direct reducing power, and so, if present in excess, may simulate sugar. Kreatin, however, is an exception, as it acts in much the same way as kreatinin, only it is very much less powerful. Since, according to Folin,¹⁰ kreatin may occasionally appear in the urine in large quantities, especially after meat diet, it is likely that in certain urines kreatin may enhance to a certain extent the effects of kreatinin. In the majority of urines the amount of kreatin appears to be small and its direct effect of minor importance. Other substances also may sometimes be present which may act to a limited extent in the same manner as kreatinin, but their influence in this direction seems to be of little consequence. Thus, it is obvious that kreatinin is the substance in urine which really constitutes the greatest interfering factor in the application of Fehling's test for the detection of small amounts of sugar; it acts by its power of apparently inhibiting the reaction with relatively small percentages of sugar, as well as by the influence it exerts in giving rise to ambiguous reactions when larger amounts are present.

On the Modifications of Colour obtained on Testing Saccharine Urines with Fehling's Solution.

Every one must have observed that certain saccharine urines give with Fehling's solution a distinctly yellow precipitate, while in others it is of a pronounced red colour. Urines containing a comparatively low percentage of sugar (say, 1 per cent. to 2 per cent.) give as a rule a yellow precipitate, while others containing three or four times this amount give a distinct red. It is generally admitted that kreatinin is the substance which is responsible for this difference in colour, but how this is brought about has, so far as I know, not been explained. That kreatinin is capable of bringing about the change is obvious from a very simple experiment. If a weak solution of sugar in distilled water be treated with Fehling's solution in the ordinary way a distinct red precipitate is

obtained; if to another portion of the same solution a little kreatinin be added, and the mixture treated as before with Fehling's solution, a yellow precipitate now results. A similar result can be seen by adding, say, $\frac{1}{2}$ per cent. or more of sugar to normal urine, when on testing the precipitate is distinctly yellow. If, again, just enough sugar be added in the latter case to give a reaction after fairly prolonged boiling, the resulting precipitate is greenish, and all gradations of colour from dirty green through greenish-yellow and yellow to red may be obtained by carefully graduating the amount of sugar added to the urine. In some of my earliest experiments on this subject I found that it was always exceedingly difficult to filter off many of these yellowish precipitates, while in cases giving a greenish substance the only result obtained was a green stain on the filter paper, the filtrate being practically unchanged in appearance; the more nearly the colour of the precipitate approached to red, the easier filtration became, the latter precipitate separating with ease. In experiments with the centrifuge the results of filtration ran parallel to those of sedimentation, the green and greenish-yellow fluids taking a very long time to yield a distinct precipitate, while red ones were very quickly separated. These and other considerations led to the conclusion that the difference in colour of the precipitate really depended on the fineness of the particles, and there can be little doubt that this is the true explanation, for it can be confirmed by microscopical examination. In the presence of kreatinin even in small amount, the precipitate with Fehling's solution is modified in that it consists of finer particles; the more kreatinin present relative to the sugar, the finer the particles are; if the amount of sugar is very large (say 5 to 6 per cent. or so), the kreatinin normally present is usually insufficient to exert any appreciable influence, and the ordinary red precipitate is obtained. As above stated, when testing a slightly saccharine urine with Fehling's solution, a point arrives at which the amount of kreatinin and its derivatives present in the mixture is just insufficient to hold the reduced suboxide in solution. Naturally this suboxide is thrown out as an exceedingly fine precipitate. If heating be continued, the kreatinin becomes less and less capable of preventing a precipitate, and a yellowish-coloured turbidity is obtained; ultimately a point is reached at which the kreatinin practically exerts no influence in inhibiting the formation of a precipitate, and the tendency now is for the suboxide to separate out as a red-coloured compound. These facts are well illustrated by an experiment in which I boiled a weak solution of sugar and kreatinin until the characteristic greenish-coloured opalescent liquid was obtained; on filtering this no change was apparent in the filtrate, the filter paper being merely stained green. After centrifuging this for two hours, a greenish precipitate was obtained, the fluid part being now distinctly blue and quite clear. This blue solution was again heated, and this time the result was a distinct greenish-yellow precipitate. This was separated by means of the centrifuge and the liquid again heated; this time a yellow precipitate was obtained, which was capable of being separated by filtration. A repetition of the process eventually resulted in the filter paper showing a distinct red precipitate. These differently coloured precipitates are therefore dependent on the degree of fineness of the particles composing them, this in turn being directly dependent on, and caused by, the kreatinin normally present in the urine according to the above explanations. In the ordinary textbooks we are generally told that the yellow precipitate is the cuprous hydroxide ($\text{Cu}_2\text{OH}_2\text{O}$), while the red is characteristic of the cuprous oxide (Cu_2O). A consideration of the chemical properties of the lower hydrate of copper at once suggests a doubt with regard to this statement; if it were true that the yellow precipitate is really the lower hydroxide, this would be an easy method of obtaining a body which is one of the most unstable of chemical compounds. If, however, the yellow is really the hydrate and the red the cuprous oxide, what composition is to be ascribed to the great number of intermediate coloured precipitates obtained between green and yellow and yellow and red? There is every reason to believe that the precipitate is in every case the comparatively stable cuprous oxide, the difference in colour being no indication of a chemical difference, but merely depending on the physical state of

the precipitate with regard to subdivision of the particles. That the yellow precipitate is really the lower oxide in modified form was suggested by Dr. Angus Fraser of Aberdeen many years ago.

If a very strong solution of sugar be made up in urine the resulting precipitate may be somewhat modified and appear as a dark brown mass. This probably depends on the fact that part of the excess of sugar present is charred, while at the same time the alkali of Fehling's solution gives, with the sugar, a dark brown liquid (Moore's test). The mixture of reduced oxide, charred sugar, and dark brown liquid naturally gives a modified result, but this is essentially distinct from that obtained with moderate quantities of sugar. If a very weak solution of sugar in distilled water be examined by Fehling's test a yellow or greenish-yellow liquid is at once obtained, but here the precipitate is really red, as can be at once proved by the centrifuge; also separation by filtration is easy, and leaves a red deposit on the filter paper.

Uric Acid.

Uric acid, present in the form of urates, is another substance credited with the power of markedly interfering with the results of Fehling's test. In all textbooks this substance when present in excess is held to be a prominent factor in simulating small amounts of sugar in urine. Uric acid reduces Fehling's solution very readily, and any interfering action it may possess is due to this property. In making some experiments on this subject I found that normal urine contained more than sufficient uric acid to give a distinct reaction with Fehling's solution; the addition of a very little kreatinin, however, was quite sufficient to prevent this reaction, and showed that a considerable amount of uric acid could be prevented from giving any reaction by a comparatively small amount of kreatinin. It was obvious that the uric acid normally present in the urine was prevented from reacting with Fehling's test by the inhibitory power of the kreatinin. Experiments were conducted with a view to saturating a urine with uric acid (or urates) in an endeavour to obtain a uric acid reaction with Fehling's test. In every case this failed, and it would seem that the maximum amount of uric acid capable of being held in solution in ordinary urine is prevented from giving any reaction with Fehling's test by means of the kreatinin present. Similar experiments, in which the actual urates obtained from concentrated urine were utilized, gave negative results. These results all point to the general conclusion that urates do not generally constitute an important interfering factor in Fehling's test for sugar when this is performed in the ordinary way. After prolonged boiling they would no doubt exert a certain influence, but when boiling is fairly restricted, as is generally the case in testing by Fehling's solution, it is doubtful whether they really often cause a reaction that would likely be mistaken for sugar. There can be no doubt of the fact that many of the urines giving an ambiguous reaction are rich in urates, or at any rate show copious deposits of this substance in the urine glass. The natural inference would, of course, be that an ambiguous reaction obtained from such a urine was due to urates; it is very questionable, however, if there is sufficient ground for such a view. In urines of the above-described type, great numbers of which I have examined during the last few years, I almost invariably found that other substances, mainly of a carbohydrate nature, were present in quantity sufficient to account for the reaction.

A few in which the kreatinin was rather low were explained by this fact, as described above, the kreatinin being insufficient in amount to prevent the reducing action of the normal carbohydrate material present. That uric acid, with the help of other substances, tends to decolorize Fehling's solution to some extent is undoubtedly true, but that it is often present in urine in amount sufficient *per se* to simulate a sugar reaction in the presence of the normal kreatinin is inconsistent with the results of many observations and experiments.

How, then, it may be asked, can the fact that many urines giving an ambiguous Fehling's reaction show a copious deposit of urates on standing be accounted for? The answer seems undoubtedly to be that these urines are really concentrated and contain an excess of normal carbohydrate reducing material, which is really mainly responsible for the action. The deposit of urates is a

natural result of concentration, but really seems to be of secondary importance in the sense of interfering with Fehling's test as generally performed. It would thus seem that the usual statement to the effect that urates often interfere with the test requires modification. The maximum amount of uric acid capable of being held in solution by urine containing the normal amount of kreatinin would not exert any appreciable influence on Fehling's solution in the sense of producing a precipitate of cuprous oxide. As will be described below, a urine rich in urates and giving a reaction with Fehling's test under normal conditions is practically always found to contain sugar in slight excess of the normal amount. In such a case no doubt the urates contribute to the effect in virtue of their reducing power, but the fact that sugar is present in excess shows that the part played by the urates is not a marked one. The statement that a urine containing the normal amount of sugar and kreatinin will not usually give an ordinary Fehling's reaction though saturated with urates may be accepted as a general fact. With a little care in performing Fehling's test in urine, any influence exerted by urates may be, in a general sense, neglected; on the other hand, if boiling is prolonged they may undoubtedly assist in causing a precipitation, but it is doubtful whether boiling should ever be prolonged in performing the test. The reason for this will be discussed below.

Phosphates.

It is not often that phosphates are mentioned as interfering with Fehling's test, but that they occasionally do constitute a source of difficulty is certain. This is due to the fact that in a urine rich in certain phosphates the addition of an alkali causes a precipitation, which, as a rule, becomes much more marked on heating. In such a urine the addition of Fehling's solution may give rise to no change in the cold, but immediately on heating a greenish-yellow fine precipitate of phosphates forms, giving the fluid quite an opalescent appearance. This reaction, however, differs from that caused by a definite reduction of Fehling's solution in the fact that the phosphatic precipitate soon becomes flocculent, and separates out more or less in distinct masses. If a precipitate of this kind is obtained, it will be found that rendering the urine alkaline with some such salt as sodium carbonate, and heating, will give a precipitate of phosphates which can be easily removed by filtration. On testing the filtrate with Fehling's solution no reaction is now obtained. This interference by means of phosphates, though not perhaps very common, is a very real one when it is encountered, and exceedingly liable to cause a mistake. The accompanying

table gives an indication of what is often obtained in urines of high specific gravity, giving a certain reaction with Fehling's test. It will be seen that the amount of carbohydrate present is sufficient to give an ambiguous reaction, except in one case in which the reaction was due to phosphates.

Glycuronic Acid.

This substance undoubtedly exercises a powerful reducing action on Fehling's solution, and if present in urine in sufficient quantity would simulate sugar very closely. In urine it always appears as the paired acid (the glycuronate), and seems to be present as such to a very slight extent in normal urine; in such minute quantity it is of course of no importance, and the question to be considered is whether this substance is often present in urine in sufficient amount to simulate sugar. In a few isolated cases it has been found in abundance in the urine of apparently healthy persons, but such cases seem to be rare. As a general fact it may be stated that, in the absence of drug treatment, the presence of glycuronic acid in urine in appreciable amount is an exceedingly rare condition. On the other hand, there can be no doubt but that in certain cases, as the result of drug treatment, it really does appear in fair amount in urine, and so tends to give a more or less well-defined reaction with Fehling's test. After such drugs as chloral, morphine, naphthol, camphor, turpentine, etc., this condition may be met with. Even in these cases, however, I find that along with this substance there is sometimes an increase in the amount of fermentable sugar as well, and in cases in which a very distinct reaction was obtained sugar was generally found to be present in excess along with an appreciable amount of glycuronic acid. On the whole, this substance really seems to be of rare occurrence in urine in amount sufficient to give a Fehling's reaction, though undoubtedly it is perhaps the most important direct interfering substance likely to be encountered in quantity sufficient to closely simulate sugar. Many methods of distinguishing between sugar and glycuronic acid have been advocated, and they are so well known as to make it superfluous to describe them here. The chief of these are the phenylhydrazine reaction and the fermentation test. Both operations are tedious, and the first often requires special manipulation and experience in order to obtain a reliable result. The simplest way to distinguish between sugar and glycuronic acid seems to be to add to the urine a little tribasic lead acetate (1 to 2 grams to 50 c.cm. urine) and boil for a little time; then cool thoroughly, allow to stand, and filter. The glycuronic acid will have been almost entirely precipitated by the lead salt, while any

No.	Specific Gravity.	Results with Fehling's Solution.				Total Carbohydrate Present.		Notes.
		Before Fermentation.	After Fermentation.	After Treatment with Basic Lead Acetate.	After Allen's Modification for Precipitation of Uric Acid, etc.	Before Fermentation.	After Fermentation.	
1	1025	Fairly distinct yellowish-green precipitate on cooling after boiling for ten to twenty seconds	Faint reaction after cooling; greenish opalescence	Reaction after cooling	Reaction after cooling, but somewhat less distinct than original urine	Per Cent 0.23	Per Cent 0.14	Sample gave marked deposit of urates on standing.
2	1028	Greenish opalescent precipitate after boiling for a few seconds; became more distinct on cooling	Much fainter reaction; no distinct precipitate to naked eye; greenish-yellow opalescence	Distinct reaction after cooling	Fairly distinct reaction; more marked after cooling	0.26	0.13	Deposit of urates.
3	1018	After heating for a few seconds became fairly opalescent; on cooling, flocculent precipitate	Flocculent precipitate as before	—	—	0.11	0.06	Sample contained excess of phosphates which gave ambiguous reaction.
4	1024	Opalescence on heating; slight greenish precipitate on standing	Very faint reaction	Opalescent solution after cooling	Opalescent solution after cooling	0.19	0.10	Deposit of urates.
5	1015	Green colour on heating; distinct greenish-yellow precipitate on standing	No reaction	Fairly distinct reaction	Fairly distinct reaction	0.23	0.98	No deposit.
6	1030	Gave slight opalescence after few seconds' heating; fairly distinct precipitate on standing	Gave slight precipitate after cooling, though much less marked than unfermented urine	Gave opalescent fluid with precipitate on cooling	Distinct reaction after cooling	0.29	0.17	Deposit of urates.

sugar present will be still in solution in the filtrate. It is important to notice that the urine be of acid reaction; of course, an ammoniacal urine would be quite unsuitable, for in this case the sugar would also be precipitated. As a rule the filtrate can be quite well tested by Fehling's solution without separating off the lead; if a distinct result is now obtained the reaction is in all probability due to sugar. I have found this method very reliable as well as quick; it can be performed in a few minutes without any special apparatus. At present I am investigating the subject of glycuronic acid, and hope at a later date to publish further results both in regard to this and other substances, such as homogentisinic acid, pentoses, lactose, etc. These are, however, of comparatively rare occurrence, and only occasionally give any trouble. Lactose is only in evidence under circumstances that would probably suggest its presence in the urine of lactating women.

The Reducing Power of Normal Urine on Fehling's Solution.

Every urine will give a distinct precipitate with Fehling's solution if boiled long enough.¹¹ The time required varies from an average of two to five minutes, or more, depending on the quantity of reducing substances normally present. Pathological urines rich in dextrose give a marked reaction even before the boiling point is reached, while the presence of a minute amount may necessitate boiling for some time in order to obtain any indication of the presence of sugar. These modifications are caused by the kreatinin present, as will be obvious from the facts already stated. If no kreatinin were present, all normal urines would at once give a fairly distinct reaction with Fehling's solution, even the normal quantity of uric acid being much greater in amount than is necessary to cause this. Besides uric acid, normal urine always contains a certain amount of reducing carbohydrate material (of which part, at any rate, is dextrose) as well as minute traces of other substances, such as glycuronic acid compounds. All these tend to cause almost immediate reduction after the boiling point is reached, and in the percentage in which they occur in urine reduce their equivalent amount of Fehling's solution in a few seconds; in the process of reduction they undergo destruction.

Kreatinin, on the other hand, while holding in solution the reduction products of the above substances, gradually exerts a reducing action of its own. This occurs slowly, however, as is proved by the fact that a solution of kreatinin containing approximately the percentage of this substance found in normal urine requires to be boiled with Fehling's solution for many minutes before any precipitate is obtained.

If, therefore, we boil a mixture of Fehling's solution and normal urine for a short time, we have a fluid containing a certain amount of cuprous oxide which is held in solution by the kreatinin; on continued boiling the kreatinin itself continues gradually to add to the amount of the lower oxide present, in virtue of its own reducing power, while, at the same time, it becomes gradually less in amount. Ultimately a point is reached at which the suboxide reduced by the uric acid, sugar, etc., of the urine, added to that reduced by the kreatinin itself, is no longer capable of being held in solution by the amount of kreatinin and its derivatives actually present in the urine at that moment; at this point a precipitation occurs.

Thus it will be seen that the reaction obtained from normal urine is essentially the same in kind, and only differs in degree, from that obtained in a urine containing more than the normal amount of sugar, the chief difference being that the more sugar present the quicker the reaction occurs.

On the General Significance of these Ambiguous Reactions.

The question of the significance of many of these modified reactions in urine probably presents itself most forcibly in the examination of cases for life insurance. In examining the urine of a candidate, it fairly often happens that a reaction is obtained with Fehling's solution, which, though not a typical sugar reaction, leaves the examiner in great doubt as to whether he is justified in saying that the urine is free from sugar. As life insurance companies generally hold that the presence of a trace of sugar in the urine is sufficient to disqualify the applicant, the question is rendered still more difficult, strictly speaking, every urine contains a certain amount

of sugar, which, as explained above, is prevented from giving a reaction with Fehling's test by means of the kreatinin present. The kreatinin, however, is not capable of inhibiting the effect of considerable amounts of sugar, even when it is itself present in increased amount, and if the percentage of normal carbohydrates be increased from any cause, then a modified result will be obtained with Fehling's test. Now, ordinary urine contains on an average carbohydrate matter to the extent of from 0.05 per cent. to 0.1 per cent., part of this being dextrose. Such a urine would normally give no reaction, for the reasons already given. If such a urine of specific gravity 1015 be evaporated to about half its bulk, its specific gravity would then be from 1028 to 1030. Such a concentrated urine would contain from 0.1 per cent. to 0.2 per cent. carbohydrate material, and the latter amount would usually be quite sufficient to give an ambiguous Fehling's reaction, though the urine was otherwise quite normal.

In testing a urine the first consideration to be taken into account is the specific gravity. It is often the case that these doubtful urines are of high specific gravity, and consequently contain more normal carbohydrate than usual. In the days when the presence of sugar in urine was denied, this concentration was of course held to have no effect, and the ambiguous reaction obtained from such a urine was either ascribed to urates or some other reducing substance, or if sugar was suspected, the urine was held to be abnormal. Many quite normal concentrated urines, however, give an ambiguous result with Fehling's test, and the following observations may help in a doubtful case.

If the specific gravity of the urine is high, it ought to be reduced by the addition of water to about 1015. In the case of a urine of from 1028 to 1030 specific gravity an equal volume of water would be required to effect this. If now it gives a distinct reaction when boiled with an equal volume of Fehling's solution for about ten seconds either immediately or on standing for a minute or two, it may generally be assumed that sugar is present in quantity above the average for normal urine. Unless at least a distinct opalescent fluid with a greenish or yellowish precipitate is obtained, the test should be held as negative for practical purposes, since the amount of sugar necessary to cause decolorization and yet not give a precipitate is likely to be of no clinical importance. In the absence of drug treatment an ambiguous reaction with Fehling's test generally indicates, in a urine of average specific gravity (1015), a slight excess of sugar. In connexion with specific gravity, it should be remembered that a urine giving an indication of 1020 when passed, will, on cooling to the temperature of the room, show a specific gravity of 1025 to 1026. In insurance cases the urine is generally tested immediately after it is passed, and this fact in regard to specific gravity is often lost sight of, and consequently leads to error.

If all urines, after cooling to room temperature, were diluted to about 1015, many mistakes might be avoided; a reaction in such a dilute urine would always be obtained with less than 0.15 per cent. dextrose. In urines containing a slightly higher percentage of sugar than normal the important point, of course, is to determine how much urine is passed per diem, and so obtain a clue as to the total sugar passed. An ambiguous reaction with Fehling's test in the urine of an individual passing less than 1,000 c.cm. per day would obviously have a different significance to that of a urine giving a somewhat similar reaction but of which the daily average amount was nearly 2,000 c.cm., or even over this. In a case giving a modified reaction with Fehling's test after dilution it is always advisable to postpone a report until a future examination can be made, for it seems certain that the presence of small amounts of sugar in urine in quantities above the normal is often caused by factors which have very little direct bearing on the probability of life. The amount of exercise taken, the quantity of carbohydrate eaten, working or sitting in badly-ventilated rooms or offices, temporary feverish or mentally-disturbed conditions, alcohol, etc., all tend to cause a slight increase in the amount of sugar excreted in the urine of many apparently healthy individuals.

Again, observation proves that sugar may be occasionally present in the urine of healthy people without any apparent cause, and that the power of assimilation of carbo-

hydrate may vary within fairly wide limits, not only in different individuals, but in the same individual. The important point to consider is whether, with ordinary diet, the urine always contains a small excess of sugar. If it does so, the case is suspicious, and the ultimate prognosis very doubtful—probably bad. If the sugar is of but temporary occurrence, it can be said that in the majority of cases its presence is of no importance. This temporary increase of sugar is often observed in cases of mental disturbance. The teaching that urines containing sugar are of high specific gravity, while generally true, is not always so. In connexion with this, it is interesting to note that a sample kindly sent me by Dr. A. W. Macintosh, of the Aberdeen Royal Infirmary, contained about 1 per cent. sugar, and yet the specific gravity was, on passing, only 1005, and after cooling, 1010. On the other hand, urines of high specific gravity, such as 1030, suggest at once a suspicion as to the probable presence of sugar. This, of course, is the ordinary textbook statement, but, unfortunately, is often misleading. If, on testing such a urine with Fehling's solution, a modified reaction is obtained, this is often held as confirmatory. As a matter of fact, almost all urines of such high specific gravity as 1030 give some reaction with Fehling's test, but a little consideration would at once show that if the high density were really due to sugar, then the amount necessary to cause this would give a very decided reaction indeed. If glycuronic acid is suspected it can be separated by basic lead acetate. In doubtful cases other tests must be made use of, and in a future publication I intend to give an account of my experience of some of these tests and their results, as well as some methods by which the exact amount of sugar present can be determined. The above will serve to show the nature of Fehling's reaction, and to render intelligible certain reactions of which I have not hitherto seen an explanation. The fact that every urine contains a certain amount of sugar; that the amount normally present varies with different individuals and probably for the same individual, and that the test is at best but a relative one, indicates that the occurrence of an ambiguous reaction in a concentrated urine is generally of no clinical importance, and though largely due to carbohydrates does not usually indicate an abnormal condition. In urines of low specific gravity these ambiguous reactions do generally indicate a slight excess of sugar, but its probable bearing and importance can only be ascertained by watching the case, and by repeated examination of the urine at short intervals. In many cases it will be found to disappear, and give no further trouble; in a few it may show the starting point of a true diabetes mellitus, and in such cases early detection of the true condition is exceedingly important.

In conclusion, I would express my thanks to the various medical men who were kind enough to supply me with samples of urine from time to time during the last few years, and would particularly mention Drs. Souter, Copeland, and Chalmers, residents at the Aberdeen Royal Infirmary.

REFERENCES.

- ¹ Simon, *Clinical Diagnosis*, 1904, p. 522. ² Pavy, *The Physiology of the Carbohydrates*. ³ MacLean, *Biochemical Journal*, 11, 4 (March, 1907). ⁴ MacLean, *ibid.*, 1, 2 (February, 1906). ⁵ Keilas and Wethered, *Lancet*, October 20th and 27th, 1905. ⁶ Huppert-Neubauer, *Harnanalyse*, 10 Aufl., S. 387. ⁷ MacLean, *Biochemical Journal*, 1, 2 (February, 1906). ⁸ Simon, *loc. cit.* ⁹ Allen, *Chemistry of Urine*, 1895, p. 61. ¹⁰ Folin, *Zell. f. Physiol. Chem.*, 1904, Bd. XII, pp. 223-242. ¹¹ MacLean, *Biochemical Journal*, March, 1907.

LIGATURE OF COMMON CAROTID FOR ARTERIAL DILATATION AND EPILEPSY.*

By RUSHTON PARKER, M.B., B.S., F.R.C.S.,
PROFESSOR OF SURGERY IN THE UNIVERSITY OF LIVERPOOL.

The following case of arterial dilatation of the left side of the head, attended by epilepsy and right hemiparesis, relieved by ligature of the left common carotid, is of interest from several points of view:

An unmarried woman, aged 32, was admitted into the Liverpool Royal Infirmary in November, 1905, suffering from a dilated and strongly pulsating condition of the arteries of the

* Read at a joint meeting of the Lancashire and Cheshire and North Wales Branches of the British Medical Association at Chester, April 10th, 1907.

left side of the head for fifteen years, epilepsy for ten years, some frontal headache, and a dull expression, with slow mental processes. It was presumed that similar vascular derangement existed inside the skull, affecting portions of the brain, and probably accounting for the epilepsy. She was put into a medical ward, under the care of Dr. Abram, for observation and report previous to operation.

It was there ascertained that the dilated vessels were first noticed on the left forehead at the age of 17, and had since spread over that side of the head, with increasing pulsation and headache. The fits began at the age of 22, and had latterly been as many as thirteen in a week. There was no aura, but complete loss of consciousness. She had bitten her tongue, but had not passed urine or faeces during a fit. There was a certain amount of right hemiparesis, but sensation was normal; Babinski's sign and ankle clonus were present in the right foot and triceps-jerk in the right arm. The pupils varied and the fundus oculi was normal. Besides the great general dilatation of blood vessels on the left side of the head, confined to the arteries, and specially affecting the main trunk, there was a thrill felt and a murmur heard. The vessels elsewhere were normal. Four days after admission she had six fits, which appeared to start in the right leg.

In the matter of treatment by operation, as the vascular changes outside the skull involved branches of the external carotid, and as those inside might possibly have concerned the middle meningeal—itsself also a branch of the external carotid—it was suggested by Dr. Abram that ligature of this division of the main trunk might wisely be attempted.

With that view on November 27th, 1905, seventeen days after admission, the left common carotid was exposed and its bifurcation sought in the usual place, but no sign of it found there. The wound had been freely made in the line of the vessel, and was now extended at right angles. It was eventually found that there was an unusually high division of the carotid at the level of the jaw. The external carotid was barely within reach of the finger and could hardly, if at all, have been accessible for ligature. Moreover, pressure on it did not control the strong pulsations outside the head. They were only allayed by compression of the common carotid, which was then tied with prepared catgut. The vessel was much widened, as surmised before operation.

The wound healed simply and quickly by first intention; the patient was immediately relieved of her headache and all other inconvenience, and remained without fits for a time. On April 5th, 1907, she came to hospital, and was then examined by Dr. Abram and Mr. Parker. Her sister said that the fits returned about three months after operation, but much more mildly and seldom. Although during the period after their return she had sometimes been free for as much as three weeks at a time, they had latterly become more frequent, and now numbered about one a week. Her memory, also, and power of speech were failing; but in other respects she was healthy and cheerful, in great contrast with her condition before operation. Examination of the fundus oculi showed again no abnormality, and Babinski's sign was no longer present in the right great toe.

The total disappearance of outside arterial dilatation, the cessation of wild pulsation, mitigation of the right-sided hemiparesis, together with the improvement in her appearance and comfort, are quite conspicuous; but the ten years' epilepsy before operation has permanently affected her mental powers, the further deterioration of which was notably arrested during the three months' total freedom from fits. The vascular changes are not likely to have been confined to those perceived outside the left side of the head; the accompaniment of right hemiparesis, epilepsy and headache having been probably due to some such internal derangement.

ON THE CULTIVATION OF THE MENINGOCOCCUS IN THE PRESENT EPIDEMIC OF CEREBRO-SPINAL MENINGITIS IN BELFAST.

BY

WM. ST. CLAIR SYMMERS, and W. JAMES WILSON,

M.B. ABERD.,
MUSGRAVE PROFESSOR OF
PATHOLOGY, QUEEN'S
COLLEGE, BELFAST.

B.A., M.B., D.P.H. CANTAB.,
RIDDEL DEMONSTRATOR
OF PATHOLOGY,
QUEEN'S COLLEGE, BELFAST.

DURING the present epidemic of cerebro-spinal meningitis in Belfast we have made bacteriological examinations of the fluid obtained by lumbar puncture during life (that is, during the first week of illness) from 31 cases, from 27 of which we obtained the *Diplococcus meningitidis* of Weichselbaum by culture on appropriate media. (See below.) One of the negative cases showed intracellular, Gram-negative cocci in the pus cells; and another negative