which the following is an example: the patient, aged 40, had true diabetes with neuritis.

TABLE III.—Blood Sugar Percentage

	Before Glucose	$\frac{1}{2}$ Hour After	1½ Hours After	2½ Hours After
Arterial	0.236	0.354	0.526	0.466
Venous	0.207	0.332	0.455	0.434
Difference... ..	+0.0.9	+0.022	+0.071	+0.032

The following figures, obtained from a patient, aged 31, with true rheumatoid arthritis, illustrate the disproportionately small amount of sugar which may be passed in the urine, in spite of high blood sugar values, showing that the renal threshold is altered in cases of this disease.

TABLE IV.—Blood Sugar Percentage

	Before Glucose	$\frac{1}{2}$ Hour After	1½ Hours After	2½ Hours After
Finger blood	0.138	0.300	0.247	0.183
Vein	0.121	0.280	0.261	0.150
Difference... ..	+0.017	+0.020	-0.014	+0.033
Urine sugar	Absent	A trace	Faint trace	Very faint trace

Curves similar to the above were obtained in practically all cases of suspected rheumatoid arthritis diagnosed clinically, and as such curves do not occur in any other type of arthritis investigated a glucose tolerance test would seem to be a valuable confirmatory means of diagnosis. Further work is being carried out on this point.

With the subsequent introduction of insulin into the treatment of rheumatoid arthritis^{2, 3} the question arose as to whether, *pari passu* with the manifest clinical improvement, any alterations in the glucose tolerance occurred such as are seen in many cases of diabetes after treatment. In several patients with true rheumatoid arthritis multiple curves were plotted at varying stages, the later curves often being after several months' treatment with insulin (in most instances with great benefit clinically), but in no case was any remarkable improvement in the glucose tolerance curve observed. Indeed, subsequent curves generally showed striking resemblances to those previously obtained. Table V shows the figures obtained in one of the few cases in which there was apparent improvement in sugar tolerance after treatment. The patient was 50 years old, and the interval between the two observations nine months.

TABLE V.—Blood Sugar Percentage

	Before Glucose	$\frac{1}{2}$ Hour After	1½ Hours After	2½ Hours After
Finger... ..	0.121	0.235	0.250	0.133
Vein	0.110	0.190	0.211	0.125
Difference... ..	+0.011	+0.045	+0.039	+0.008
Nine months later				
Finger... ..	0.111	0.166	0.220	0.162
Vein	0.083	0.150	0.181	0.146
Difference... ..	+0.028	+0.016	+0.016	+0.016

We wish to record our thanks to our colleagues on the medical staff of the B.R.C.S. Clinic for rheumatism for permission to investigate and record findings on their patients in addition to our own.

REFERENCES

- ¹ Pemberton, R.: *Arthritis and Rheumatoid Conditions*, Lea and Febiger (U.S.A.), 1929.
- ² Howitt, F. D., and Christie, W. F.: *Lancet*, December, 1931.
- ³ Copeman, W. S. C.: *British Medical Journal*, December 19th, 1931.

FATAL STAPHYLOCOCCAL SEPTICAEMIA WITH PSEUDOMYXOMA PERITONEI

BY

OCTAVIA WILBERFORCE, M.R.C.S., L.R.C.P.

ASSISTANT PHYSICIAN, NEW SUSSEX HOSPITAL FOR WOMEN,
BRIGHTON

The following case seems worth recording on account of: (1) the abdominal condition: few symptoms or physical signs were evident till the disease was advanced—papillomatous cystadenoma of ovary followed by pseudomyxoma peritonei; (2) the vomiting, acidosis, and unexplained mental upset with subsequent complete restoration of clearness of mind; and (3) the rapid, fatal staphylococcal infection which spread to the brain following what the patient called a "common cold sore" on the upper lip.

History.—The patient, an unmarried woman of 58, consulted me in October, 1930, for indefinite symptoms of headache, fatigue, slight loss of weight, occasional backache, heartburn, and flatulence. She said she had always been subject to gastric trouble, and for several years had been obliged to take aperients. She did not complain of abdominal pain; recently she had not been sleeping well. In 1905 she was operated on for "appendicitis and colitis." A horseshoe kidney was found; the appendix was not removed. Though never a robust individual, she had had no other serious illnesses. Menopause occurred several years past; no discharge since.

On examination I could find nothing definitely abnormal. I prescribed dyspeptic mixtures, and decided to suggest a barium meal if symptoms failed to improve. I saw her on two further occasions before Christmas; while her weight and general condition seemed to have undergone no marked change, she was no better. I next saw her on January 12th, 1931. She had had a slight blood-stained discharge per vaginam on the previous day, and had been passing very little urine. A large cystic abdominal tumour, filling nearly the whole of the right side of the abdomen, was palpable, and I arranged for her admission to the New Sussex Hospital. The blood picture on January 16th was: red cells, 5,152,000 per c.mm.; haemoglobin, 88 per cent.; colour index, 0.86; white cells, 9,800 per c.mm.—polymorphs 60 per cent., small lymphocytes 32 per cent., large lymphocytes 3 per cent., hyalines 4 per cent., and eosinophils 1 per cent.

OPERATION

An operation was performed on January 17th. A paramedian subumbilical incision was made, and the peritoneal cavity was found to contain a pint or more of blood-stained free fluid. A large right papillomatous cystadenoma of ovary, about the size of a cricket ball, was removed after ligation of the pedicle. A left ovarian cyst of similar consistency, although smaller, was also resected. Small nodules of friable pseudo-mucinous growth covered the whole of the broad ligaments, bladder, and uterus, and extended as a widespread jelly-like peritonitis over the pelvic peritoneum, mesentery, mesocolon, omentum, and left lobe of the liver. There were many adhesions, and there was considerable matting together of the intestines. The uterus was normal in size and position. The appendix was enlarged, being 3 cm. in diameter, and was covered with similar growth. An appendicectomy was performed. There was a large horseshoe-shaped single kidney, consisting of a large left and small right; the median part of it curved downwards, the convexity being downwards and forwards.

X-RAY TREATMENT

In view of the marked tendency of the tumours to extend to adjacent parts by implantation and invasion of the surrounding tissues, and of the fact that such pseudomyxoma peritonei may become clinically malignant, it was decided to give intensive x-ray treatment, in the hope of preventing further extension. Accordingly, on March 26th a total depth dosage of 110.6 per cent. was given through two large anterior and two posterior fields; and on July 31st a total depth dose of 94.3 per cent. through one anterior, one posterior, and two lateral fields. A voltage of 186,000 was employed—the focal

distance being 30 cm.—and a filter of 0.5 mm. copper + 2 mm. aluminium was used.

Sections showed papillomatous cystadenoma of ovary. Some of the cysts revealed small papilliferous ingrowths, and in other parts there was a more glandular structure, but no evidence of malignancy. Appendix showed chronic obliterative appendicitis, with growth adherent to the surfaces. The patient made an uneventful recovery.

SECOND ADMISSION TO HOSPITAL

On June 18th she came to me complaining of heartburn, fatigue and bad nights; she had gained 2 lb. in weight. The abdomen was not, however, markedly fuller. I saw her again on June 28th: she felt ill and had vomited slightly. On July 2nd she was very poorly, with slight pain in epigastrium; she had had one large vomit. There was much distension, and she was constipated in spite of many aperients. She was admitted to hospital on July 9th. The bowels were eventually opened, but a good deal of distension remained. She was seen in consultation on July 11th and a large mass palpable in Douglas's pouch was found. For the preceding eight days she had been on light diet, fluids, glucose in her drinks, and two-hourly feeds. By July 13th the abdominal distension was definitely increasing although the bowels were acting well.

On July 16th the patient had a disturbed night; she talked nonsense and was very restless. Sedobrol was given, and later luminal, but without effect; finally, at 1.30 a.m., an injection of 1/100 grain of hyoscine hydrobromide was given, but the patient did not quiet down until 3.45. When I saw her later in the morning she was much distressed, said people were saying she was out of her mind, that she saw coloured lights, and that there had been a great noise in the night, etc. Her blood pressure was 120/90, pupils dilated, disks normal, and I could find nothing abnormal in her central nervous system. On July 18th the mental condition was very bad, and for the first time acetone was present in her urine. The following day the mental condition was still very bad, and the pulse rapid—110, and soft.

On July 20th the urine contained no acetone, and the patient's mental condition became quite rational. Acetone was again present the next day, but in lesser degree. The patient's mind was clear, and indeed never became clouded again.

During the period July 13th to 20th the temperature rose to 99° on several evenings, and the pulse ranged from 92 to 110. The distension increased rather than lessened, but gradually she was able to take light, solid food, which was digested without discomfort. She had a long talk with me as to the queer "nightmares," the details of which she remembered clearly. She left the hospital on July 29th and had more intensive x-ray treatment, which to some extent reduced the abdominal distension.

FATAL STAPHYLOCOCCAL INFECTION

On September 9th I was sent for urgently to Worthing, where she had just moved for a month's change of air. Five days earlier she had noticed what she called a small "cold sore" on the outer right side of her upper lip. This she had not squeezed, but had painted with tincture of iodine; later, she had used hot fomentations. The swelling had suddenly spread upwards into the neighbouring cheek, her temperature had risen to 101°, pulse to 112, and the general appearance of the face at first glance almost suggested erysipelas, but there was no defined margin or raised edge. I moved her again into the New Sussex Hospital, and the following day an incision under N₂O was made. Film and culture showed a growth of *Staphylococcus aureus*. Sodium cacodylate hypodermically, mercury perchloride hypodermically, salines and glucose per rectum and later intravenously, entirely failed to check the infection, which spread rapidly, causing proptosis of both eyes just before her death on September 14th. Her temperature swung in a typically septic fashion, rising finally to 107.4°, accompanied by a relative increase in pulse rate.

ABSTRACT OF POST-MORTEM FINDINGS

The septic condition of the face extended through both cavernous sinuses, which contained pus and ante-mortem thrombi, to the brain, where pus was found on the under surface of the pons. There were petechial haemorrhages in the gastro-intestinal tract, which contained altered blood, and

the spleen was diffuent. A metastatic abscess was found on the right middle lobe of the lung. The peritoneum contained several pints of yellow fluid, and the omentum was thick with a myxomatous material resembling sago or frog spawn, and lined by a single layer of flat cubical epithelium. A horseshoe kidney was present, the ureter being from the left half, which was in normal position, the right half being on the aorta.

I am indebted to my colleagues Dr. Louise Martindale for the detailed report of her operative measures and intensive x-ray treatments, and Dr. Mary Leslie-Smith for the pathological and post-mortem findings.

Memoranda

MEDICAL, SURGICAL, OBSTETRICAL

ACUTE STREPTOCOCCAL PERITONITIS IN CHILDREN

In the *British Medical Journal* of October 15th, 1932 (p. 704), Mr. L. E. Barrington-Ward states:

In my experience, recovery with or without subacute encysted intraperitoneal collections never occurs. Operation has no effect. In fifteen years there have been twenty-two cases at Great Ormond Street: twelve were operated upon, ten were not, but all terminated fatally.

Although we agree with Mr. Barrington-Ward that acute streptococcal peritonitis is one of the most tragic of all abdominal catastrophes, we think there is frequently a chance of recovery. During the last four years we have seen eight cases, three of which have lived. All were treated by operation.

CASE I

Female, aged 3. She had complained of increasing abdominal pain for the previous twelve hours. On admission to a nursing home (January 28th, 1929) the temperature was 101.5° F., the pulse rate 120, and the respirations 26. Her chest and back were covered by a scarlatiniform rash. Rigidity and tenderness were present over the lower abdomen. Immediate operation disclosed an early pelvic peritonitis with pus. The appendix, which appeared normal, was rapidly removed and the pelvis drained. During the operation the child had clonic contractions of both legs and arms, and a convulsion which lasted for almost fifteen minutes. Following the operation, she received large doses of anti-streptococcal serum, and, after an acute illness of seven days, made a rapid recovery. The pus gave a pure culture of haemolytic streptococci.

CASE II

Male, aged 7. He had had acute abdominal symptoms for two days. On admission to hospital (July 8th, 1929) the temperature was 103.4° F., the pulse rate 128, and the respirations 38. When the abdomen was opened a large quantity of pus escaped. The appendix, which was not inflamed, was removed and the pelvis drained. In three days he appeared to be out of danger, and made an uneventful recovery. Non-haemolytic streptococci, of the *Streptococcus faecalis* type, were grown from the pus.

CASE III

Male, aged 4½. He was admitted to a medical ward on April 4th, 1932. The boy complained of pain in the right loin, of drowsiness, and inability to walk, and had had two convulsions the day before. The temperature was 102° F., the pulse rate 126, and the respirations 28. Nothing abnormal was found on examining the chest or central nervous system, but the right side of the abdomen was rigid and tender. Three weeks later the abdominal picture had not changed. The temperature was 103°, the pulse rate 144, and the respirations 40. Operation was decided on, and an abscess, involving the omentum, was found beneath the liver on the right side. After being acutely ill for two weeks, he gradually recovered. Non-haemolytic streptococci were grown from the pus.

We are of the opinion that a suitably timed operation distinctly improves a child's chance of recovery.

G. GORDON BRUCE.
NORMAN J. LOGIE.

Aberdeen.