we found a leaking aneurysm of the left side of the circle of Willis, which had burst.

In another case in my department, under Mr. Angell James, mastoiditis was simulated and an operation performed; the lesion was an aneurysm of the basilar artery.

Oöphoritis appears occasionally to produce earache. Sometimes we are compelled to regard the earache as functional in origin, as in the case described below.

CASE IX.—*Neurotic "Earache"*

R. N., a female aged 24, was admitted as a case of mastoiditis, with severe left earache, intense tenderness to lightest touch on mastoid and behind, giddiness, and falling. No nystagmus or nausea. Temperature 98.6° F., pulse 76; hearing normal, tympanic membrane normal. She seemed quite quiet and intelligent, and anxious to have an operation—"anything to get rid of the pain." Careful examination revealed no physical signs. I made the diagnosis of quasi-hysterical neurosis, which rather startled her own doctor. When told that no operation was necessary she lost all pain. Later I found out that she had just lost her job. Three years before she had suffered from hysteria, her betrothed having married her best friend just before the date fixed for her own wedding.

REFERENCES

- ¹ Clark and Taylor: Tic Douloureux of the Genuiculate Ganglion, *Journ. Nerv. and Ment. Dis.*, 1910, 242, 511; Harris, W.: Genuiculate or Otic Neuralgia, *Journ. Amer. Med. Assoc.*, 1914, 1728.
- ² Hunt, J. R.: Sensory Field of Facial Nerve, *Brain*, 1915, 418. See also Aitken, R. S., and Brain, R. T.: *Lancet*, 1933, i, 19.
- ³ McKenzie, D.: Herpes Oticus with Recurrent Laryngeal Paralysis, *Journ. Laryngol.*, 1915, 339.
- ⁴ Tilley, H.: Symptoms, etc., of Chronic Suppuration in Sphenoidal Sinus, *British Medical Journal*, November, 1905.
- ⁵ Watson-Williams, E.: Mastoid Disease without Otorrhoea, *ibid.*, December, 1929.
- ⁶ *Idem*: Septic Meningitis, *Practitioner*, December, 1931, 659.

THE LATENT MASTOID AS A CAUSE OF CHRONIC AURAL SEPSIS

BY

GILBERT CHUBB, D.Sc., F.R.C.S.

SURGEON IN CHARGE OF THE EAR AND THROAT DEPARTMENT,
WESTMINSTER HOSPITAL

In the vast majority of cases of acute otitis media the path of infection is via the Eustachian tube. Owing to the early closure of the latter, the inflammatory exudate is under pressure. It must be rare indeed for the mastoid antrum to escape infection.

The acute mastoiditis which occasionally supervenes may present the classical signs of mastoid involvement. Often, however, these are conspicuous by their absence. The only definite physical sign may be the persistence of the meatal discharge. Reluctantly we operate, and we find a frankly purulent mastoid. Many such cases must be missed. What becomes of them? The answer is given by the chronic discharging ears that come to the radical mastoid operation. Whatever may be the condition determining operation in these latter cases, however completely mastoid signs and symptoms may be absent, they almost all agree in presenting at the operation unequivocal indications of old-standing mastoid infection.

When we consider the occasional, almost fortuitous, discovery of the more or less latent mastoid infection in the recent case, and the almost constant indications of prolonged latency of the mastoid infection in the old-standing case, is not the conclusion inevitable that these latter represent the mastoid extension of the original acute infection which, having run throughout a latent course, has been missed? If this is so, if these latent mastoid infections remain after the subsidence of the acute otitis media, then our failure to diagnose them is responsible for the great bulk of chronic aural sepsis, with its progressive deafness and potential dangers.

How far the mastoid bone with its diploic tissue or air cells is involved will depend on the virulence of the infection, and on the resisting power of the patient. It will, however, depend also on the rapidity with which drainage is established. Rupture or paracentesis of the drum is at the best imperfect drainage for the antrum. With partial obstruction of the aditus it becomes still less efficient. With actual involvement of the mastoid bone this drainage can do little more than serve to maintain the latency of the mastoid infection.

The discharge of chronic aural sepsis may show prolonged periods of complete remission. A scarred drum alone remains to tell the tale of previous active sepsis. As there is no reasonable ground for denying this complete latency in the case of the mastoid infection, while admitting it, as we must, in the case of the tympanic cavity, we are driven to the conclusion that even the absence of a discharge is no certain ground for assuming a non-infected mastoid. Indeed, we have only to postulate the early and complete closure of the aditus to envisage the possibility of a chronic but entirely latent mastoid infection, coexisting side by side with a normal middle ear, normal drum, and normal hearing. The two following cases suggest that this actually occurs:

CASE I

A boy of about 12 was sent in with pain, tenderness, and marked swelling over the right mastoid, with a corresponding displacement of the pinna. The drum and external meatus were, however, absolutely normal. So also was the hearing. There was nothing on the scalp, and no concurrent disease to suggest a blood-borne infection. Operation disclosed a mastoid filled with pus.

CASE II

A woman of about 40 came to me with pain in the right ear of recent onset. There was no previous history of ear trouble. She had a red and bulging, but otherwise normal, drum, which I incised. Some fourteen days later there were signs of mastoid involvement. At the operation the mastoid presented every appearance of an infection of many years' standing.

Neither of these cases definitely proves that the mastoid infection originated in an acute otitis media, which, as far as the middle ear was concerned, cleared up completely. Blood-borne infections of the mastoid are, however, so rare and infections via the Eustachian tube so common that this explanation is at least a probability.

These cases must be rare, and would, in any case, be difficult to diagnose till physical signs arose. Cases in which a discharge persists after the acute stage has subsided are very much more common. They represent nearly all cases of chronic suppurative otitis media. In how many of these is the persistence of the discharge due to the presence of residual mastoid sepsis—sepsis latent save for the discharge? My experience suggests that it is in a very large proportion of them. The following cases lend support to that view:

CASE III

A girl of 12, with a copious discharge from both ears, dating from an attack of measles eighteen months previously, developed a swinging temperature. As nothing could be found to account for this, I was called in because of the aural discharge. By that time the temperature had been swinging for eleven days. A bilateral cortical operation showed both mastoids full of pus and the left lateral sinus exposed in the abscess. She recovered.

CASE IV

Another child, of about the same age, came to the outpatient department with a history of a discharging ear and some very doubtful signs of mastoid involvement. She reappeared a week later, desperately ill, with marked adenitis round the right jugular and a swinging temperature. Immediate operation on the mastoid and thrombosed lateral sinus failed to save her, and she died of meningitis.

CASE V

In pleasant contrast to the two preceding cases this boy of 16 was sent in on the strength of one sharp rise of temperature the same day. He had a small perforation in the left drum and a history of intermittent discharge from that ear. There was no tenderness or other sign of mastoid involvement. Immediate operation disclosed a septic mastoid, with pus and granulations on the exposed lateral sinus. He recovered.

Such cases could be multiplied indefinitely. They carry a twofold and very obvious lesson—namely, that a suppurating mastoid can run a completely latent course save for the ear discharge, and that when complications arise they may do so without any warning from the mastoid.

CASE VI

A boy of 10 was sent to me from the ophthalmic department with the urgent request that any focal sepsis should be discovered and eliminated. All that I could find was a small perforation in the left drum and a history of discharge. There was no tenderness, no pain, and no discharge when I saw him. In view of the urgency of the eye condition I opened the mastoid. It was full of pus.

CASE VII

A man of 20 came under my care with bilateral acute otitis media. A paracentesis of both drums resulted in his temperature becoming normal. Pain disappeared, and the left ear recovered without further intervention. The right ear on the tenth day still showed a pulsating discharge in the tympanic perforation. There was no pain, very doubtful tenderness, and the temperature was normal. Operation disclosed a frankly purulent mastoid.

CASE VIII

A girl of 6 had had pain in the left ear for some days. There was a slight temperature and some discharge from this ear. The meatus was narrowed, and there was some excoriation of its external opening. There was no tenderness. There was a history of acute earache some years previously, followed by paracentesis, but whether of one or both ears was not known. There had, however, been pain each winter, sometimes in one ear, sometimes in the other. There were no physical signs referable to the right ear at the time of examination. Both mastoids contained pus, that in the left being under pressure.

CASE IX

A girl of 8, sister of the previous case and seen a few weeks later. She had had pain in the left ear for three days. There was no temperature, no tenderness, and she was cutting a tooth on that side. The tonsils and adenoids had been removed five years previously. Two years ago there had been earache, both sides (?). There was no history of any discharge. In the left ear there was an accumulation of epithelial debris and, in the antero-inferior quadrant, a puckered area and some granulation tissue (?) suggesting a partially closed perforation. A similar area, but without granulations, was observed in the same position on the right drum, which was thickened. Shrapnell's membrane on the right side was red, but not bulging, and there was no tenderness; there was slight deafness. On the following day there was slight tenderness over the tip of the left mastoid, none over the right. Temperature still normal. Both mastoids were filled with granulation tissue and pus.

In both these children (Cases VIII and IX) the ears recovered completely, and even a long spell of whooping-cough the same winter failed to produce aural symptoms.

CASE X

A boy of 12. A recent attack of acute tonsillitis had resulted in an exacerbation of a bilateral ear discharge, which had been intermittent since acute earache following measles two years previously. The temperature was raised, and there was definite tenderness over the right mastoid. There were no physical signs referable to the left ear other than the discharge. At the operation there was nothing to choose between the two mastoids, both being frankly purulent.

CASE XI

A hospital nurse was warded for an attack of (?) influenza. There was a slight initial rise of temperature, which quickly subsided. Ten days later, while still in bed, she had an acute otitis media on the right side. Rupture of the drum resulted in the temperature again becoming normal, and it remained so. Nevertheless, three weeks later, the discharge, which in the meantime had steadily diminished in quantity, could still be seen pulsating at the perforation. There were also vague pains referred to that side of the head. There was no tenderness, and x rays failed to show any difference between the two mastoids. At the operation there was a considerable quantity of pus under pressure, and a very forwardly placed lateral sinus was already exposed to the sepsis. The organism was *Streptococcus haemolyticus*.

These cases are not exceptional. On the contrary, they represent my general experience. They must represent the experience of many others also. Whether or no latent mastoid sepsis is really as often the cause of chronic aural sepsis as these cases would suggest should be speedily settled once we are alive to its possibility. In all the cases in which I have performed the cortical mastoid operation solely on account of the persistent discharge there have been only two in which the mastoid was not frankly purulent. It cannot, however, be supposed that every case of persistent discharge following an acute otitis media is due to the presence of mastoid sepsis. At present every such case must be considered on its merits. I would suggest that in doing so the probable frequency of unsuspected mastoid sepsis should be borne in mind.

THE TREATMENT OF CHRONIC AURAL SUPPURATION WITH IODINE POWDER

BY

R. SCOTT STEVENSON, M.D., F.R.C.S.Ed.

SURGEON TO THE METROPOLITAN EAR, NOSE, AND THROAT HOSPITAL

The purpose of this short article is to emphasize the value of a simple method of treating chronic middle ear suppuration by insufflation of so-called iodine powder, which consists of iodine 0.75 per cent. in boric acid powder. Although I began to use this powder tentatively and without much conviction of its value two years ago, I now use it as a routine for every case of chronic middle ear suppuration, first mopping or sucking out the discharge and then blowing the powder through the perforation into the middle ear and until the deepest part of the meatus is full. Nascent iodine apparently gets into the furthest corners of the affected middle ear. A simple powder blower with a glass or wooden stem and a rubber bulb is quite adequate for the patient's own use, but a special all-glass or vulcanite powder insufflator should be used by the practitioner.

I first came across this method of treatment in Chicago and New York in the autumn of 1930. The iodine powder used was an American proprietary preparation, originated by Nathan Sulzberger, a New York chemist, about 1915, and was called by him "iodosan." It was used first of all for suppurative conditions of the nasal cavities, and only afterwards for chronic middle ear suppuration. The percentage of iodine in the powder varied from 10 to 0.75, but it was found more effective and much less irritating in the smaller percentages; in America it is now used commonly as 0.75 per cent. for the initial treatments and as 2 per cent. after a number of treatments have been given.

Many practitioners are under the impression that the treatment of ear discharge by powder insufflations is dangerous, as hard concretions may form from the powder and the drainage become obstructed, especially with small