had always stressed in early editions of *Tropical Diseases* that the life-history would in the main be analogous to that of the liver-fluke, *Fasciola hepatica*, whose cycle in a pond snail—*L. truncatula*—had first been worked at by Thomas in 1883. The first inkling of the true life-history was afforded by Ogata in 1914 for the allied species *Bilharzia japonica*, which was afterwards confirmed by R. T. Leiper and E. L. Atkinson. In 1915 R. T. Leiper, in Egypt, not only discovered the life-history of this parasite in the freshwater snail *Bullinus contortus*, but was also able to separate the intestinal form of bilharzia—*B. mansoni*, of which the lateral-spined eggs occurring in the faeces had been recognized in 1903 by Manson, and which was found to possess a distinct morphology and to develop independently in a snail belonging to a different genus, *Planorbis*. The method of successful treatment by means of continued injections of antimony tartrate in increasing strength we owe to J. B. Christopherson, who worked it out in Khartum in 1917, and it has hardly been improved upon up to the present day.

COUGH IN CHILDHOOD *

BY

ALAN MONCRIEFF, M.D., M.R.C.P.

PHYSICIAN FOR DISEASES OF CHILDREN, BOLINGBROKE HOSPITAL,
LONDON

The symptomatology of diseases in childhood is rendered difficult by the absence of accurate self-expression on the part of the patient; in consequence, any symptom which can be carefully described by other observers must be regarded as of very great value. It is probable that in this fact lies the reason for "cough" being one of the commonest complaints made by parents on behalf of their children, and the frequency of this symptom justifies a more detailed analysis than is usually made in the standard textbooks. It will probably serve to introduce the subject in a more orderly manner if a definition of "cough" is first attempted, followed by a brief summary of the modern views on the physiology of the condition.

A cough is a modified expiration of an explosive character designed essentially to remove a foreign body from the mucous membrane of the respiratory tract.¹ Coughing may be voluntary or it may be purely an involuntary reflex act. Usually it is a mixture of the two: a reflex act reinforced by volition.² Thus, even in the voluntary act, there is a reflex element, and it is therefore important to consider the possible sites where the primary stimulus may arise. Broadly speaking, the afferent portion of the reflex arc runs in the glossopharyngeal nerve from the upper parts of the respiratory tract and in the vagus from the lower parts. A great deal of work has been carried out in recent years on the relative sensitivity of the various zones, and it seems conclusive that there are definite areas in the pharynx and in the region of the larynx which are more sensitive than others. The region of the bifurcation of the trachea is also much more sensitive than the rest of this structure, and as the respiratory mucous membrane is traced further and further down the bronchi sensitivity gradually disappears. The lung parenchyma is probably almost completely insensitive, although the evidence is a little contradictory. Perhaps the difficulties here can be explained, as Chevalier Jackson has suggested,² on the grounds that "tolerance" is very quickly established in the bronchial system, and after a foreign body has been inhaled the initial stage of coughing or choking is followed by a silent period as

long as the foreign body remains fixed. If movement occurs a new area of mucous membrane is involved, and this sets up a reflex cough, followed again by a period of silence. After pus forms, the cough reappears because fresh areas are stimulated by the inflammatory process. Here again there are difficulties, but, according to the same authority, in any inflammatory condition of the lungs or bronchi the sensitivity of the mucous membrane seems to be diminished. With diminished sensitivity of the smaller bronchi and absolute insensitivity of the lung parenchyma, it is not surprising that in lobar pneumonia, for example, very little cough may be present; this disease, especially in childhood, may pass almost unnoticed, so little is there to direct attention to the lungs. Pleural involvement leads to the classical restrained type of cough, since the pleura is another area from which the cough reflex may be initiated. It should also be noted that, experimentally, there appears to be a strong reflex connexion between the nasal mucous membrane and the bronchial motor system, so that cough may be initiated by the effect of irritant gases in the nose. Similarly, cough may be produced experimentally by certain forms of stimulus applied to the oesophagus or stomach, probably because of the common vagal nerve supply.

It follows from this brief summary of the origin of the cough reflex that the upper part of the respiratory tract down to the level of the bifurcation of the trachea is of very much more importance than the portions of the respiratory apparatus below this point, and that, especially in childhood, coughs have much less to do with the "chest" than is commonly imagined. It is not an exaggeration to say that the vast majority of coughs in childhood are characterized by a complete absence of physical signs on examination of the chest.

CAUSATION

In considering the possible causes of cough in childhood it will be convenient to proceed anatomically from the upper regions of the respiratory tract downwards, adding comments on special features as the long list of possibilities is set out.

Inflammation of the Nose and Nasopharynx.—Acute inflammatory conditions of the nose and nasopharynx are frequently associated with a cough, but in such instances it is almost impossible to exclude inflammation of other portions of the respiratory tract, and the cough may have a complex origin. Similarly with the difficult question of adenoid tissue in the nasopharynx, this is so frequently associated with tonsillar hypertrophy or pharyngitis that it is impossible to be certain whether adenoids alone can cause a cough. In babies with the comparatively rare condition of congenital adenoid enlargement, a cough may occur, and this, I believe, is an example of a pure "adenoid cough." A foreign body in the nostrils is said to cause a cough in some instances, but I have not yet seen an example of this. With chronic inflammation of the nose and nasopharynx, with or without "adenoids," it is quite common to see pus trickling down the posterior pharyngeal wall; cough in such cases may in part be due to the irritation of this inflammatory material in the lower parts of the pharynx. To sum up the evidence here, I should say that to attribute a cough to a purely nasal or nasopharyngeal origin is usually unwise. The possibility, however, of a cough due to an ear disease must be mentioned. Here again the exact aetiology is complicated by the concomitant tonsillitis or pharyngitis, and I have not yet seen a child with a cough which could be directly attributed to middle-ear disease alone.

Coughs of Pharyngeal Origin.—These are very common, and need little discussion. It should be remembered that a young child seldom complains of a sore throat, and

* An address to the South-West London Medical Society.

acute inflammatory conditions of the tonsils, including diphtheria, may have a cough as the dominating symptom. This is a characteristic barking cough which is easily recognizable, and it is just this type of cough, due to recurrent acute tonsillitis, which is usually labelled "bronchitis" by the mother. Even a cursory examination of the chest will exclude this diagnosis, and a view of the throat soon settles the matter. It is important to make this diagnosis accurately and early, not only because a diphtheritic membrane may be missed, but also because the stock "bronchitis" mixture with ipecacuanha wine is useless for treating inflamed tonsils.

Larynx.—We come next to the larynx, and here the characteristic "croupy" type of cough usually makes diagnosis easy. Most laryngeal affections are a combination of inflammation and spasm, varying from almost pure laryngitis through laryngitis stridulosa, where the spasm is playing a part, to laryngismus stridulus, where underlying tetany produces an almost purely spasmodic condition. There are three conditions of the larynx and neighbouring structures in which difficulty may arise: (a) a croupy cough in measles before the rash appears, due probably to the occurrence of Koplik's spots on the laryngeal mucosa; (b) retropharyngeal abscess, where some oedema of the larynx is not uncommon and a croupy type of cough occurs; and (c) laryngeal diphtheria, which must always be suspected until it can be excluded either on clinical or bacteriological grounds. The differential diagnosis of all the various laryngeal affections would take us too far from the main purpose of this paper. It is comparatively easy, however, to localize the origin of the cough as laryngeal where the curious husky bark is present. Retropharyngeal abscess is readily excluded by an exploring finger.

Trachea.—Coughs due to affections of the trachea form an important group. The harsh, painful, ineffective cough of a tracheitis is a comparatively common sequel to a "cold in the head," and in adults the retrosternal discomfort makes the diagnosis easy. In the young child no help can be obtained in this way, but sometimes it is possible to hear an occasional coarse rhonchus over the upper part of the sternum. Pressure on the trachea may cause a cough, and there are three main conditions in childhood in which such pressure may occur. In babyhood an enlarged thymus sometimes accounts for a loud stridor and a loud brassy type of cough. During the last few years I have seen several with marked symptoms and good x-ray photographs of thymic enlargement; under radiation treatment (radium or x rays) the symptoms and shadow simultaneously disappeared. In later childhood enlargement of the mediastinal glands may give rise to a harsh, hacking, and spasmodic cough, which is sometimes very puzzling. I have discussed elsewhere³ the diagnosis of the condition, which is undoubtedly not easy. The proper eliciting of D'Espine's sign is, I think, of most value; bronchial breathing and whispering pectoriloquy heard below the level of the fourth and fifth thoracic spinous processes in a child is very suspicious, and x rays can be used in confirmation, especially with the oblique view to show the posterior mediastinum. A third cause of tracheal pressure is advanced heart disease, with or without pericarditis, and in many instances the cough is, I believe, due to enlarged lymphatic glands in the anterior mediastinum. The diagnosis is usually easy.

Bronchitis.—Inflammation of the bronchi frequently follows tracheitis as the "cold spreads down to the chest," and the physical signs of true bronchitis require no particular comment. The rule is, roughly speaking, rhonchi on the large tubes and rales in the smaller tubes, and the finer the rales the more anxious one becomes lest bronchopneumonia should develop. Notice should be taken whether the added sounds are mostly inspiratory or

expiratory in type: if the latter, then some bronchial spasm is probably present, and treatment must be directed against this. Other affections of the bronchi giving rise to cough need little comment. Bronchiectasis is characterized by showers of rales at one or other base (usually the left) after coughing has emptied some of the secretion out of the dilated bronchi and allowed air to enter. Asthma is easily recognizable during the attack, and between attacks there may either be no physical signs or those of bronchitis. It should be remembered that the younger the subject the more likely is recurrent bronchitis to be of asthmatic origin, and inquiry should always be made for any history of nocturnal wheeziness.

Inflammatory Conditions of the Lung.—Coming to the lung parenchyma, I have already stressed the fact that pneumonia may be a "silent" disease in the child as far as cough is concerned. This is true only of the primary lobar type, the common type in childhood, for in the secondary catarrhal or bronchopneumonic form there has usually been a previous bronchitis—and the alveolar involvement is secondary to this. Chronic inflammatory conditions—non-tuberculous fibrosis of the lungs—are common causes of cough in childhood, especially where the cough follows measles or whooping-cough. The physical signs of this condition are varied, but I pay particular attention to diminished movement and air entry at the base of one or other lung, especially if this is associated with persistent catarrhal signs. X-ray examination will confirm the diagnosis in most instances. Tuberculosis as a cause of cough in childhood is comparatively rare. Acute miliary forms give rise to little or no cough, while of the chronic type, tuberculous mediastinal glands are very difficult to diagnose, and cough may or may not be present. Fibrocaseous tuberculosis of the adult type does occur sometimes in childhood, but cough is never the same dominant symptom as it is in adult patients. It is usually easy to exclude this type of the disease on physical examination, followed by an x-ray picture in cases of doubt.

Pleurisy, serous or purulent, as a cause of cough in childhood presents no essential difficulties. A restrained irritating cough will occur during the early "dry" phase and a period of silence usually follows. For this reason it is a common occurrence for empyema in children to be missed and to assume a chronic form; the condition then presents itself for loss of appetite or anaemia!

From the nose to the pleura completes the anatomical survey, and I fear the list has been a long one. There still remain one or two coughs to be discussed. First let us consider "pertussis," for, I suppose, the question which bothers us all most in the coughing child is to decide whether or not this is whooping-cough. Once the whoop has occurred diagnosis should, of course, be easy, but mothers are extraordinarily reluctant to admit the existence of a whoop, even when you and she have listened to a perfect exhibition of a typical paroxysm together. Before the whoop occurs suspicion must be attached to a nocturnal cough which is spasmodic and associated with vomiting. This is especially likely to be confirmed if no physical signs can be found in the chest and no disease of the upper respiratory tract. The sublingual ulcer is always worth looking for, and sometimes gives help in a difficult or important case. In one patient seen recently the ulcer was on the side of the tongue instead of beneath it, and this was due to the position of the teeth, the lower central incisors being out. Bacteriological methods of diagnosis are probably quite useful, especially the procedure recently described by Gardner and Leslie,⁴ but quite inapplicable to the conditions of most out-patient departments or of many practices. If in doubt the case should be regarded as whooping-cough; a week or ten days generally settles the matter.

Then there is the "stomach" cough: on the authority of Hess there are physiological experiments to prove its possibility, and coughs have been produced artificially by appropriate stimulation to the oesophagus or stomach. I think the most usual explanation is that the attack of gastritis, or whatever it is which is the stomachic element of the cough, leads to a general catarrhal state of the mouth and pharynx, also seen in the coating of the tongue, and this causes the cough. I never diagnose the condition unless I see the coating of the tongue extended right over the root of this organ and a red, glazed condition of the posterior pharyngeal wall. On the whole, it is a diagnosis to make with great caution, as it is a difficult one to live down with parents, and is apt to be sneered at.

Lastly, there is the question of cough as a habit spasm. Following an attack of whooping-cough a child is liable to go on coughing and whooping for months, for no more obvious reason than another child will go on twitching its mouth long after a patch of impetigo has vanished from the lip. To diagnose a cough as a habit spasm, however, necessitates a very careful examination of all other possible causes, and, as with any diagnosis resting on exclusion, it has inherent weaknesses. In this category also is the so-called barking cough of puberty, a cough which is labelled as hysterical in some of the textbooks. Here again diagnosis is largely by exclusion, and to attribute a cough to a purely psychical origin requires a very close examination of the whole environment of the child. As in *Alice*, it may be that

"He only does it to annoy
Because he knows it teases."*

TREATMENT

If the origin of the cough has been accurately localized the question of treatment becomes relatively simple. Obviously, since the prime object of a cough is to remove a foreign body from somewhere in the respiratory tract, the cough should be encouraged when there is the possibility of achieving this, and discouraged otherwise, as in most of the instances I have outlined. In other words, measures should be taken to stop almost all coughs originating above the bifurcation of the trachea: this also holds true for many coughs below this point. It follows that expectorants have a very limited use, and in my opinion too much attention has been directed in the past, as far as children's medicine is concerned, to the question of "depressant" and "stimulating" varieties. For most coughs in children a sedative linctus is best, such as tinct. camph. co. mx, oxymel scillae mx, glycerinum mx, spir. chlorof. mij, aq. ad. ʒi, for a child of 2 years, as required.

Obviously, the first step in the treatment of cough is to eliminate or treat the cause as far as possible. As this will mean measures varying from radiation of an enlarged thymus to evacuation of an empyema it is not possible to discuss each type of cough in detail in this respect. I would like to emphasize, as I have already done elsewhere,⁵ the importance of irritant gases as a cause of inflammation of the respiratory passages. In London the acidity of the atmosphere is sufficiently powerful to make serious inroads upon the stonework of the Houses of Parliament, and it is not surprising that many infants cough from their earliest days. Still⁶ has also called attention to the possible irritation produced by a smoking fire in the nursery. Adequate treatment of coughs in childhood requires, in certain instances, a study of home surroundings.

The steam-tent in the treatment of respiratory disorders of childhood has a very limited field of usefulness and

many drawbacks. In some animal experiments I carried out last year* it was found that hot moist air quickly caused fatal oedema of the lungs if the temperature was allowed to approach that of the body. The most beneficial effect in the relief of artificially produced spasmodic conditions of the bronchi was obtained when the air was merely kept fully saturated with water vapour at room temperature. This is best achieved by a kettle on the hearth, well away from the child's bed; no steam should be visible in the room at all. Used in this way moisture is of assistance in the relief of spasm, whether this be in the larynx or in the bronchi. It should be remembered in this respect that bronchitis with spasm is frequently a manifestation of asthma in the young child, and remedies based upon modern work should be employed—namely, correction of constitutional errors in digestion, removal of all possible sources of animal emanations and dust, and drugs to stop the acute attack.

Lastly, the treatment of bronchitis requires a brief word. Many of the drugs used, such as ammonium carbonate, squill, and ipecacuanha, have a direct irritant effect upon the alimentary tract if used in anything but small doses, and in young children the use of these drugs may lead to a troublesome and often dangerous gastro-enteritis. When the secretion is thick, during the early stages, alkalis are most valuable, and a good combination is vin. ipecac., mij, pot. cit. gr. v, tinct. camph. co. mv, glycerinum mx, aq. ad. ʒj, three times a day, for a child of 1 year. Later on a little ammonium carbonate (gr. 1/2) may replace the potassium citrate, and if much wheeziness is present a couple of minims of tincture of stramonium should be added. The use of atropine to "dry up the secretion" in such cases is not recommended; it has, indeed, been condemned by Chevalier Jackson. It is doubtful whether much can be done to aid nature in the mechanism of expectoration, but harm can certainly be done by ill-considered interference.

REFERENCES

- ¹ Hess, W. R.: *Die Regulierung des Atmung*, Leipzig, 1931, p. 91.
- ² Jackson, C.: *Journ. Amer. Med. Assoc.*, 1922, lxxix, 1399.
- ³ Moncrieff, A.: *Post-Grad. Med. Journ.*, 1930, vi, 58.
- ⁴ Gardner, A. D., and Leslie, P. H.: *Lancet*, 1932, i, 9.
- ⁵ Moncrieff, A.: *Journ. State Med.*, 1932, xl, 336.
- ⁶ Still, G. F.: *Common Disorders of Childhood*, London, 1927, fifth edition, p. 434.

AETIOLOGY AND SYMPTOMS OF MITRAL STENOSIS

A REVIEW OF THREE HUNDRED CASES

BY

E. WYN JONES, M.D., D.P.H.

SENIOR MEDICAL REGISTRAR AND TUTOR, LIVERPOOL ROYAL INFIRMARY

It has been stated by Coombs¹ that every case of mitral stenosis is a case of advanced cardiac rheumatism, and this writer has produced clinical and pathological evidence in support of the statement. It does not, however, appear clear in his monograph in what percentage of cases of mitral stenosis a rheumatic history was obtainable. Price² states that mitral stenosis is *usually* due to rheumatism in one or other of its protean manifestations.

The following analysis of 300 cases of mitral stenosis was undertaken on clinical lines in order to assess the importance of the history of rheumatic fever and allied conditions in the causation and aggravation of the disease. During the investigation a number of other factors have been noted. The age and sex incidence of mitral stenosis are discussed briefly, and the symptomatology is also considered.

* Before accepting this modern psychological dogma too hastily the chorus following these immortal lines should be recalled—"Wow, wow, wow!"

* Working with a part-time personal grant from the Medical Research Council.