question, as behind most of those I have touched on, and many others such as the relation between industrial medicine and the other services, lies the deeper question of the aims of national policy.

Nowhere in the Statute Book can I find any definition of aims that would give us a unifying line through our problems of organization. The truth is, as we all know, that the health services have grown up haphazard. From crude beginnings at control of infection they now embrace a range of scattered activities that include sanitation, infectious diseases services, with special branches for tuberculosis and venereal diseases, school medical service, maternity and child welfare, industrial medical inspection, national health insurance, and several others. Each development of a service or addition to the number of services was made in response to some special need or feeling of the times. The result is a congeries of disparate activities. The insurance practitioner works in his field, the child welfare officer in his, the school medical officer in his, and none is inspired by the feeling that he is linked to the others by a common purpose.

It is my own belief, and the belief of many others, that we cannot and should not expect to solve our problems of defective results or organization until we have arrived at a definition of policy that would unify, so far as is possible, the whole range of activities bearing on the health of the people. That is not a job solely or even primarily for medical men; but the medical man must make his contribution along with the biologist, sociologist, administrator, and the plain citizen. The need for a clear definition of policy has been reinforced by the

development of biological science in recent times. I do not mean by biology the crude propaganda that masquerades as eugenics, although the criticism that in some respects we are striving officiously to keep alive the weak at the expense of the strong, and thus doing injury to biological principles and to the race in the long run, has to be met. But biology, the science of life, and that branch of it, social biology, which deals with the numbers and quality of population, has much to say to us that could not be said when many of the present services and much of our existing organization were established. It will be strange if, in the light of modern biological knowledge, we do not find it necessary to adapt our services and organization so as more directly to promote healthy living. I do not imply that philanthropy should be shut out from the health or other social services; far from it. Philanthropy is a good old word, much abused, and the spirit it expresses, I hope, will live on in all our activities. What I imply in my plea for a biological assessment is a clearing of our minds so that we shall know when we are serving a biological and when a philanthropic purpose, and we shall regulate our activities accordingly.

The present is a time for stocktaking in medicine and in health policy. We have to discover our aims; we have to assess the worth of our present activities and modify them where necessary, so that they may most fittingly achieve the ends of defined policy. From such a stocktaking there will come, I believe, not merely improved organization, but also a new inspiration, which will carry us to achievements in public health greater than any we have known.

THE PRESENT POSITION OF SEROLOGY IN RELATION TO THE PATHOGENIC COCCI*

BY

H. J. PARISH, M.D., M.R.C.P.ED.

WELLCOME PHYSIOLOGICAL RESEARCH LABORATORIES, BECKENHAM

During the past decade there has been a rapid expansion of our knowledge of the pathogenic properties of the cocci which cause disease in man—more especially of the Gram-positive group. The chief advance has come from the realization that staphylococci and streptococci are true toxin-producing organisms in the same sense that the diphtheria bacillus and tetanus bacillus are toxin producers. Although no true toxin has been discovered for the pneumococcus, improved methods of production of antibacterial serums have brought a new hope in the treatment of pneumonia. I propose to discuss the bearing which these and other advances have had upon the serology of the pathogenic cocci in relation to the recognition, prevention, and treatment of the human diseases for which they are responsible.

STAPHYLOCOCCUS

It has frequently happened in the history of mankind that a calamity has been turned to good account through the stimulus it has provided for more thorough investigation and research. While the existence of staphylococcal toxins had been fully realized and many of their properties had been described (Neisser and Wechsberg,¹ 1901; Kraus and Pribram,² 1906; Nicolle and Cesari,³ 1914; Parker,⁴ 1924), a disaster at Bundaberg, Queens-

land, in 1928, led to a more intensive study of staphylococcal toxin and antitoxin by Burnet^{5, 6, 7} (1929, 1930, and 1931). In this disaster twelve of twenty-one children, who had been inoculated with contaminated diphtheria toxin-antitoxin, died from acute staphylococcal septicaemia. Subsequent to the work of Burnet in Australia, the investigations of Weld and Gunther⁸ (1931) in America, Parish and Clark^{9, 10} (1931), Panton, Valentine, and Dix¹¹ (1931), and Panton and Valentine¹² (1932) in England, amongst others, served to make the properties of staphylococcal toxins and antitoxins more apparent.

Toxic Products

Toxins prepared by a special method kill rabbits, guinea-pigs, and mice rapidly, produce skin lesions in guinea-pigs and rabbits (necrotoxins or dermatotoxins), and have a destructive action on leucocytes (leucocidins) and erythrocytes (haemotoxins or haemolysins). An enterotoxic substance, which caused vomiting and diarrhoea in monkeys, has also been described by Woolpert and Dack¹³ (1933), who have made a special study of strains of staphylococcus from cases of food poisoning. There was a rough correlation between the amount of this food poison and the amounts of killing toxin, dermatotoxin, and haemolysin. Although Burnet formerly regarded the various toxic constituents as manifestations of a single toxin, the work of Weld and Gunther⁸ (1931) and Woolpert and Dack¹³ (1933) suggests plurality of toxins, and Glenny (unpublished) has found at least two haemolysins with specific antibodies. In human disease the various toxic properties, which have been demonstrated in animals and *in vitro*, undoubtedly play their part; the lethal effects were only too clearly demonstrated at Bundaberg. There is no skin test for susceptibility or immunity to staphylococcal infections in human subjects analogous to the Schick and Dick tests. The reactions

* Read in opening a discussion in the Section of Pathology at the Annual Meeting of the British Medical Association, Dublin, 1933.

which follow the intradermal injection of toxic filtrates cannot be correlated with the amount of circulating antitoxin (Bryce and Burnet,¹⁴ 1932).

Toxoid

Staphylococcal toxins may be rapidly converted into toxoid by formaldehyde. Toxoid is innocuous to laboratory animals, but can still bind antitoxin *in vitro*, and is an efficient immunizing agent, producing an increase in the amount of circulating antibody. Encouraging results have been reported from its use in the treatment of persistent, intractable, or recurrent staphylococcal diseases (Panton and Valentine,¹² 1932, Dolman,¹⁵ 1933). We have provided some clinical friends with toxoid for use in cases of the kind under controlled conditions in an endeavour to confirm these observations.

Antitoxin

Antitoxin which is also an antihæmolyisin can be prepared by the inoculation of horses and rabbits with a culture filtrate, but not with washed whole staphylococci. The serum can be concentrated. Apart from its capacity to neutralize toxin *in vivo* and *in vitro*, its most interesting property is that when it is given intravenously to rabbits with, or a few hours after, a certainly fatal dose of virulent living culture it prolongs and in some cases saves life. It is perhaps not generally realized that the use of antitoxic serum in staphylococcal infections is a recent development, for the older anti-staphylococcal serums were produced in response to immunization with suspensions of staphylococci and not toxic culture filtrates. The excellent results which have been obtained with antitoxin in laboratory experiment, and the promise which has attended its use in a small number of patients, justify more extensive clinical trial, especially in staphylococcal septicaemia. It is likely to be most effective in cases in which toxæmia is predominant, and should be given early, in large doses, and by the intravenous route. It is a rational form of therapy, for, theoretically at least, a serum which neutralizes toxins impedes one of the more important modes of attack by the staphylococcus, and thereby aids the natural defences of the tissues. On the other hand, one has to admit that contradictory reports have been made as to its usefulness. Its effect on the invasive properties of the coccus has not yet been determined, and it cannot be expected to take the place of surgery in such conditions as osteomyelitis, and staphylococcal abscess of the kidney, although it may be a useful adjunct. I have already referred to the plurality of toxic constituents in staphylococcal filtrates; antibodies to these substances are developed in different amounts in the serum of horses undergoing immunization. Further clinical investigation is necessary in order to assess the importance of these differences in antitoxic serums.

STREPTOCOCCUS

Hæmolytic Streptococcus Exotoxin and its Specificity

As in the case of the staphylococcus, the early literature of the hæmolytic streptococcus contains many references to hæmolyisin, leucocidin, and the toxic symptoms produced after the injection of culture filtrates into animals. A new era of research followed the discovery by the Dicks¹⁶ (1924) of the rash-producing (erythrogenic) property of the exotoxin of a scarlet fever strain. Filtrates were subsequently prepared not only from streptococci isolated from scarlet fever, but also from septic sore throat, erysipelas, puerperal fever, and other infections. There is still some difference of opinion as to the inter-relationship of these toxins, but it is coming to be recog-

nized that in most cases there is, if not identity, at least a common valency. "A clear-cut disease specificity of hæmolytic streptococcus exotoxin quite definitely does not exist" (Williams,¹⁷ 1932). Parish and Okell¹⁸ (1928) and Okell¹⁹ (1932) have stressed the essential similarity of toxins of the majority of strains from different sources. These workers found that scarlet fever antitoxin, prepared by the immunization of horses with filtrates of the Dochez strain of *S. scarlatinae*, could counteract toxæmia in rabbits infected not only with cultures of hæmolytic streptococci from scarlet fever, but also from erysipelas and puerperal septicaemia. Homologous antitoxins, prepared with the less toxic filtrates of erysipelas and puerperal strains, were also effective, but to a less extent than the scarlet fever antitoxin. McLachlan²⁰ (1927), J. Smith²¹ (1927), and other workers have performed cross-neutralization tests in Dick-positive subjects, and also concluded in favour of the very close relationship of the toxins used. Wadsworth^{22, 23} (1932 and 1933) has, however, noted that marked qualitative variations may be encountered in the toxins, even of two closely related strains, but that with selected cultures—for example, the Dochez N.Y.5—a serum could be prepared which neutralized the toxin of a large number of strains. Wadsworth agrees with Parish and Okell that there is no known method of distinguishing between the so-called *S. scarlatinae* and other human hæmolytic streptococci, such as *S. erysipelatis*, and that "phasic variations" on the part of the streptococcus developing under special conditions of susceptibility on the part of the host would explain the apparently "specific" clinical conditions resulting from infection. He distributes for the treatment of all streptococcal infections a monovalent serum prepared by the immunization of horses with a scarlet fever strain of streptococcus—the Dochez N.Y.5—a procedure which has also been recommended by the English workers.

Antitoxin Treatment of Scarlet Fever

There would appear to be little doubt in this country and in America as to the value of scarlatinal antitoxin in the treatment of the initial toxæmia of scarlet fever, but there is very conflicting evidence as to its usefulness in preventing the septic complications. In the laboratory we found that while we were able with antitoxic serum to protect rabbits from the acute phases of the septicaemia resulting from the intravenous injection of culture, the majority of rabbits developed arthritis towards the end of the first week or later. Protection was never obtained against this phase of the infection with any antitoxic or so-called antibacterial serum we used. It is perhaps not surprising, therefore, that the incidence of septic complications has not been greatly influenced in hospital wards. Banks^{24, 25} (1932 and 1933) has had excellent results following the intravenous use of antitoxin; septic complications were almost wholly prevented, and the period of morbidity was reduced to just over a fortnight. On the Continent, however, where more serious outbreaks have occurred, the newer therapy has not established itself as an unqualified success, even in combating the initial toxæmia. It is possible that difficulties in connexion with titration, and thus with ensuring supplies of potent serum, have here played a part. Veldee, Stevenson, and Mitchell²⁶ (1931) have also concluded that "there is not an irrefutable case for the use of streptococcus antitoxin in the treatment of scarlet fever," but it should be noted that the majority of their cases were mild, and the dosage for the more severe cases would be considered inadequate by most clinicians. From a general review of the clinical and laboratory evidence I have little doubt that the treatment of scarlet fever with antitoxin is a rational procedure and has come to stay.

Passive Immunity and Active Immunization

McClean²⁷ (1927), Joe and Swyer²⁸ (1929), Nabarro and Signy²⁹ (1931), and many others have injected antitoxin prophylactically in doses of 5 to 10 c.cm. into all the Dick-positive contacts of cases of scarlet fever in hospital wards. Passive immunity is conferred which lasts for about a fortnight. At Great Ormond Street Hospital the method has been highly successful; nearly all cleft-palate and hare-lip patients are Dick-tested, and, if positive, given antitoxin before operation; scarlet fever rashes do not occur as in the pre-antitoxin era. Operations on the mouth region are carried out almost with impunity in any ward where scarlet fever is present. While serum has been used prophylactically in many other institutions in England with highly satisfactory results, it is, rather surprisingly, not employed to the same extent in the United States of America.

Dick-positive persons can be actively immunized with toxin, and become in this way negative (Dick and Dick,³⁰ 1924). Williams¹⁷ (1932) recommends five subcutaneous injections of 500, 2,000, 10,000, 20,000, and 40,000 S.T.D. (skin test doses), and retests after one to six months. The majority of fever hospital superintendents are emphatic that active immunization is worth while, and state that nurses naturally Dick-negative do not contract scarlet fever.

Serum Treatment of Streptococcal Infections other than Scarlet Fever

The efficacy of haemolytic streptococcal antitoxin in combating non-scarlatinal infections has also been the subject of inquiry, and, although a beneficial effect has sometimes been reported, there have been many failures, and some years must elapse before a final judgement is possible. Wadsworth²³ (1933) has had striking improvement in erysipelas, the effect on the toxæmia preceding by a short interval in most cases the improvement in the local lesions; in other cases, however, serum therapy was without avail.

(Incidentally, in rabbit experiments, Parish and Okell¹⁸ (1928) could find no evidence of specific blocking with antitoxin of the skin reactions produced by haemolytic streptococci injected intradermally, although, as stated above, they were successful in controlling the toxæmia following the intravenous injection of haemolytic streptococci from various sources. Their results regarding the influence of antitoxin on skin lesions were thus at variance with those of Birkhaug³¹ (1925).)

In miscellaneous streptococcal infections, with and without bacteraemia, Wadsworth has had good results, provided the appropriate surgical measures were also employed whenever indicated, but his most encouraging cases were those of septic sore throat, in which the response to serum therapy was "even more striking and spectacular than in scarlet fever or erysipelas." The last claim is surprising, for one would hardly expect passive antitoxic immunity to have much influence on the local faucial manifestations of streptococcal disease. Wadsworth insists that in all streptococcal diseases early administration and adequate dosage are all-important if success is to be achieved.

Another recent paper to which I must refer is that by Benson and Rankin³² (1933), who deprecate the use in *puerperal septicaemia* of large doses of expensive serum, which have not only been ineffective, but have caused unpleasant symptoms, especially after intravenous injection. They make an exception for its use in the rare cases of puerperal scarlet fever. Since the vast majority of haemolytic streptococci from whatever source are capable of producing erythrogenic toxin *in vitro*, the use of antitoxin serum is theoretically sound treatment, even in those cases in which rash is absent. One might expect the antitoxin in

certain instances to "tip the beam" in favour of recovery. However, some clinicians of repute, though not all (see Wadsworth,²³ 1933), have come to the same conclusion as Benson and Rankin, as to the inadequacy of streptococcal antitoxin in puerperal cases. Apparently the extent and severity of the local condition, in particular the presence or absence of dying or necrosed tissue, are much more important factors in determining the prognosis than is the amount of erythrogenic toxin passing into the circulation. In this connexion it is of interest to note that there is no significant difference between the liability of Dick-positive and Dick-negative pregnant women to invasion with the haemolytic streptococcus (Joe,³³ 1929). On reviewing the situation I feel I cannot lightly dismiss the presumptive evidence that scarlet fever antitoxin may be a useful therapeutic measure in the early high fever of puerperal streptococcal infections.

Although an antibacterial action on a homologous strain has been attributed to serums prepared experimentally by Lancefield and Todd³⁴ (1928), Downie³⁵ (1930), and others, no *antibacterial serum* is at present available to combat the pyogenicity and invasive properties of the streptococcus. Recently hope was raised by the preparation of an antihaemolysin of high titre by Todd^{36 37 38 39} (1932 and 1933). Unfortunately "haemolysin" cannot be identified with the pyogenic principle.

Titration of Antitoxin

Titration of antitoxin is still a matter of considerable difficulty, no method having yet met with international acceptance. The skin neutralization test on human subjects—determining the quantity of antitoxin which will neutralize a known amount of toxin on intradermal injection—is much used in America. Veldee¹⁰ (1932) has recently devised a similar skin neutralization method on the inner surface of the rabbit's ear. This method is still in the experimental stage, but has not proved very satisfactory in our hands. It is apparent from recent publications (Wadsworth,²³ 1933) that there is difference of opinion in America as to the minimum standard of potency for issue, and that many serums have been issued in the past of relatively low value. In our own laboratory the Parish-Okell⁴¹ rabbit method—finding the quantity of antitoxin which will neutralize a culture of the Dochez streptococcus, or a dose of toxin given intravenously—has been used for many years, and a review of the evidence gives reason for believing that the rabbit method has ensured the production in England of serums which have been satisfactory clinically; tested by this method they have corresponded in potency to the best serum made abroad. My colleagues, Dr. Buttle, Dr. O'Meara, and Miss Whatley, have recently been engaged in investigating other and less expensive methods of titration. Dr. Buttle has been able to evaluate serums on the skin of selected rabbits—not necessarily chinchillas—on the lines originally suggested by Fraser and Plummer⁴² (1930); (see also Kolchin: *Journ. Immunol.*, 1933, xxiv, 397), and Dr. O'Meara has had encouraging results with a flocculation test akin to the Ramon method for diphtheria toxin and antitoxin. Although it cannot be assumed that the reacting substances in the flocculation test are identical with the erythrogenic toxin and antitoxin, the method should prove a useful adjunct to other tests, even if it cannot replace them. A similar method has been used in Park's laboratory in New York (see Williams,¹⁷ 1932), to determine tentatively the strength of toxins, but O'Meara has shown that the test can be applied to both unconcentrated and concentrated antitoxins. Research is constantly in progress to improve the potency of streptococcal toxins by studying the growth factors, as has been done for diphtheria toxin. Another recent development has been the selection of the most suitable horses for immunization, by ascertaining the presence or absence of natural

antibody. It is hoped that the future application of knowledge acquired in these studies will ensure the preparation of antitoxin of much higher potency than hitherto. With improved serums some advance should be possible in the therapy of streptococcal disease.

Precipitation and Agglutination

Time does not permit of a consideration of the analysis by Lancefield^{43 44} (1928 and 1933) of the antigens present in the haemolytic streptococci, nor of the careful work on agglutination by F. Griffith^{45 46} (1926 and 1927) and others. The observations of Glover and Griffith⁴⁷ (1931) on the common streptococcal infections of school age are, however, of considerable epidemiological importance. In the words of Okell¹⁹ (1932), the present position with regard to agglutination is "much the same as with the meningococci and pneumococci where in larger epidemics certain types tend to recur, yet in interepidemic periods the organisms may be serologically heterogeneous."

Rheumatism

Acute rheumatism is now regarded by some workers as a condition of hypersensitiveness induced by some focus of streptococcal infection usually assumed to be in the tonsil. The allergic state can be set up by a large variety of streptococci (Swift et al.,⁴⁸ 1931), of which the haemolytic group is probably the most important (Coburn,⁴⁹ 1931). Glover and Griffith⁴⁷ (1931) and Coburn and Pauli⁵⁰ (1932) regard as significant the comparative incidence of haemolytic streptococcal tonsillitis and rheumatic manifestations. In this country and in America investigations are now in progress on the dermal reactions frequently obtained with extracts of streptococci in rheumatic subjects (Collis,⁵¹ 1931). It has been shown by Todd^{37 38} (1932 and 1933) that abnormally high anti-streptolysin titres may be present in the serum of rheumatic fever patients: this is further evidence of the association of acute rheumatism with previous haemolytic streptococcal infection. Antistreptolysin and antitoxin are apparently independent antibodies.

Non-haemolytic Streptococci

Apart from their possible role in the aetiology of rheumatic affections, the non-haemolytic streptococci may produce sub-acute infective endocarditis and occasionally septicaemia. On the whole, measures of treatment so far adopted—vaccines, immuno-transfusion, and the administration of serums prepared against the organisms isolated—have been singularly unsatisfactory.

PNEUMOCOCCUS

I propose to restrict my remarks to certain aspects of the serology of the pneumococcus which have specially interested me, and to leave the more general discussion to Dr. Cruickshank, who has for some years been concerned in an investigation into the value of the serum treatment of pneumonia in Glasgow, and who gave the Milroy Lectures this year on pneumococcal infections.

Preparation and Titration of Serum

Antipneumococcus serum is prepared by the intravenous injection into horses of suspensions of dead virulent pneumococci. In some laboratories living organisms are subsequently injected to complete the course of immunization. The serum, which is type-specific, can be concentrated (Felton,⁵² 1925). Titration of serum is really limited to the mouse protection test, the usual technique being an adaptation of the League of Nations method—that is, determining the smallest amount of serum which will protect mice against 0.01 c.cm. or 0.001 c.cm. of a culture of which the minimum lethal dose is 0.0000001 c.cm. to 0.000000001 c.cm., all the

injections being given intraperitoneally (Felton,⁵² 1925; Parish and Trevan^{53(a) 53(b)} 1930). Goodner⁵⁴ (1928) has suggested a method of titration in rabbits involving the intravenous injection of serum following the intradermal injection of culture, but this test is impracticable for the routine examination of large numbers of samples of serum. Sabin⁵⁵ (1933) has confirmed Goodner's observation that the rabbit intradermic method may give a different evaluation of potency from the mouse intraperitoneal method. *In vitro* methods of titration have also been investigated. The agglutination titre does not correspond with the antibody content measured by the mouse protection test. The precipitation value obtained with the soluble specific carbohydrate (Wilson Smith,⁵⁶ 1932) shows a much closer correlation with the mouse test values than does the agglutination titre, and it is therefore possible that the precipitation test may be useful in controlling the routine immunization of horses. Very recent work concerns the preparation and titration of a stable dried standard of potency based on the Felton standard, Type I (Hartley, Parish, Petrie, and Wilson Smith—*Lancet*, 1933, ii, 91). Dr. Hartley is now engaged in preparing a similar dried standard serum, Type II. A unit is defined as a certain definite weight of the dried product, and not, as in the case of the original Felton liquid standard, as that amount of serum which will protect against a million fatal doses of culture. By the use of this stable standard more accurate potency estimations of commercial serums should be possible. It follows that the labelled unitage of serums will be more correctly stated, and the clinician will thus be better equipped to estimate his dosage.

Treatment with Serum

The very high degree of protection obtained in mice would lead one to expect better results than have been reported in the treatment of pneumonia in man. In the mouse, however, the type of disease produced is not pneumonia, but a general septicaemia. When the serum is given intraperitoneally to mice at the same time as a million fatal doses of the homologous organism the animals can be saved; administered up to four hours afterwards it is still possible to "catch up" this massive infection (Parish,⁵³ (a) 1930). Petrie and Morgan⁵⁷ (1932) have made an exhaustive study of the therapeutic efficiency of serum against much smaller infecting doses, and have saved mice twelve hours after infection, even when there was demonstrable bacteraemia. One is therefore encouraged to continue research on methods of preparing the most suitable suspensions for the immunization of horses, and on the concentration of serum. By the use of improved methods of immunization it has already been possible to prepare, in selected horses, unconcentrated serum of unusually high potency. This unconcentrated serum has the outstanding merit of cheapness. It is now generally agreed that serum reduces the mortality from Type I pneumonia in man, and probably produces a slighter, but still appreciable, effect on Type II cases. The therapeutic efficiency of serums prepared against other pneumococcal types is now being investigated in America. The serum should be given intravenously in large and, if necessary, repeated doses. The patient's clinical condition is usually regarded as the best indication as to further dosage, but Francis⁵⁸ (1933) has recently made use of a test injection of the soluble specific carbohydrate, given intradermally six to eight hours after the administration of serum. If no reaction is observed within half an hour a further amount of serum is given, but if the response is positive additional serum is unnecessary, for recovery has already begun. The work of Alston and Lowdon (*Brit. Journ. Exper. Pathol.*, 1933, xiv, 1) does

not support the view that positive skin reactions are associated solely with recovery from pneumonia.

Reactions After Intravenous Injection

In the early days of concentrated purified serum (Felton,⁵² 1925) the reactions which occurred after intravenous injection were a serious drawback to its clinical use. With improved methods allergic manifestations have seldom been severe, and have yielded to the administration of adrenaline. For some years we have used mice for the detection of batches of serum of abnormal toxicity. A group of ten mice receives intravenously doses of 0.5 c.cm. of each concentrated serum, and the immediate and late effects are observed; if more than two of the animals die the batch is regarded as unsuitable for clinical use. We believe that this test has materially reduced the chance of reactions following the intravenous use of serum.

Serological Typing

One of the great deterrents to the use of serum has been the necessity for the serological typing of the infecting organism. Speedy and more accurate typing is now possible by such methods as those described by Sabin⁵⁹ (1930), Armstrong⁶⁰ (1932), Logan and Smeall⁶¹ (1932), and Valentine⁶² (1933). (See also Sabin: *Journ. Amer. Med. Assoc.* 1933, c, 1584.)

MENINGOCOCCUS

The serology of the meningococcus is on a much less satisfactory basis than that of the organisms we have just considered, for up to the present we have been severely handicapped by the lack of a reliable animal test for the differentiation of toxic products or the titration of serum. The position, however, will be radically changed if recent work, to which I shall shortly refer, is developed and confirmed.

Endotoxin and Exotoxin

The toxin of the meningococcus has usually been regarded as a typical endotoxin which can be liberated from the bodies of the organisms by autolysis, solution in 0.025 N NaOH, lysis by fresh guinea-pig serum, and various other methods (Besredka,⁶³ 1906; Gordon,⁶⁴ 1920; Murray,⁶⁵ 1929; Malcolm,⁶⁶ 1932). In comparative tests in mice, guinea-pigs, or rabbits no very reliable or constant results have been obtained with endotoxin or with dead cocci, although there is no doubt about their toxicity in sufficiently large doses. For many years all attempts to prepare a soluble toxin from the meningococcus have been unsuccessful, but recently Ferry, Norton, and Steele⁶⁷ (1931), Ferry⁶⁸ (1932), and Krestownikowa, Belkina, Dosser, and Lasowsky⁶⁹ (1933) have prepared filtrates from cultures on hormone broth, which were not only toxic to laboratory animals but were also antigenic. Rake⁷⁰ (1933) has studied the biological properties of "fresh" and "stock" strains, although not from the standpoint of toxin production. He refers to considerable differences between recent and old strains, the failure of recent strains to adhere to rigid type-specificity on subculture, and the ready growth, with pellicle formation, of the "smooth" recent strains on hormone broth. It is therefore possible that many workers in the past have failed to obtain a potent exotoxin because they have used "rough" cultures or unsuitable conditions of growth.

Preparation and Titration of Serum

Polyvalent antimeningococcus serums are prepared by injecting horses intravenously with suspensions of from four to eight strains, representative of the different serological

types, and replaced every few months as more recent strains become available. The use of living cultures is emphasized by some serologists. A long course of immunization is advisable, since there is no completely satisfactory test of potency. It is true that Schwartzman^{71 72} (1929 and 1931) has spent considerable time in elaborating the method which is known by his name, and which consists essentially in the intracutaneous injection of the test materials followed by an intravenous injection which makes skin reactions develop. For intravenous injection a weak solution of agar (Sickles,⁷³ 1933) is as useful as a specific meningococcus filtrate. The method is not an easy one to carry out, and requires a careful adjustment of the conditions; nor is there any evidence that it is an accurate measure of antibody. Murray⁶⁵ (1929) refers to the power of serum (1) to neutralize endotoxin (Gordon,⁶⁴ 1920), (2) to promote phagocytosis, and (3) to produce lysis of the meningococcus. He states that on the basis of these properties it should be possible to assess the therapeutic efficiency of a serum. Wadsworth⁷⁴ (1932) has fallen back on the agglutination test, which is admittedly unreliable, and merely indicates that horses have received immunizing injections with a number of strains. We also employ this method, and, like Wadsworth, give a potency index to each serum by averaging the agglutination titres for a few strains. Very recent work concerns the preparation of antitoxin by injecting toxic filtrates into horses (Ferry, Norton, and Steele,⁶⁷ 1931; Krestownikowa et al.,⁶⁹ 1933). The antitoxin content of the serum is apparently titratable clinically and in the laboratory.

Treatment with Serum

Ferry has claimed that antitoxin prepared as described above had a curative and protective action in guinea-pigs injected with a lethal dose of culture, while the ordinary antimeningococcal serum had little or no effect. He states that antitoxin was also of value in experimental meningococcal meningitis in the monkey. If this work should be confirmed and applied clinically to human infections a great advance will have been achieved. There is a good deal of evidence that antimeningococcal serum prepared in the ordinary way is of value clinically, but, as Topley and Wilson⁷⁵ (1929) have stated, it

"has never been given under conditions which from a statistician's point of view could be considered satisfactory. To do this, it would be necessary to give the serum to alternate cases in an epidemic, and to treat both series of cases in other respects in identically the same way. This would mean performing a lumbar puncture on the control cases with the same frequency as on the serum-treated cases, since the effect of this measure alone appears to be of value. Until such a series is tried, we shall not be able to form a just opinion of the actual worth of the serum itself."

I give their view verbatim, for it illustrates the difficulty which the clinical observer may encounter who sets out to form a just opinion of the value of a new remedy. Methods of concentration of serum have recently been introduced, but in the absence of a satisfactory method of titration it is impossible to be certain that some essential constituent has not been lost during the process of refinement. The practical value of the concentrated product has not yet been demonstrated on a large scale. Another form of treatment which is being given a trial at present is the intrathecal injection of antimeningococcus serum to which fresh unheated human serum ("complement") has been added (Fairley and Stewart,⁷⁶ 1916; Bunim and Wies,⁷⁷ 1933). The rationale of this therapy is that, since complement is virtually absent from the cerebro-spinal fluid, and absent from antimeningococcal serum, some benefit might accrue from

making good the deficiency; the mechanism responsible for the sterilization of the spinal fluid might possibly be assisted.

Precipitation

Zozaya and Wood⁷⁸ (1932) have stated that the soluble specific carbohydrates and "nucleoproteins," obtained from the various types of meningococci and other Gram-negative cocci, have very similar properties and are not type-specific. In their experience the precipitable substance did not appear in the serum of horses undergoing immunization until the fourth month or later. Agglutination and precipitation did not run parallel.

GONOCOCCUS

Toxic Products

Tulloch⁷⁹ (1929), writing in the *System of Bacteriology*, has stated that we have "frankly to admit ignorance of the precise nature both of the invasive mechanism and of the disease-producing mechanism of the gonococcus." Cultures, filtrates, and autolysates are all toxic to laboratory animals, the maximum toxicity being after prolonged incubation—which perhaps suggests that so-called "endotoxins" liberated by autolysis of the cocci are the essential toxic constituent.

Serums and Vaccines

As in the case of antimeningococcus serum, there is no adequate method of titration of the so-called "antibacterial" serum used in the treatment of gonorrhoea and its complications. Satisfactory evidence as to its clinical efficacy is also lacking. Vaccines may be of some value in the prevention and treatment of complications, although the published evidence does not lend itself to statistical analysis.

Precipitation, Agglutination, and Complement Fixation

I have previously mentioned that the soluble specific substances and nucleoproteins of all the Gram-negative cocci have very similar properties and are not type-specific (Zozaya and Wood,⁸⁰ 1932). As regards agglutination, serological types exist for the gonococcus as for the other organisms we have considered, but the difficulties in the technique of the agglutination reaction are considerable, and it is not always easy to reconcile the results of different workers. The complement-fixation test (see Tulloch,⁷⁹ 1929; Price,⁸¹ 1930; Osmond and Oliver,⁸² 1929) is of value in the diagnosis of gonorrhoea, but great care and experience are necessary in the technique of the reaction and in its interpretation. It is generally negative in the early stages of the disease, and positive for some time after cure has been effected. It is almost always positive when complications are present, and is therefore of considerable value in the diagnosis of arthritic and gynaecological diseases. The reaction ceases to be of diagnostic significance if specific vaccine treatment has been employed.

CONCLUSION

It will be seen from this brief review of the serology of the pathogenic cocci that considerable advances have been made of recent years in connexion with the Gram-positive group. As regards the meningococcus and gonococcus, the present position is much less satisfactory, but there are already some indications that more efficient antimeningococcal serums may become available. So far the serologist's most valuable contribution to the gonococcus problem has been the complement-fixation test.

REFERENCES

- Staphylococcus*.—¹ *Zeit. f. Hyg. u. Infektionsk.*, 1901, xxxvi, 299.
² *Wien. klin. Woch.*, 1906, xix, 493. ³ *Ann. de l'Inst. Pasteur*, 1914, xxviii, 219. ⁴ *Journ. Exper. Med.*, 1924, xl, 761. ⁵ *Journ. Path. and Bact.*, 1929, xxxii, 717. ⁶ *Ibid.*, 1930, xxxiii, 1. ⁷ *Ibid.*, 1931, xxxiv, 471. ⁸ *Journ. Exper. Med.*, 1931, liv, 315. ⁹ *Journ. Path. and Bact.*, 1931, xxxiv, 593. ¹⁰ *Ibid.*, 1932, xxxv, 251.
¹¹ *Lancet*, 1931, ii, 1180. ¹² *Ibid.*, 1932, i, 506. ¹³ *Journ. Infect. Dis.*, 1933, lii, 6. ¹⁴ *Journ. Path. and Bact.*, 1932, xxxv, 183. ¹⁵ *Journ. Amer. Med. Assoc.*, 1933, c, 1007.
Streptococcus.—¹⁶ *Journ. Amer. Med. Assoc.*, 1924, lxxxii, 301.
¹⁷ *Streptococci in Relation to Man in Health and Disease*, 1932.
¹⁸ *Lancet*, 1928, i, 746. ¹⁹ *Ibid.*, 1932, i, 761, 815, 867. ²⁰ *Journ. of Hyg.*, 1927, xxvi, 84. ²¹ *Journ. Path. and Bact.*, 1927, xxx, 651.
²² *Trans. Assoc. Amer. Phys.*, 1932, xlvii, 161. ²³ *Canadian Public Health Journ.*, 1933, i. ²⁴ *Public Health*, 1932, xlv, 101. ²⁵ *Journ. Hyg.*, 1933, xxxiii, 282. ²⁶ *U.S. Public Health Reports*, 1931, xvi, 3023. ²⁷ *Lancet*, 1927, i, 483. ²⁸ *Ibid.*, 1929, i, 177. ²⁹ *British Medical Journal*, 1931, ii, 599. ³⁰ *Journ. Amer. Med. Assoc.*, 1924, lxxxii, 544. ³¹ *Bull. Johns Hopkins Hosp.*, 1925, xxxvii, 307. ³² *Lancet*, 1933, i, 848. ³³ *British Medical Journal*, 1929, i, 995. ³⁴ *Journ. Exper. Med.*, 1928, xlviii, 769, 751. ³⁵ *Journ. Path. and Bact.*, 1930, xxxiii, 563. ³⁶ *Journ. Exper. Med.*, 1932, lv, 267. ³⁷ *Brit. Journ. Exper. Path.*, 1932, xiii, 248. ³⁸ *Journ. Path. and Bact.*, 1933, xxxvi, 201. ³⁹ *Ibid.*, 1933, xxxvi, 435. ⁴⁰ *U.S. Public Health Reports*, 1932, xlvii, 1043. ⁴¹ *Journ. Path. and Bact.*, 1927, xxx, 521. ⁴² *Brit. Journ. Exper. Path.*, 1930, xi, 291. ⁴³ *Journ. Exper. Med.*, 1928, xlvii, 91, 469, 481. ⁴⁴ *Journ. Exper. Med.*, 1933, lvii, 571. ⁴⁵ *Journ. of Hyg.*, 1926, xxv, 386. ⁴⁶ *Ibid.*, 1927, xxvi, 363. ⁴⁷ *British Medical Journal*, 1931, ii, 521. ⁴⁸ *Amer. Journ. Med. Sci.*, 1931, clxxxi, 1. ⁴⁹ *The Factor of Infection in the Rheumatic State*, 1931. ⁵⁰ *Journ. Exper. Med.*, 1932, lvi, 609. ⁵¹ *Lancet*, 1931, i, 1341.
Pneumococcus.—⁵² *Journ. Infect. Dis.*, 1925, xxxvii, 199. ⁵³ (a) *Journ. Path. and Bact.*, 1930, xxxiii, 729. ⁵⁴ (b) *Ibid.*, 1930, xxxiii, 739. ⁵⁵ *Journ. Exper. Med.*, 1928, xlviii, 413. ⁵⁶ *Ibid.*, 1933, lvii, 139. ⁵⁷ *Journ. Path. and Bact.*, 1932, xxxv, 509. ⁵⁸ *Brit. Journ. Exper. Path.*, 1932, xiii, 96. ⁵⁹ *Journ. Exper. Med.*, 1933, lvii, 617. ⁶⁰ *Journ. Infect. Dis.*, 1930, xlv, 469. ⁶¹ *British Medical Journal*, 1932, i, 187. ⁶² *Ibid.*, 1932, i, 188. ⁶³ *Lancet*, 1933, i, 22.
Meningococcus.—⁶⁴ *Ann. de l'Inst. Pasteur*, 1906, xx, 304. ⁶⁵ Special Report Series, Medical Research Council, 1920, London, No. 50. ⁶⁶ Medical Research Council, *System of Bacteriology*, 1929, ii, 291. ⁶⁷ *Journ. Bact.*, 1932, xxxii, 98. ⁶⁸ *Journ. Immunol.*, 1931, xxi, 293. ⁶⁹ *Ibid.*, 1932, xxxii, 315. ⁷⁰ *Zeit. f. Immunitäts. u. Exper. Therap.*, 1933, lxxviii, 451. ⁷¹ *Journ. Exper. Med.*, 1933, lvii, 549, 561. ⁷² *Journ. Amer. Med. Assoc.*, 1929, xciii, 1965. ⁷³ *Journ. Infect. Dis.*, 1931, xlviii, 339. ⁷⁴ *Amer. Journ. of Hyg.*, 1933, xvii, 412. ⁷⁵ *Trans. Assoc. Amer. Phys.*, 1932, xlvii, 161. ⁷⁶ *Principles of Bacteriology and Immunity*, London, 1929, ii, 934. ⁷⁷ *Commonwealth of Australia Quarantine Service*, 1916, Publication No. 9. ⁷⁸ *Journ. Amer. Med. Assoc.*, 1933, c, 178. ⁷⁹ *Journ. Infect. Dis.*, 1932, l, 177.
Gonococcus.—⁸⁰ Medical Research Council, *System of Bacteriology*, 1929, ii, 239. ⁸¹ *Journ. Infect. Dis.*, 1932, l, 177. ⁸² *Journ. Path. and Bact.*, 1930, xxxiii, 493. ⁸³ *Brit. Journ. Ven. Dis.*, 1929, 281.

AN INVESTIGATION ON THE PHARMACOLOGICAL ACTION OF SOME NEW SUBSTITUTES FOR MORPHINE AND HEROIN

BY

G. NORMAN MYERS, M.D., Ph.D., M.Sc.

BEIT MEMORIAL RESEARCH FELLOW

(From the Pharmacological Laboratory, Cambridge)

The fact that new substances akin to morphine are always being produced by the chemical industry is due to the desire to find, by means of fresh chemical combinations, a substance which is as useful as morphine but has not its harmful effects. During the past few years three new synthetic drugs have been manufactured and extensively used on the Continent as substitutes for morphine and heroin. These drugs are being manufactured in Germany, and are marketed under the trade names of "dilaudid," "dicodid," and "eukodol" respectively. The clinical literature on these drugs is fairly extensive, but little has been published with regard to their pharmacological action. Eddy,¹ however, gives an extensive survey of the literature bearing on dilaudid. The object of this investigation was to determine some of the more important pharmacological actions of these drugs and to compare them with morphine and heroin.