THE

British Medical Journal

THE JOURNAL OF THE BRITISH MEDICAL ASSOCIATION

LONDON: SATURDAY, JULY 1st, 1933

A Paper

ON

OBSTETRICAL ERRORS*

BY

E. FARQUHAR MURRAY, M.D., F.R.C.S., F.C.O.G.
HONORARY GYNAECOLOGIST, ROYAL VICTORIA INFIRMARY; OBSTETRICIAN,
PRINCESS MARY MATERNITY HOSPITAL, NEWCASTLE-ON-TYNE

It is important that the art of obstetrics should be as highly organized as possible. If engaged for a case we must assume full responsibility and not relegate to a nurse or midwife those matters which are essentially medical and obstetrical. Again, if we are usually called in by a certain midwife and find a condition which could have been recognized, and possibly prevented, at an earlier stage, then it is our duty to educate that midwife to the value of closer co-operation. One of the biggest errors made, and still being made, by the profession is that of allowing obstetrics to drift into the hands of midwives, clinics, and whole-time officers. Some of this may be inevitable and even desirable, but doctors are losing the supervisory and consultative interest in this work to the detriment of their practices. The backbone of a successful practice is midwifery, followed by child welfare.

ANTE-NATAL CARE

The supervision, manipulations, and decisions concerned with ante-natal care form one of the most fascinating studies in medical art. The absence of this care is a big error. The multitude of normal labours which occur without such supervision tend to belittle its value and to make people somewhat sceptical. Sooner or later a most devastating tragedy will shock the most casual in this regard. Such comparatively simple and not infrequent matters as the recognition of an occipito-posterior, a breech in a primipara, or a contracted pelvis which invites obstruction may, when properly dealt with, make all the difference between a triumphant success and maternal and foetal exhaustion, bruising, and even death. Ante-natal care is not merely safeguarding the lives of the mother and her child, but also making certain that they will emerge from their joint experience—the confinement—as undamaged from wear and tear as possible. What we have to aim at is that every baby is presenting head first as a polar lie and with its back anterior, and that the relation of the head to the pelvic cavity shows no unfavourable disproportion.

Occipito-posteriors are best dealt with by applying a pad over the child's back, and it is fixed in position with a tight binder. The bladder and bowels are kept empty,

and the patient remains in bed for twenty-four hours. If the procedure fails, repeat it at the next week-end; and if it still fails, then apply it at the onset of labour.

In breech cases the error is often in diagnosis: the buttocks are mistaken for the shoulders. Palpate for the head at the fundus, and dip the fingers into the pelvis behind the pubic bones to feel for a deeply engaged head. Finally, make a vaginal examination. By one or other of these means it should be possible to arrive at a diagnosis. In cases of breech with extended legs, the lower pole of the foetus feels very like the foetal head. If faced with a breech presentation in a primipara ask yourself "Why is it presenting as a breech?" and exclude the presence of a contracted pelvis. If the pelvis is badly contracted leave alone and arrange for a Caesarean section, but if normal or doubtful then do a version and test the head to the brim. In carrying out a version always lubricate the abdomen with soap or oil; this makes the version easier. If failure results then try under anaesthesia. If there is still failure, then by means of external measurements and a careful estimation of the pelvic capacity per vaginam decide whether or not there is sufficient room for the breech to come through, always allowing for manipulations for possibly extended arms or legs. Failure of judgement in this respect will end in a very difficult extraction with a stillborn child, or, in gross cases, in retention of the head above the brim. Lubrication of the hands, gloved or otherwise, with ethereal soap makes the corrective manipulations during delivery much easier.

Contracted pelves are usually contracted at the brim or throughout. Some are only contracted at or towards the outlet, and they are the more treacherous, as the head engages well and gives an appearance of normality, while late in labour the moulding and caput may suggest proximity and invite a delivery with forceps. Pelvic measurements, therefore, should always be taken of the outlet as well as of the brim. With the more common types of contracted pelvis the head is high and even overriding the brim. This in a primipara is always suspicious, as the head is usually deeply engaged for some weeks before term. Measure the pelvis and test the head to the brim, if necessary under anaesthesia. With a reasonable degree of care it is possible to divide cases into those that *will*, those that *won't*, and those that *may* come through. The first are normal, the second are for Caesarean section, and the third group should be as small as possible. Such a thing as a trial labour should not be allowed to occur outside a home or hospital, unless expert assistance is available. There is a place for induction of premature labour, but only in cases with minor degrees of contraction, occasionally in primiparae, and rarely in atypical forms of deformity.

* Read at a meeting of the Border Division, March 23rd, 1933.

Absence of ante-natal care may allow the toxæmia of pregnancy to produce the kidney of pregnancy and its major crisis, eclampsia. Regularity of the bowels, routine urine testing every fourteen days till the last weeks, when it is done weekly, and, most important of all, encouraging the patient to report any unusual symptom will modify this danger. Oedema is not, as commonly believed by the laity, due to pressure, but is really a manifestation of renal deficiency. Don't hesitate to starve an albuminuric patient. Fluids are essential, and food may be withheld for a week or more with impunity. Fruit juice and stewed fruit and sugar may be given after a period and before proteins are added to the diet. In actual practice the routine testing of urine specimens would be best dealt with by the nurse if it can be so arranged.

The utmost need for ante-natal care is in the case of primiparae, as their first pregnancy and labour is a trial of their pelvic girdles and their metabolism. Multiparae must be watched carefully also. Their kidneys may be damaged, their lax uteri may allow malpresentations. Four such, with previously normal histories, arrived at a hospital with ruptured uteri. They had malpresentations which had been corrected by version. Three of them died. The malpresentations or the versions should have been dealt with earlier.

LABOUR

The dramatic obstetrics which impressed us as students, the difficult versions, embryotomies, decapitations, and so on, were largely owing to the absence of organized ante-natal care. The contrast between a timely Caesarean section in a case of badly contracted pelvis and a similar case dealt with late in labour *per vias naturales* requires no comment.

A great error during labour is to withhold chloroform. The patient's cries are unnerving the household, the atmosphere is tense, the doctor may be urged to terminate labour. Delivery may be the worse possible treatment in such a case. The patient, a primipara, may be only half-way through the first stage of a perfectly normal labour. Put her and the family to sleep by administering chloroform to the patient. Do not forget to do this when called in by a midwife to a case which you have never seen before. Quieten the household, get the history, and make a careful abdominal examination; then wash up and make a careful internal examination, and arrive at your final decision after sizing up the general condition of the mother and the child. Morphine may be given, and also bromides and chloral; but morphine should not be given if delivery is likely to take place within four hours of the injection.

Failed forceps, when analysed, are divisible into those due to errors of omission and those due to errors of commission. The former are neglect of ante-natal care, and the latter are forceful attempts to anticipate natural processes or incorrect attempts to remedy the neglect of ante-natal care. Cases admitted to hospital with the os uteri one-third or one-half dilated are merely given time to complete the natural process. The injection of pituitary extract is open to criticism. Having known of it being given to primiparae, and even primiparae with badly contracted pelvises, I believe that the following is fairly sound advice: never give pituitary extract until you are fully conversant with obstetrics, and never to a primipara. Reserve its use for multiparae with a fully dilated os, and a head well engaged or low down, when it is merely a question of pushing or gently pulling the head over the perineum. Always dilute the extract and inject it gradually under anaesthesia. Another common cause of failed forceps is failure to recognize that the baby's head in an occipito-posterior case is not yet in the antero-posterior of the outlet. It is not always easy to detect, especially if there is much of a caput. It is still more

difficult if the patient is lying in the left lateral position. Great help is obtained by delivering these patients in the cross-bed position, using either Clover's crutch or the obstetric straps to obtain the lithotomy position. The exact relation of the head to the outlet is then easily detected by inserting the left hand in the vagina in order to palpate it thoroughly, or even to feel for the ears. Once the condition is recognized, rotate into the correct position and apply forceps, but if impossible to correct manually then correct with the forceps and reapply them, after correction, before extracting the head. Another error is to attempt extension of the head with the forceps before the occipital protuberance on the child's head has descended below the symphysis pubis: feel for it. In the third group of failed forceps are degrees of pelvic contraction. Gross cases are obvious by abdominal examination alone. Moderate cases require a careful examination under anaesthesia of the pelvic girdle and foetal head. Ask yourself what has already been done and what remains to be done. The only possible alternative at this stage, provided the child is alive, may be a Caesarean section or, occasionally, a version. Do not be misled by the small size of the os in bad cases of contracted pelvis into imagining that the patient is still in the first stage. The os may never dilate any further owing to non-descent of the head.

It is a mistake to allow tears to occur, to fail to repair those that do occur, and to admit that they have occurred. The vexed question of tears is so bound up in the public mind with neglect that it is useful to be frank with ourselves and admit that they do arise in a very large number of primiparae. Much can be prevented, much is inevitable. The really important point is to realize when a tear is likely to be inevitable, and to anticipate it by a timely episiotomy, which should be median in position. The indication for this is a tense ostium vaginae which has produced a secondary caput on the child's head, and where, with each pain, there is blanching of the vestibule. This may, if neglected, lead to two sequels: a bursting tear of the perineum, vagina, and labia, and such laceration of the sphincter urethrae as to result in laxity and incontinence of urine later. Failure to repair tears adds to the risk of puerperal sepsis, and possibly, later, of prolapse developing. As a gynaecologist as well as an obstetrician, and knowing how uniformly well properly repaired lacerations heal, I cannot but believe that, whether the blame rests with the profession or, more likely, with the midwives, there is simply no attempt to repair much vaginal trauma. One error might be excused—namely, failure to recognize that an intact perineal surface is associated with concealed lacerations. Always examine for them. Repair cannot, with rare exceptions, be accomplished by means of silkworm-gut sutures; they produce an external appearance of approximation, but they rarely manage to produce actual anatomical apposition. In order to effect this catgut should be used, either buried or superficial. Repair the vaginal tube, repair the rectal tube and sphincter when necessary, bring the levators together between them, and finally repair the skin edges. Tell your nurses not to count the stitches or to admit tears. Tell them to admit bruising; the patient and her husband will thoroughly understand, and their minds be peaceful. This is not duplicity, it is common sense. A "tear" sounds horrible, a "bruise" sounds reasonable. When the baby has arrived examine at once for tears. If trivial, repair at once with catgut; but if severe, then wait until the placenta has arrived and do a proper and deliberate repair in the cross-bed position.

Another error is mismanagement of the third stage. The main danger is at night when time hangs heavy and one wants to "get it over." We are all human; but

unless an anaesthetic has been administered, then it is useful to note the time of delivery and to allow at least fifteen to twenty minutes to elapse before worrying about the placenta. Leave the uterus alone, or at most feel it from time to time. It will not let you down. If an anaesthetic has been given, then the uterus *may* let you down, so it is best to massage it gently and note when the placenta has left. Should it be necessary to go in for the placenta, then always give an intrauterine douche afterwards. Failure to do this is asking for sepsis. Never allow a binder to be applied until at least one hour after delivery, and palpate the uterus from time to time.

PUERPERIUM

The first three days are probably the most important. Get the bowels open early. An enema is an adjuvant, not an essential. Opening medicine should be given in twenty-four hours' time or earlier. Always be on the look out for sepsis. Influenza, pneumonia, debility, yellow atrophy of the liver, and breasts are often provided as an explanation for a sequence of events which are in reality due to septic processes. The headaches, shivers, malaise, and temperatures of a septic process are often attributed to influenza if it is prevalent. Chest conditions of septic origin are often manifest within three days of delivery, and to the inexpert might be believed to be of accidental origin. Rapidity of pulse, with practically no other symptom than increasing weakness, can be due to hyperacute sepsis, but may be attributed to shock or haemorrhage. Jaundice, with only moderate pyrexia, or even in the absence of pyrexia, can be due to a hyperacute haemolytic streptococcal infection and not to hepatic derangement. Septic temperatures and engorgement of the breasts are common on the third day. Always be on your guard about sepsis; always suspect it; and always look at the perineum and lochial discharges. Never give a vaginal douche without clear proof of vaginal sepsis. In the vast majority of cases it is merely perineal sepsis: therefore arrange for local swabbing with peroxide, 20 volumes, three times a day, followed by a jug douche of boric or weak lysol; and this is all that is needed. If there is vaginal infection apply similar treatment to the vagina. Very occasionally the uterus is to blame, but only rarely in the first instance. The difficulty of adequate nursing in outlying areas is obvious. The fact remains that such may be essential.

Three errors remain to be dealt with, although there are many others. Lysol, or any other such substance, is quite useful as an inhibitor of bacterial activity. It does not, except in its pure state, act as a reliable antiseptic. The only reliable antiseptic is a preparation of mercury, such as the perchloride (1 in 1,000) or the biniodide (1 in 1,000). Swabs, hands, and individuals passing from case to case can never be efficiently disinfected if they depend on lysol. With rare exceptions, never examine a patient when she comes to book you. She is nervous and self-conscious, especially if she is a primipara. Internal examinations are a rarity under modern conditions of ante-natal care. Do not tell a woman that she must never have another baby unless such an edict is based on the soundest medical, surgical, or obstetrical grounds. Such advice—in cases met with—has rarely been justified. The patient fears falling pregnant, and is panicked if she does. If such advice is given, whether justified or not, it is strangely out of place that no further advice follows. Finally, we must realize that the very essence and ethics of our existence are based upon the conservation of life itself. Therefore, safeguard the mothers with ante-natal care. Remember, when you are called in to see a case, the importance of time, sedatives, and careful decisions in labour, and always be on the look out for sepsis during the puerperium.

PANCREATIC LITHIASIS

BY

S. N. SENNETT, M.B., B.S.LOND., M.D.BERL.

CAPETOWN

Pancreatic lithiasis is a rare condition; the symptomatology is bewildering and the diagnosis beset with many difficulties. A review of the history and symptomatology may therefore prove interesting inasmuch as it may enable a diagnosis to be made more frequently in the living. Although pancreatic lithiasis was first observed by Regnerus de Graaf in 1664, its great rarity is such that nearly two and a half centuries later (in 1903) Lazarus was able to collect only eighty such cases described in the literature. In most of the cases the calculi were found at necropsy; in some the diagnosis was made of chronic pancreatitis of probable calculous origin, and in several instances the diagnosis was only made at operation (Lindsay, Link, Seeger, Hartman, etc.). In recent years, however, with the aid of roentgenology, a few more cases have been described, so that the condition is not quite so rare as had been supposed. Out of 11,000 necropsies at Guy's Hospital there were three cases only, and Opie in 1910 found two among 1,500 necropsies at the Johns Hopkins Hospital, all in elderly men. Rindfleisch, in 2,000 necropsies at Königsberger Medical Clinic, found only three cases. In the Mayo Clinic, where at every laparotomy of the upper abdomen the pancreas is palpated, eight cases have been discovered in fifteen years. Pancreatic calculi occur more frequently in men than in women: this is also consistent with the observation that chronic pancreatic disease is much more frequent in males than in females. Among the fifty-seven cases collected by Lazarus, forty-seven were in men and ten in women, and of these, thirty-seven were between the ages of 30 and 50 years.

The number of calculi varies greatly: usually five to ten are present; in rare instances there is a solitary calculus, but as many as 300 have been found (Lancereaux). A characteristic of multiple pancreatic calculi is that they are usually not faceted (unlike gall-stones). They vary in size from mere sand to masses the size of a walnut, and may be smooth or rough, soft or hard, and white, brown, or yellow. They consist chiefly of calcium carbonate or calcium phosphate, or even occasionally of calcium oxalate, which is somewhat surprising, as the pancreatic juice does not contain these salts, and is strongly suggestive of an inflammatory aetiology. This fact is, however, of great importance, as it is the lime salts which make the stones opaque to x rays. Rarely they consist of organic substance, and occasionally of lime salts, together with organic protein debris. It is of interest to note that pancreatic calculi have also been repeatedly found in animals, particularly in cattle.

Calculi probably never form in a healthy pancreas. All the evidence points to primary inflammatory changes in the ducts and in the parenchyma of the gland, with occlusion of the ducts (stasis) or invasion by bacteria as a secondary factor. Giudiceandrea demonstrated the colon bacillus with numerous long bacilli in pancreatic calculi, while in a second case he found numerous cocci and bacilli in the centre of a stone. On the other hand, Lindsay could not discover any bacteria in his calculi.

SYMPTOMS

Pancreatic lithiasis may be unaccompanied by noteworthy local symptoms indicating disease of the organ. Lazarus gave an account of such a case, in which death occurred with diabetes and pulmonary tuberculosis. At necropsy the pancreas was found to be converted into a