however—namely, the group of marathon runners; in this group the heart was very much larger than the body weight would have led one to predict. It would be impossible to find any group of men whose circulatory mechanism is functionally more efficient than that of Olympic marathon runners, and it would be absurd to regard the cardiac enlargement in this case as pathological. Similar findings have been reported by other workers^{4, 5} in men who participated in other types of sport entailing prolonged and severe exertion, notably ski-running and cycling. The same is true throughout the animal kingdom, the hearts of athletic animals being heavier in proportion to their body weight than those of animals which lead a more sedentary life. One is bound, therefore, to conclude that slight cardiac enlargement is, by itself, insufficient evidence on which to base a diagnosis of heart disease; although, when considered in conjunction with the other clinical findings and the habits of the patient, it may be of great value in diagnosis.

I have confined my attention to cardiac enlargement because I do not believe that one is ever justified in regarding a heart which is too small as pathological, unless there be additional evidence to support the diagnosis. Bedford and Treadgold² found a heart with a very small transverse diameter in a member of the Schneider Cup team, and Ellis and I³ found a similar heart in the Olympic champion for the 100 and 200 metres!

In conclusion, let me summarize my remarks by endeavouring to answer the question, "What can x-ray

examination contribute to the diagnosis of pathological cardiac enlargement?" I would submit that it can make at least three important contributions.

1. It can tell us all that we can learn from percussion and palpation regarding cardiac enlargement, and it can do so with a much higher degree of accuracy than either of these methods.

2. It can give us reliable information regarding slight changes in heart size and the slighter degrees of cardiac enlargement, even under the most unfavourable circumstances. Although slight cardiac enlargement by itself is not sufficient ground on which to base a diagnosis of heart disease, it may sway the balance when considered in conjunction with evidence from other sources.

3. X-ray examination can reveal changes in the shape of the heart outline. These changes tell us which chambers are enlarged, and so indicate the probable cause of the enlargement. This, in my opinion, is its most important contribution. The frontal silhouette alone gives us a far more accurate idea than does percussion of the shape of the heart, whereas examination in the two oblique positions yields additional information of which percussion can tell us little or nothing.

REFERENCES

- ¹ Hodges, F. J., and Eyster, J. A. E.: *Arch. Int. Med.*, 1926, xxxvii, 707.
- ² Bedford, D. E., and Treadgold, H. A.: *Lancet*, 1931, ii, 836.
- ³ Bramwell, C., and Ellis, R.: *Quart. Journ. Med.*, 1930-1, xxiv, 329.
- ⁴ Deutsch, F., and Kauf, E.: *Heart and Athletics*, 1927.
- ⁵ Herxheimer, H.: *Klin. Woch.*, 1924, iii, 2225.

SOME OBSERVATIONS ON RICKETS, WITH SPECIAL REFERENCE TO ITS OCCURRENCE IN PIGS

BY

JOHN POOL MCGOWAN, M.A., M.D.

ABERDEEN

(With Special Plate)

In describing these observations on rickets in pigs, which were made during the period 1922-31, the question of the causation of the disease, whether due to deficiency of calcium, phosphorus, or vitamin D, comes under discussion.

CALCIUM DEFICIENCY

Since rickets was first observed the question of calcium deficiency as a cause has always been a major point for debate—for example, by Korenchevsky (1922).

Hutyra and Marek (1913) deal at great length with this aspect of the disease in pigs. Elliot, Crichton, and Orr (1922), also working with pigs, alleged that rickets in these animals could be prevented by adjusting the CaO percentage of their ration to that of the P_2O_5 . As, however, 10 c.cm. per pig per day of the antirachitic substance cod-liver oil was administered in the basal ration, the condition resulting from the calcium deficiency could not have been rickets. Evidence will be adduced later to show that, in all likelihood, the lesions produced were those of osteoporosis or, very exceptionally in the circumstances, of osteoporosis complicated by a condition of rickets already present before the animals were submitted to the experiment.

During the years 1922-6 the opportunity occurred of observing, among large breeding stocks with an annual population of hundreds of susceptible pigs, the effect of "balancing" in the ration the CaO with the P_2O_5 . Rickets as a natural disease in pigs occurs most frequently in January and February, and in the stocks just alluded to, fed on this "balanced" ration and during these months, hardly a pig which could be the subject of the

disease escaped it. Zilva, Golding, and Drummond (1924) performed experiments on pigs with results which confirmed these observations, and showed that, while the administration of cod-liver oil prevented the occurrence of rickets, the "balancing" of the CaO with the P_2O_5 of the ration was quite ineffectual. Subsequent to the publication of their paper a "repeat" of their experiments confirmed these results. Later still, the specimens from an experiment, in which groups of "summer" pigs (free, therefore, from "latent" rickets) were fed, some on a calcium-deficient ration, others on this ration with the addition of a vitamin D preparation, and others again on this ration with cod-liver oil added, were examined by me. Advanced osteoporosis with fractures of the bones was present in all groups, and particularly in that receiving cod-liver oil. Rickety changes were entirely absent. In 1931 I was requested by Dr. R. D. Sinclair* (who has allowed me to refer to his results) to report on the specimens derived from a somewhat similar experiment. Groups of pigs were fed, some on a calcium-deficient ration, others on the same ration with exposure to sunlight, others on the same ration with exposure to ultra-violet light, and others again on the same ration with the addition of cod-liver oil.† Osteoporosis of an advanced type, often with fractures, was present in the bones of all the groups: rickety changes were absent (see Special Plate).

The following provisional interpretation of these collected observations seems justified—namely, that calcium deficiency produces osteoporosis which cod-liver oil is practically powerless to prevent; and that, while administration of calcium does not prevent "latent" rickets

* *Scientific Agriculture* (1933, xiii, 489) records the other details of this experiment.

† Observations were also made on a parallel series of groups where the P_2O_5 was "balanced" with $CaCO_3$. Osteoporosis, less in degree than in the calcium-deficiency series and worst in the basal group, was present except in the cod-liver oil group. The most feasible explanation of these phenomena would appear to be along the lines of a condition of alkalosis interfering with assimilation, remedied to some extent by a state of acidosis produced by vitamin D (*vide infra*).

from developing into "patent" rickets, or rickets itself from appearing, cod-liver oil is very effective in this regard.

Hess (1930) and Park (1923) stress the point that deficiency of calcium in the diet does not produce rickets and that the administration of calcium will not cure or prevent it. On the other hand, cod-liver oil has no effect on the condition caused by deficiency of calcium. A type of so-called rickets—the low-calcium high-phosphorus variety—has been described (McCollum and Simmonds, 1925) as occurring in experimental rats. The histological description of this condition as a whole is indicative of osteoporosis: some slight appearances present, which might conceivably be regarded as being of a rachitic nature, could be attributed to the animals having already been the subjects of the disease before they were placed on experiment. This further evidence from the literature therefore goes to confirm the conclusions already stated. Korenchevsky (1922) believes that osteoporosis may indeed occur in the course of rickets, being induced by a state of cachexia, debility, and lack of nutrition developing in the course of the disease. This, however, is an accident, and not the essence of the condition. The basic histological changes present in such states as scurvy, scurvy rickets, general cachexia, and the calcium deficiency just referred to—conditions markedly different in regard to aetiology and cure—are atrophy and osteoporosis, while in rickets they constitute pronounced hypertrophy. All these diseases have much in common as far as the symptomatology of the bones is concerned (however they may be modified by intercurrent influences, such as movement, fractures, stresses, and strains), because of the interference with the integrity of the epiphyseal grid, as described (McGowan, 1924). It should be realized, however, that rickets is aetiological and pathogenetically—as it is morphologically—quite distinct from the other conditions just referred to above.

THE AGE FACTOR

Before considering the possible relation of phosphorus or vitamin D deficiency to the occurrence of rickets some fundamental points in connexion with the disease may be dealt with. Hess (1930) has emphasized the point that rickets occurs much earlier in infants than is generally imagined. Schmorl (1909) found it present histologically in four children of about one and a half months old, and Park (1923) quotes Dunham as having observed a very advanced case in a premature child of one month. The subject of "latent" rickets, where the disease is usually observed accidentally post mortem, and that of "patent" rickets, where symptoms are exhibited during life, is thus introduced again. These conceptions are matters of great moment in an understanding of the disease. A concrete example in connexion with pigs, recorded elsewhere (McGowan, 1926), illustrates this. At that time a considerable number of cases had been collected (these have been added to in the course of the present investigations) in which histological evidence of the presence of rickets in pigs was forthcoming at the tenth to the twentieth day of life, while clinical symptoms do not usually appear till the animal is from 10 to 12 weeks old. Such occurrences have important implications. In this way one may explain, as already mentioned, the finding at times of morphological appearances of rickets in pigs which have been fed on a calcium-deficient ration. Further, all experiments hitherto conducted on the aetiology of rickets in rats, dogs, pigs, etc., have necessarily been performed on weaned animals, and the fact that such experimental animals might quite well have been the subjects, at the commencement of the experiment, of rickets acquired naturally in this way introduces serious complications in regard to interpretation of results. A converse situation exists in which rats and their young, supplied without the knowledge of the experimenters with vitamin D, have created difficulty in vitamin D assay by their failure to become rachitic on the diets usually employed for this purpose.

The possible reasons for the appearance of rickets during the suckling stage have been discussed (McGowan, 1926) in relation to the occurrence of an allied disease in pigs—"iron deficiency." The conclusion was arrived at that iron deficiency was due to a deficiency of iron in the milk relative to its protein content and to the growth potential of the young animal. These inimical conditions could be overcome by ensuring that the sows were "good milkers," and so supplied large quantities of milk to their offspring, the excess protein of which would be burnt off, leaving, however, the large quantity of iron of the excess milk to obviate the occurrence of anaemia. A parallel was drawn in regard to the occurrence of rickets, and an instance was quoted in which rickets in lambs born in January from mothers grossly fat (and poor milkers) was prevented by the simple expedient of providing an abundant supply of milk from foster mothers, all other conditions remaining the same. In this connexion it is also of interest to note the practice of Shorthorn breeders, who rear their December-born bull calves indoors each on the milk obtained from two or more foster mothers. By this means large, overgrown specimens of the breed are obtained, yet with an entire absence of rickets, in spite of the indoor life and absence of sunlight.

CALCIUM AND PHOSPHORUS DEFICIENCY

It now becomes necessary to discuss the particular ingredient of milk which is deficient when rickets occurs. Sow's milk contains 0.4 per cent. CaO and 0.35 per cent. P_2O_5 . Korenchevsky (1922) pointed out that human milk contains no excess of CaO over the minimum requirements of infants, and that with a growth potential over the average a condition of relative calcium deficiency could easily arise. In view of the excessive rate of growth in modern breeds it is reasonable to assume that the same conditions exist to an even greater extent in pigs, and that the CaO content of pig's milk represents an irreducible minimum. If this is the case, considering the smaller absolute amount of P_2O_5 compared with CaO in the milk, and considering the fact that CaO is mainly employed in the formation of bone, that CaO and P_2O_5 collaborate in the body* in this function, and that the body has a very great need for P_2O_5 in connexion with the formation of the soft tissues, it would seem that deficiency of P_2O_5 in milk must be a contingency more to be expected than deficiency of CaO.

As regards the protein content of milk in relation to the disease, it should be noted that swine colostrum contains 15.56 per cent. of protein as compared with the 6.6 per cent. of ordinary sow's milk.

Hess (1930) has stressed the importance of albuminized milk in causing rickets. He points out that egg white increases the intensity of the disorder because it increases the growth-promoting properties of the diet: the protein of cow's milk has come under a ban for similar reasons.

INFLUENCE OF GROWTH RATE

Hess (1930), Park (1923), and Mellanby (1922,* 1925) have emphasized the great importance of growth—especially in cases where there is growth in bulk without proportionate growth in bone—in connexion with the appearance of rickets. It has been stated that without excessive growth rickets could hardly appear. The great susceptibility of premature infants because of their rapid growth in proportion to size has been emphasized by everyone. Moreover, Park (1923) has stated that the liability to rickets varies directly with the youngness of the organism and the speed of new tissue formation.

* It is necessary to mention that in Mellanby's experiments (1925) on the "cereal" effect in rickets his diets are very deficient in calcium, and that his illustrations of sections of rib junctions show osteoporosis and atrophy *only*, without, so far as one can judge, any trace of rickets. It should also be noted that the worst offender as regards production of "rickets"—oatmeal—contains 16.1 per cent. protein and 7.2 per cent. fat, as compared with the 7.9 per cent. protein and 1.4 per cent. fat of the much less noxious substance, wheat-flour.

Again, the frequency of the appearance of rickets during periods of rapid growth has also been remarked upon. On the other hand, Park avers that the older the animal—and hence slower growth—the more difficult is the production of rickets, and the more independent of the antirachitic vitamin the animal seems to be. Vitamin A, by favouring growth, has also a furthering influence on the appearance of rickets. Finally, Mellanby (1922) points out that the administration of thyroid to dogs increases the rachitic changes by increasing metabolism, an observation in harmony with the finding of John Thomson that cretins, if given thyroid, develop rickets, and for this reason require the simultaneous administration of cod-liver oil.

All modern breeding tendencies are towards the development of an animal which will grow fast and mature quickly. In such animals the growth is disproportionately great in the earlier stages. Thus, while a newborn pig may treble its original size in two weeks, it requires the next six weeks to treble in turn the size at this stage, showing thus the greater comparative rapidity of growth at the earlier period. This is a point of considerable importance in dealing with the early appearance of rickets in pigs. From such considerations as the above the conclusion was drawn (McGowan and others, 1931) that rickets was due not to a relative deficiency of calcium but to a relative deficiency of phosphorus.

TRUE FUNCTION OF VITAMIN D

We have now to consider the part that vitamin D plays. Because vitamin D can cure and prevent rickets it would be illogical to conclude from this alone that a deficiency or lack of it was the cause of the disease: on such reasoning syphilis could be ascribed to a deficiency of salvarsan. Theiler and Green and Du Toit's (1932) observations on aphosphorosis in cattle, and Wilson's (1932) observations on osteomalacia, two conditions allied to rickets, which develop in spite of strong insolation, show that something more fundamental than vitamin D—in this instance lack of phosphorus in both cases—is involved in their initiation. Further, the occurrences in lambs and Shorthorn calves already referred to demonstrate that vitamin D is unnecessary for normal or even excessive healthy growth when other conditions—especially an adequate amount of P_2O_5 —are optimal.

It must be emphasized that these observations are not meant to minimize the important part that vitamin D plays in rickets. It is a question of the logical ascription of a consequence to a cause. To use a crude simile, vitamin D might be regarded as playing the part of governor of a steam engine, while phosphorus would be something more vital—say, the piston. Elsewhere (McGowan and others, 1931; McGowan, 1933) evidence has been adduced to show that the probable mode of action of vitamin D, in its preventive and curative roles, is by the setting free of phosphoric acid from the lipins, thus increasing the acidity as well as the available phosphate of the body. The acidity itself may have an important effect (through rendering possible increased secretion of HCl in the gastric juice) on the intestinal reaction, and hence on absorption of adequate amounts of P_2O_5 (and CaO). Browning (1931) quotes several observers—for example, Bacharach and Jephcott (1929)—as having shown that vitamin D renders the contents of the intestine acid.

DISCUSSION

Histological and other observations made during the years 1922-31 showed that a fair proportion of the rickets cases in pigs appeared in "latent" form during the suckling stage, and that the subsequent feeding of pigs on a diet high in $CaCO_3$ —especially in winter conditions—resulted in the appearance of "patent" rickets,

due either to an aggravation of the already existent "latent" disease or to a *de novo* initiation. Elsewhere (McGowan, 1933) evidence is forthcoming that the mere ratio of CaO to P_2O_5 in the diet is of little importance in relation to the ultimate metabolism of these substances, and that the two matters of real moment are the acidity of the gastric juice and the occurrence of great excess of CaO ($CaCO_3$) in the diet. Other important factors, however, require detailed mention—namely, the percentage composition of the diet as regards CaO and P_2O_5 and the related circumstance of the amount of food that has to be eaten before the animal is "filled" and its appetite appeased.

On the average, milk may be taken as an example of the ideal food. As already stated, it contains, in pigs, 0.4 per cent. CaO and 0.35 per cent. P_2O_5 ; so that for the introduction of a comparatively small amount of CaO and P_2O_5 , a considerable bulk of food requires to be taken to satisfy the animal's appetite. In spite of this consumption of an apparently dilute fluid, in the fastest growing period of the young pig (from birth to the age of 3 weeks) a kilogram of sow's milk, containing 4.0 grams CaO, causes an increment of 250 grams in the body weight (Prazmowski, 1932). Similar conditions hold good for another natural food—grass—in which the composition as regards CaO and P_2O_5 is of the order 0.25 per cent. Suggestions of this kind from Nature cannot be lightly disregarded, and they would appear to give a clear indication of the "fuel mixture," for which the body machine is built. If one turns to a consideration of the "concentrates" employed in the unnatural and artificial feeding of animals by man, one finds that in these—on the basis of what has just been said, and considering the bulk required to satisfy the appetite—about three or four times more P_2O_5 is used than is necessary. At the same time, in such a ration only one-quarter of the necessary CaO is present—taking milk as a standard. When, however, such a ration is "balanced" by means of $CaCO_3$ so that the CaO/ P_2O_5 ratio is unity, the diet contains about three to four times the requisite CaO. Under these conditions a simple calculation will show that a pig of 100 lb. weight, eating 5 lb. of such food per day, would require to secrete more than 20 litres of gastric juice per day before its gastric contents could give a reaction to the acid side! Such considerations afford a ready explanation of the experimental production of the high-calcium low-phosphorus type of rickets, and also of the failure of the procedure of "balancing" the P_2O_5 by means of $CaCO_3$ to prevent the occurrence of rickets in the cases referred to at the beginning of this communication. They likewise indicate a gross economic waste*—directly, because the added $CaCO_3$ renders useless and unabsorbable the P_2O_5 supplied in an amount three to four times more than necessary; indirectly, because of the production of disease by such manipulations.

Tetany and convulsions were observed very frequently in the rachitic pigs on this "balanced" ration. Shohl and others (1931) have shown that the condition of the blood in rickets is one of alkalosis, which brings about a lowering of the ionized calcium in the blood; and Zucker, Johnson, and Barnett (1922), and Zucker and Matzner (1923) have stressed the fact that in rickets the reaction of the intestinal canal is alkaline. In the present instance the administration of such quantities of $CaCO_3$ must tend to exaggerate further this alkalotic condition of the body and its juices. Thus, it can easily be imagined that in such circumstances a condition of "achlorhydria" could supervene, with the result that there would be difficulty in absorbing not only P_2O_5 but even CaO.

* The advisability of feeding concentrates incorporated with a bulky non-concentrated food such as turnips, cabbages, etc., seems apparent.

The findings recorded elsewhere (McGowan and others, 1931; McGowan, 1933) in connexion with rabbits fed on high CaCO_3 diets are also of interest in this connexion. Here large quantities of calcium are excreted in the urine as CaCO_3 by an interaction in the kidney between CaCl_2 and NaHCO_3 , whereby CaCO_3 on the one hand and NaCl and HCl on the other are formed. The consequent replacement of the alkaline NaHCO_3 by NaCl and HCl brings about a tendency to reduction of alkalosis and to a swing over towards acidosis, such as is seen in the administration of salts of calcium with fixed acids.* An example of this is seen in calcium chloride therapy, where the acidosis so produced plays a considerable part in the cure of tetany by increasing the ionization of calcium (Gamble, Ross, and Tisdall, 1923).

The peak of occurrence of rickets in the investigation I am recording took place in January and February. In the months immediately prior to this there were fewer "patent" cases because sunlight had acted as a protection in the suckling stage, while the relative infrequency of the disease in the months immediately succeeding may be attributed to sunlight acting as a curative agent to the older pigs. It goes without saying that in the summer months both curative and protective roles would be operative.

Reference has already been made to aphosphorosis in cattle and its relation to rickets. Theiler (1932) has shown that the histological appearances in this condition are essentially those of rickets, modified, of course, by the fact that it occurs in adult animals in which the epiphyses are no longer actively functioning. Its occurrence is due to lack of phosphorus in the pasture; administration of phosphorus cures it; and it develops in spite of abundant insolation. The significance of these phenomena in relation to an understanding of vitamin D action has already been indicated. Osteomalacia in human beings has many points in common with this disease. It can also occur in spite of considerable insolation (Wilson, 1932), and in such circumstances it is amenable to the administration of tricalcium phosphate. Usually, however, osteomalacia presents a wider picture. The clinical symptoms, including as they do tetany, may be regarded as a mixture of those associated with aphosphorosis and acalcicosis. The history of its onset indicates a deficiency not only of P_2O_5 but also of CaO in the diet. There is hypocalcaemia, while the histological appearances of the bones, with the presence of osteoid tissue coupled with the extensive removal of the bony trabeculae, may be interpreted as a mixture of those appropriate to rickets and to osteoporosis. As regards treatment the beneficial effect, in most cases, of administering P_2O_5 (and vitamin D) is considerably added to by the simultaneous exhibition of CaO .

It has already been stated that rickets is of frequent occurrence in premature infants. In such cases it is very often associated with tetany, and it seems possible that there is here, also, a mixture of aphosphorosis and acalcicosis, brought into being by the needs of the rapidly growing body outstripping the supply both of P_2O_5 and of CaO contained in the milk.

SUMMARY

In pigs deficiency of calcium in the diet produces osteoporosis, and cod-liver oil, vitamin D, and sunlight are practically without avail in preventing or curing this condition. On the other hand, the "balancing" of the P_2O_5 of the ration with the CaCO_3 (especially when the ration consists of "concentrates") not only fails to prevent the appearance of rickets, but would seem to aggravate the condition if already present in a "latent" form,

or to initiate it. Cod-liver oil, vitamin D, and sunlight, however, prevent and cure rickets.

Emphasis is laid on the early appearance of rickets in pigs, and the bearing of the existence of this "latent" form on an understanding of the disease and on the interpretation of experimental results discussed. Rickets is regarded as being due to a deficiency of phosphorus in the diet relative to its protein content and the growth potential of the animal, and its origination in the suckling animal is discussed from this standpoint. The role of vitamin D—a secondary one—is also discussed, and its probable mode of action, as previously defined, reaffirmed.

In connexion with the question of CaO and P_2O_5 absorption, emphasis is laid on gastric acidity and its neutralization by excess of CaCO_3 in the diet, together with the percentage composition of CaO and P_2O_5 in relation to the amount of food required to appease the animal's appetite. The ratio of CaO to P_2O_5 in the diet, as such, is considered to be of little importance in connexion with the absorption and utilization of these substances.

The economic waste that may arise from feeding "concentrates" and the hazards attributable to balancing the P_2O_5 in such cases with CaCO_3 are indicated. Observations on tetany and alkalosis in rachitic pigs with large quantities of calcium in the diet are recorded. The seasonal incidence of the disease in these animals is also dealt with.

It would seem permissible to regard osteomalacia as essentially a combination of aphosphorosis and acalcicosis due to deficiency of both phosphorus and calcium in the diet.

In dealing with the histological problems which have arisen in the course of discussion, I would like to acknowledge the great help received from Dawson's monograph (with Struthers), *Generalized Osteitis Fibrosa* (1923), and to record the fact that the specimens from the cases of early rickets were submitted to him for verification. Specimens illustrated on the Special Plate (except that from Pig 10) are from pigs with calcium-deficient diet of experiment 1930-1. The basal ration consisted of ground oats, barley and sharps, with meat, linseed, and alfalfa meals. It contained 0.21 per cent. CaO and 0.82 per cent. P_2O_5 . The photographs are from rib junctions cut fresh and treated with von Kossa's silver nitrate method.

Addendum.—Since this was written my attention has been called to a paper, *The Significance of Calcium and Phosphorus in the Development and Growth of Pigs*, by Reimers and Smuts (*Archiv f. Tiernahr. u. Tierzucht*, 1932, Ab. B. vii, 471). It is impossible to discuss it with advantage as, amongst other things, there is no description of the histopathology of the bones—a very grave omission.

BIBLIOGRAPHY

- Bacharach and Jephcott: *Journ. Biol. Chem.*, 1929, lxxxii, 751.
 Browning: *The Vitamins*, Pickett-Thomson Lab. Rep., 1931, i, 150.
 Dawson: *Edinburgh Med. Journ.*, 1923, xxx, 421.
 Elliot, Crichton, and Orr: *Brit. Journ. Exp. Path.*, 1922, iii, 10.
 Gamble, Ross, and Tisdall: *Amer. Journ. Dis. Child.*, 1923, xxv, 455, 470.
 Hess: *Rickets, Osteomalacia, and Tetany*, Kimpton, London, 1930.
 Hutya and Marek: *Spec. Path. and Therap. of Dis. Dom. Animals* (Mohler and Eichorn), vol. i, Baillière, Tindall and Cox, London, 1913.
 Korenchevsky: *The Aetiology and Pathology of Rickets*, Medical Research Council Report No. 71, 1922.
 McCollum and Simmonds: *The Newer Knowledge of Nutrition*, Macmillan and Co., New York, 1925.
 McGowan: *Journ. Path. and Bact.*, 1924, xxvii, 409.
 Idem: *Pernicious Anaemia, Leukaemia, and Aplastic Anaemia*, Lewis, London, 1926.
 Idem: *Biochem. Journ.*, 1933, xxvii, No. 3, 924, 943.
 McGowan, Cunningham, and Auchinachie: *Ibid.*, 1931, xxv, 1925.
 Mellanby: *British Medical Journal*, 1922, ii, 490; *Journ. Physiol.*, 1922, lii, Proc. ii.
 Idem: Medical Research Council Report No. 93, 1925.
 Park: *Physiol. Rev.*, 1923, iii, 106.
 Prazmowski: *Nutrit. Abs. and Rev.*, 1933, ii, 885.
 Schmorl: *Ergebn. d. inn. Med. u. Kinderheilk.*, 1909, iv, 403.
 Shohl and others: *Journ. Biol. Chem.*, 1931, xcii, 711.
 Theiler: *Denkschr. Schweiz. Naturf. Gesellsch.*, 1932, lxxviii, Abt. 1.
 Theiler and Green: *Nutrit. Abs. and Rev.*, 1932, i, 359.
 Wilson: *Lancet*, 1932, i, 1143.
 Zilva, Golding, and Drummond: *Biochem. Journ.*, 1924, xviii, 872.
 Zucker, Johnson, and Barnett: *Proc. Soc. Exp. Biol. and Med.*, 1922, xx, 20.
 Zucker and Matzner: *Ibid.*, 1923, xxi, 186.

* Other explanations of this phenomenon have been offered—for example, Berg (1923), *Vitamins* (Allen and Unwin), and Haldane, Hill and Luck (1923), *Journ. Physiol.*, lxxii, 361.

JOHN POOL McGOWAN: SOME OBSERVATIONS ON RICKETS

FIG. 1.—Fig 47. Basal ration: showing advanced osteoporosis, with fractures and disintegration.

FIG. 2.—Fig 162. Basal ration with sunlight: as in Fig. 1.

FIG. 3.—Fig 45. Basal ration with ultra-violet light: advanced osteoporosis.

FIG. 4.—Fig 10. Basal + CaCO_3 + cod-liver oil—normal rib junction: note that the line separating the cartilage from the dense calcified bony trabeculae is straight or slightly concave.

JOHN POOL McGOWAN: SOME OBSERVATIONS
ON RICKETS

FIG. 5.—Fig 44. Basal ration with cod-liver oil: advanced osteoporosis, with fractures at X.

A. WILSON GILL AND A. D. FRAZER:
SYPHILITIC SPONDYLITIS

X-ray of spine, showing increased density of third, fourth, and fifth lumbar vertebrae with flattening of bodies—osteoarthritic ridges joining articular margins.