Carcinoma of Parotid.—Six were operated upon without operative death. Two were alive and well after three years, and also at the end of five years.

Carcinoma of Stomach.—Out of 132 cases only eighteen were found suitable for operation. Eight died of the operation, two were alive at the end of three years and also at the end of five years.

Carcinoma of Colon.—Out of seventy-eight cases thirty-eight were submitted to operation, of whom eleven died. Sixteen were free from the disease at the end of three years, and thirteen at the end of five years.

Carcinoma of Rectum.—Out of 123 cases twenty-eight only were submitted to operation, of whom nine died. Five were alive at the end of three years and two at the end of five years.

Squamous Carcinoma of the Skin.—Sixteen were operated upon with no deaths. Nine were alive at the end of three years and also at the end of five years.

CONCLUSION

I have chosen these figures from the complete investigation as affording some evidence of the results of surgery (1) in the field in which radium is making definite claims—namely, the skin, mouth, and breast—and (2) in organs in which radium has so far claimed but little—the stomach, colon, and rectum. It is a striking fact that it is in the first group that surgery can show encouraging results, and in the latter it is so disappointing. If a much larger collection of surgical results were examined I believe that the same would hold good. Evidence is so far lacking that either radium or x rays are more than local remedies. It would appear that anatomical and pathological considerations impose much the same limitations upon their usefulness as they do upon that of surgery. The great advantages they offer over operative treatment, in the way of complete freedom from risk to life, avoidance of a period of invalidism, of mutilation, and of the infliction of pain, are in these days of heavy dosage, prolonged treatment, marked local reactions, and complementary surgical procedures, less convincing.

Radium and x rays are unquestionably of service as "palliative" agents in the more advanced cases of malignant disease in those organs in which surgical treatment of the disease when less advanced is most hopeful. How far they are producing lasting freedom from the disease which equals or exceeds that resulting from surgical methods alone, subsequent speakers in this discussion will doubtless endeavour to make plain.

THE RADIATION TREATMENT OF MALIGNANT DISEASE*

BY

OLIVER CHANCE, B.A., M.B., D.M.R.E.

DUBLIN

As the other two openers of this discussion are a surgeon and an x -ray therapist, I will, as far as possible, confine my remarks to radium therapy. At the outset it may be well to stress the fact that radium is not a substitute for either surgery or x rays in the treatment of malignant disease. All three methods play their parts, and a correct choice can be made between them only if the principles underlying each are understood.

A study of cancer statistics shows that in the treatment of many forms of malignant disease there is little difference between the results obtained by surgery, by x rays, and by radium when each is in the hands of an expert. This is really not surprising, for success in medicine is obtained by the skill and judgement of the individual

worker, and not by the media in which he works. On account of this, the treatment of cancer will never be standardized; for example, should a clinic be fortunate enough to possess an outstanding surgeon or x -ray therapist cases will naturally be referred to him which under ordinary circumstances would be treated with radium, and vice versa. Of course, when treatment is planned, material and equipment must be considered besides individual skill. Limitations in this direction often make it impossible to use an ideal technique.

Any consideration of radium therapy must begin with the all-important fact, so well demonstrated by Regaud and his fellow workers, that γ rays act on the dividing cell and not on the resting one. It would seem safe to assume from this that an actively mitotic tissue will be sterilized by a dose which will affect relatively stable tissue to a much lesser extent; it is on this assumption that modern radium therapy is based. While rapid cell division and malignancy are usually found in the same tumour, active mitosis does not necessarily connote malignancy. In general, the less complex the function of the cell and the more primitive the tissue from which it arises the shorter is its life, and consequently the higher its karyokinetic activity. Thus, in order to permit growth mitosis must be very active in a rodent ulcer, for it is composed of primitive basal cells which wear out rapidly. On the other hand, a tumour such as a gastric carcinoma is composed of complex cells which wear out slowly, so that, although as compared with a rodent ulcer the rate of cell division is usually low, the malignancy or rate of spread is very high. But we know that the effect of radium depends largely on the rate of cell division, and this explains why a rodent ulcer, which is relatively benign, is often more amenable to radium treatment than a malignant gastric carcinoma.

It is important to remember that γ rays merely sterilize, or partially sterilize, the tumour cells; they do not actually destroy the growth, but they do decrease the rate of cell production in it that the natural resistances of the body and the tissues of the tumour bed are able to effect its destruction. Bearing this in mind, it will be seen that a tumour is most easily destroyed by radium in healthy vascular tissue with good lymphatic drainage. To illustrate the importance of a good blood supply it may be stated that the application of radium to a growth involving bone should, if possible, be avoided, because the dose which sterilizes the malignant cells as a rule damages the blood supply of the bone, causing osteonecrosis.

Local sepsis and inflammation are unfavourable because they interfere with lymphatic drainage; hence radium should not be used to treat a cancer on a tongue which shows signs of severe chronic glossitis. Such tumours rarely respond well to irradiation, and other methods of treatment are preferable.

Another important consideration is the radio-sensitivity of the surrounding normal tissue. Such organs as the intestine and the liver are extremely sensitive to γ rays, and it is almost impossible to give an adequate dose to tumours arising in, or lying adjacent to, these organs without causing them permanent damage. Epidermis is particularly resistant, and even when destroyed by radium it regenerates, whereas other normal tissue becomes necrotic when heavily irradiated. The patient's general condition is also of great importance; to irradiate a patient who shows signs of general carcinomatosis or cachexia is merely to make matters worse.

The radium worker's difficulties do not end when he encounters a radio-sensitive tumour lying in healthy tissue. All parts of the growth must be given an adequate dose of γ rays, and at the same time permanent damage to surrounding healthy tissue must, as far as possible, be avoided. This is comparatively simple when the growth

* Read in opening a discussion at a combined meeting of the Sections of Radiology and Surgery at the Annual Meeting of the British Medical Association, Dublin, 1933.

is an accessible one, such as cancer of the skin, cervix uteri, mouth, or breast. In addition, ingenious operations have been devised by which laryngeal and rectal growths can be made accessible. In most other regions it is not possible to subject a tumour to an adequate homogeneous γ irradiation, unless perhaps in those few clinics which possess very large quantities of radium.

Radium may be placed against the surface of a growth, or needles containing it may be implanted into and around the tumour. Seeds are also used in this manner. In addition, in a few of the larger clinics teletherapy is available. In this method a large quantity of radium in the form of a bomb is held several centimetres from the body and used as a very powerful and inexhaustible x -ray tube. There is a fourth method of application—namely, cavitation, in which tubes containing radium are placed in the natural cavities of the body; but this is now little used, except in the treatment of cancer of the cervix uteri, when it is usually combined with surface therapy.

Implantation is the method which most concerns the surgeon, and the rest of radiotherapy (estimated by Forssell as about 75 per cent. of its entire scope) is of minor interest to him. It is peculiarly fitting that this method should be discussed in Dublin, for it was here that implantation originated. I refer to the work of Professor Joly and the late Dr. Stevenson, who were the first to introduce a tube of emanation into a metal needle. This marked one of the greatest advances in the treatment of cancer, and from it all modern implantation techniques have arisen. At present interstitial irradiation is a rather unscientific procedure, which is just as well carried out by a surgeon as by anyone else. Indeed, it can be argued that owing to his training a surgeon will insert needles with greater skill than a mere radiologist. It is likely, however, that as our knowledge of radio-physics and radiobiology increases, implantation will become just as exact a method of treatment as teletherapy or x -ray therapy, in which case only persons with special knowledge will be capable of using it properly. That these will be radiologists seems obvious. Radium is only implanted in a few sites, and it seems likely that a radiotherapist, working in a special clinic and seeing many more malignant cases than the average surgeon, will develop an excellent technique even if he does not possess special surgical knowledge. He will also know when implantation should not be used and when other methods of application or deep x -ray therapy are preferable. Hence we may conclude that radium therapy will be best developed if it is left in the hands of trained specialists who devote their whole time to it, and who have the necessary knowledge and training to keep abreast of the rapid advances which are taking place.

Poor results are frequently reported from the use of a technique which has been highly successful in other hands. This is because the result of treatment does not depend so much on the technique itself as on the intelligence with which it is employed. Not only must the treatment of each case be properly planned, but it must be carefully carried through with due regard to the reaction of the individual patient. In practice this is only possible in a properly organized and completely equipped clinic directed by an experienced radiotherapist.

The demand for a special radiotherapeutic clinic may seem extravagant, but if one pictures a surgeon who has to operate on patients sent from other departments without having the opportunity of first making a diagnosis or of following up the result of his surgical treatment, one has some idea of the difficulties of the radiologist who does not work in a special department. Radiotherapeutic technique is no easier than that of surgery, and the results obtained by it and the harm it may do are only apparent after a considerable time. Therefore, like any other specialist, the radiologist must follow up and study

his cases. Each patient must be examined at regular and frequent intervals, not only for the good of the individual, but also so that the value of the treatment may be assessed. To do this, statistics must be carefully prepared and should be in such a form that they are not misleading and can be compared easily with those from other clinics. The following up of patients depends on careful internal organization of the clinic, and close co-operation between it, the patient himself, his doctor, the staff of other hospitals, and the state medical authorities.

The foregoing remarks have been made much more interestingly many times before. Indeed, they may be described in Mr. Punch's words as "blinding glimpses of the obvious." But until they are recognized and acted upon, the obvious cannot too often be repeated.

THE RADIATION TREATMENT OF MALIGNANT DISEASE*

BY

WALTER M. LEVITT, M.B., M.R.C.P., D.M.R.E.

MEDICAL OFFICER IN CHARGE OF THE RADIOTHERAPEUTIC DEPARTMENT,
ST. BARTHOLOMEW'S HOSPITAL

The object of this discussion is, I suppose, to attempt to answer the question: When, and in what form, may radiations best be applied in the treatment of malignant disease? I propose, first of all, briefly to summarize the various methods of application of radiations, and then to consider the indications for their use.

METHODS OF APPLICATION OF RADIATIONS

Radiations are commonly used in the treatment of malignant disease as x rays or as the gamma rays of radium and radon: (a) by interstitial methods, the radium or radon being inserted into the tissues enclosed in needles or "seeds"; (b) by short-distance surface application, the radium being applied to the surface of the body on applicators or other apparatus moulded to the part; (c) intracavitary application, the radium being introduced in suitable containers into the natural body cavities; (d) by long-distance surface application, the so-called telecurie- or bomb-therapy, a large quantity of radium being massed in a single container, which is applied at a distance of several centimetres from the surface.

I would like to make it clear that wherever x rays are mentioned in this paper I refer to high-voltage, heavily filtered rays as used in modern deep x -ray therapy. While it is true that there are still one or two radiologists who cling to the use of soft, non-penetrating radiations, even in the treatment of malignant disease, agreement is practically universal that it is only by the use of the hardest radiations that the best results are to be obtained.

The broad principles of the use of all these methods are well known and do not need recapitulation; what concerns us here is the relative value of the methods in the treatment of cancer. If we compare these methods of application of radiations it will soon become apparent that their scope is very different. If we may make the reasonable assumption that it is essential in the irradiation of every malignant growth to apply an adequate dose of radiation throughout the whole extent of the growth, we find, for example, that the proportion of malignant tumours that can be adequately treated by interstitial methods is very limited, principally by anatomical considerations limiting the surgical access to the growth for the purpose of inserting the needles. Again, in the case of short-distance radiotherapy the depth at which the requisite dosage can be obtained is limited to a few centi-

* Read in opening a discussion at a combined meeting of the Sections of Radiology and Surgery at the Annual Meeting of the British Medical Association, Dublin, 1933.