improbable that the houses to be built will be let at less than 13s. a week, which will usually be beyond the reach of the poor.

3. There is no official definition of the word "slum," but dictionaries concur in giving two factors in the making of a slum: (1) the character of the building—the slum dwelling; and (2) the character of the occupant—the slum dweller. From the medical point of view, the second, or human, factor is probably of greater importance than the first.

4. Overcrowding—an aspect of the human factor—is one of the worst features of the slum problem. We shall not really solve the slum problem until we can deal effectually with overcrowding. It is a serious defect of the 1930 Housing Act that while it makes provision for dealing with the "individual" insanitary house, it makes no provision for dealing with "individual" cases of overcrowding. Overcrowding can only be dealt with under the Act when it occurs in a clearance or improvement area, and many bad cases of overcrowding occur outside such areas.

5. In any attack on the slum problem the claims of the children should come first. It is a waste of municipal resources to allow three-bedroomed municipal houses to be occupied by people without children.

6. People with large families of dependent children constitute the class whose housing need is greatest, but whose margin for rent is smallest. To enable this class to take proper advantage of municipal houses some system of differential renting is essential, preferably by means of temporary rent rebates based on an approved scale.

7. To compensate for the loss in rental consequent on these rebates, such municipal tenants as no longer really need to be subsidized should be called upon to pay the full economic rent.

VALLECULAR DYSPHAGIA

BY

JOHN E. G. MCGIBBON, M.B., B.S., D.L.O.

HONORARY ASSISTANT LARYNGOLOGIST, ROYAL SOUTHERN HOSPITAL,
LIVERPOOL

AND

J. H. MATHER, B.Sc., M.B., Ch.B., D.M.R.E.

HONORARY RADIOLOGIST, ROYAL SOUTHERN HOSPITAL, LIVERPOOL

(With Special Plate)

The elucidation of the causes of dysphagia arising in the lower pharynx has always presented difficulties. Following the perfection of the indirect laryngoscope during the past fifty years, gross, neoplastic and ulcerative lesions of the lower pharynx and upper oesophagus have been recognized. But there still remained a large number of undiagnosed cases which were termed "functional dysphagia" and "globus hystericus."

Within recent years, owing to the advances in radiological technique and the more universal use of peroral endoscopy, hitherto unrecognized causes of such dysphagia have been described—for example, spasm of the cricopharyngeus muscle, post-cricoid webs and strictures, exostoses of the cervical vertebrae, etc. Recently Dr. R. R. Woods¹ reported a case which he had called "vallecular dysphagia." His case was that of a man, aged 54, who gave a history of eighteen months' dysphagia. Investigation showed that, on swallowing, masticated food filled the patient's valleculae, so that his epiglottis was forced backwards against the posterior pharyngeal wall, impeding breathing and causing dysphagia. This report threw light on a number of unexplained cases of dysphagia which have been seen by the present writers during the past five years. These latter cases showed no lesion by direct or indirect examination of the pharynx, larynx, or oesophagus, but, on careful radiological examination, "hold-up" and abnormal passage of the opaque material in the lower pharynx was observed. These contradictory findings were explained

by Dr. Woods's report, and, although the present observers' cases exhibited less violent symptoms, they are convinced that their patients' complaints were caused by a similar mechanical abnormality, and that the condition described below under the title "primary vallecular dysphagia" is a definite entity of not infrequent occurrence, and one which may easily, but erroneously, be attributed to a "functional" disorder.

ANATOMY OF THE VALLECULAE

The valleculae are two fossae situated in the anterior aspect of the lower pharynx, between the pharyngeal portion of the tongue and the epiglottis (Fig. 1). They are bounded

FIG. 1.—Pharynx opened from behind. (Modified from Cunningham.)

anteriorly by the base of the tongue covered with lingual tonsil. The mucous membrane of this region of the tongue passes to the anterior surface of the upper free half of the epiglottis, forming the floors of the two depressions. The fossae are separated from each other by a prominent fold of mucous membrane—the glosso-epiglottic fold. Laterally they are bounded by less prominent ridges of mucous membrane—the pharyngo-epiglottic folds—which pass from the free lateral borders of the epiglottis to the lateral pharyngeal walls. Between each of the two layers of mucous membrane which constitute these three folds there is a small amount of elastic tissue. The floors of the valleculae lie immediately above and behind the hyoid bone, at the level of the third cervical vertebra (shown in Fig. 2), and sometimes the cornua of the

FIG. 2.

FIG. 3.

FIG. 4.

FIG. 2.—Diagram of normal pharynx (resting).

FIG. 3.—Diagram of Phases 1, 2, and 3 in deglutition. Nasopharynx shut off by Passavant's ridge. Mouth closed—larynx raised and closed—pharyngeal cavity obliterated—epiglottis and vallecula compressed.

FIG. 4.—Diagram of Phases 4, 5, 6, and 7 in deglutition. Nasopharynx, mouth, and larynx closed—negative pressure in pharyngeal cavity—larynx dropped. Laryngo-pharynx remains high up—food sucked into pharynx and oesophagus not touching epiglottis.

hyoid bone are visible through their mucous membrane. Indirect laryngoscopy shows that in a small number of cases the fossae may be partially or completely hidden by the base of the tongue, and that they vary very much in depth in different individuals. Their mucous membrane is less red in colour than that of the surrounding structures, and it may be tightly stretched, crinkled, or even raised by large veins into prominent ridges. The valleculae are developed in the third cleft, between the field formed by the fused ventral ends of the second and third arches in front, and the central mass of the central end of the fourth arch behind. Their sensory nerve supply is derived from the third and fourth arches—that is, the glosso-pharyngeal and the superior laryngeal nerves. The writers have been unable to find any record of developmental abnormality occurring in this situation.

NORMAL MECHANISM OF DEGLUTITION.

In order to appreciate any abnormalities of swallowing, it is necessary to consider the known normal process. Deglutition is a complicated reflex act, the afferents being the second division of the fifth, the ninth, and the pharyngeal branches of the superior laryngeal nerves. The efferents are the third division of the fifth, the seventh, ninth, tenth, eleventh, and twelfth cranial nerves. The centre is situated in the medulla oblongata, and may be considered as a chain of associated centres, stimulation of one of which fires off the others in orderly sequence (Starling).

Descriptions of the mechanism of swallowing, which were formulated by physiologists, have become greatly modified by accurate radiological investigation of the process, notably by the researches of A. E. Barclay.² Similar though less detailed studies have been made by Mosher³ and Pancoast.⁴ Barclay's findings have been summed up,⁵ and the phases of swallowing are described as follows:

1. The mouth and nose are closed.
2. The larynx is raised and closed and its upper part is obliterated by the back of the tongue.
3. The pharyngeal space is obliterated by (a) raising the larynx, (b) retraction of the tongue. (The posterior wall of the pharynx does not come forward: the pharynx is not constricted, but is compressed from before backwards against the posterior wall—that is, against the vertebra.)
4. The pharynx is opened up, and with the three outlets closed a negative pressure is obtained by (a) dropping the larynx, and (b) allowing the tongue to go forward.
5. During this movement the laryngeal pharynx still remains high up, in close relation to the epiglottis. It is in an opened-mouth condition.
6. Food is tipped over the back of the tongue.
7. Food is sucked into the open mouth of the laryngeal pharynx, which, as it receives the bolus, drops from the epiglottis down to its normal position, thus opening up the larynx again after food has passed.
8. The food is probably sucked some distance down the oesophagus.

During the entire act the epiglottis remains upright; in the third phase it is pressed between the base of the tongue and the posterior pharyngeal wall, and in the sixth phase ordinary food passes over without touching it (Figs. 3 and 4). These observations have proved conclusively that the epiglottis does not fold backwards, as a trap-door, over the laryngeal orifice during swallowing—although this is still figured in physiological textbooks—and they have altered entirely the views previously held concerning deglutition. The older explanation, that the bolus of food is seized by the pharyngeal constrictors and forced into the oesophagus, where it is passed on by peristalsis, is disproved. As long ago as 1911 Lexer,⁶ while watching a case on which he had performed a Wullstein operation for carcinoma of the oesophagus, made the important observation that the absence of peristalsis in the skin tube did not have an adverse effect on the ingestion of food. Saint,⁷ commenting on this, concluded that the descent of food through the oesophagus is "due to the impetus given to it by the pharyngeal muscles." More recently Sampson,⁸ who performed a similar operation on a case of simple oesophageal stricture, showed that the force of gravity is not necessary for swallowing through a skin tube oesophagus, and he suggested that the passage of food through such a tube may be accomplished by air pressure (swallowing air). Evans,⁹ who used an extrathoracic rubber oesophagus following excision of a carcinomatous gullet, emphasized the fact that "the propulsive force of the mouth suffices to ensure the conveyance of food to the stomach through a patent inactive tube." It may therefore be accepted that the passage of food through the lower pharynx and upper oesophagus is accomplished by a combination of propulsion and negative pressure. One statement made in the summary of Barclay's observations is in question—namely, that the posterior pharyngeal wall does not come forward. Brown¹⁰ has drawn attention to the fact that a small ruck—Passavant's ridge—is formed in the upper part of the superior pharyngeal constrictor, and that this ruck *does* come forward, to form the posterior half of the nasopharyngeal sphincter, for closure of the nasopharynx during deglutition (Figs. 3 and 4).

RADIOLOGICAL EXAMINATION OF THE ACT

The writers have made careful screen examinations of the act of swallowing in more than 100 normal males and females. The usual bread-and-milk opaque meal was used. In approximately 50 per cent. of the cases examined the bolus of food was swallowed, leaving only a trace in the mouth and pharynx, none being seen to enter the valleculae. In the other 50 per cent. a little opaque material entered the valleculae, but was evacuated almost immediately by compression of the valleculae during succeeding acts of swallowing. Apparently, therefore, in normal cases a little food may be retained for a very short time in the valleculae. In these cases the valleculae are demonstrated in the lateral and oblique positions as a small semilunar shadow, and in the "standing facing" position as two small semilunar shadows, one on each side of the mid-line at the level of the disk, between the third and fourth cervical vertebrae.

VALLECULAR DYSPHAGIA: PRIMARY AND SECONDARY

Symptoms and Aetiology

The writers have observed fourteen cases of what they term "primary vallecular dysphagia," and a very considerable number of cases which they classify as "secondary vallecular dysphagia."

Primary vallecular dysphagia is clinically important. Of the fourteen cases seen, five were females and nine males. Their ages ranged from 35 years to 79 years. These patients all came to the throat department complaining of discomfort on and after swallowing, which they referred to the hyoid region, and which had been present for periods extending from fourteen days in the shortest up to "about two years" in the longest: in some cases the discomfort was intermittent, and occasionally persisted for some long time after eating. The nature of onset varied—in some cases it was insidious, in others sudden, and with no known cause; in three cases the symptoms were ascribed to swallowing a bone, which the patients imagined had become lodged in their throats. The above-mentioned three patients stated that they felt as though they had a foreign body in their throats; the remainder were obviously apprehensive and suffering from carcinophobia. None of them complained of dyspnoea or such marked symptoms as occurred in Dr. Woods's case. One man, aged 35 years, complained of excessive salivation. This was the only case in which masticated food lying in one vallecula was actually seen by indirect laryngoscopy. Indirect laryngoscopy in the remainder of cases revealed no local lesion, and these negative findings were confirmed later in all cases by direct examination.

X-ray investigation showed that in cases of primary vallecular dysphagia, when watched on the screen, there was a definite inability to swallow the bolus of opaque food. When told to "swallow," the patients made repeated but ineffectual attempts to pass the bolus of food from the mouth into the pharynx. At each attempt a little food passed (1) into the valleculae, where it remained, (2) into the right and left pyriform fossae, and (3) into the pharynx and oesophagus. The pharynx was dilated and food adhered to its walls. Gradually, and after repeated "swallows," the patients succeeded in emptying their mouths. The paste was then slowly evacuated from the pharynx, but was retained in the distended valleculae for as long as forty-five minutes after the first attempts to swallow.

Figs. 1 and 2 (on Special Plate) clearly demonstrate the above points; they are oblique and antero-posterior views of the pharynx and upper oesophagus of a typical case of primary vallecular dysphagia, taken after the patient had succeeded in emptying his mouth.

General clinical examination was negative for any discoverable systemic disease, and some of these cases have been kept under observation for five years without manifesting any lesion which might have been latent at the time of their first examination.

In what we have termed *secondary vallecular dysphagia* the valleculae were filled in a similar manner, but this abnormality was due to a manifest lesion somewhere within the reflex arc, or in the pharynx or oesophagus themselves. In these cases the presenting symptoms were those of the primary lesion, and the vallecular filling and retention was discovered only on routine examination, the x-ray findings being similar to those occurring in the primary form of dysphagia, but less marked. Thus we have seen secondary vallecular dysphagia in patients suffering from carcinoma of the upper and middle thirds of the oesophagus, post-cricoid carcinoma, and pharyngeal pouch. Mosher published a plate showing a similar condition in a case of bulbar paralysis, although he did not apply this title to the condition; and vallecular filling and retention have been described by Assmann¹¹ in bilateral recurrent nerve paralysis secondary to a mediastinal tumour. A similar condition was observed by him in a case of tabes with paralysis of the vagus, and in a case in which botulism had injured the vagus nerve. Reiche¹² has reported retention of food in the valleculae following diphtheritic paralysis, and Weingaertner¹³ and Desseker¹⁴ have each described a similar retention in cases of carcinoma of the upper part of the oesophagus with paralysis of the vagus nerve. In these latter cases the vallecular contents overflowed into the trachea and bronchi. A similar case has been observed by the present writers.

From our observations it would appear that secondary vallecular dysphagia occurs only with lesions of relatively rapid evolution. In non-progressive or slowly advancing lesions which have been present for many years the neuro-muscular apparatus has apparently accustomed itself to its abnormal conditions and has overcome the mechanical difficulties present. Thus we have found it absent in post-cricoid spasm, webs, and strictures; in congenital oesophageal stenosis, traction pouch of the oesophagus, etc.—all conditions of lengthy duration. It has been present in pharyngeal pouch, but such cases come for advice only at the onset of troublesome dysphagia—that is, when there is a relatively rapid exacerbation of their symptoms.

PATHOLOGY

The pathology of primary vallecular dysphagia is speculative only, but the known pathology of the secondary form suggests that there must be somewhere a failure in the neuro-muscular mechanism of swallowing. Ledoux,¹⁵ in a study of the mechanics of deglutition, has classified the elevators of the larynx into two functional groups— anterior and posterior. The anterior group consists of the mylo-hyoid, the genio-hyoid, the posterior fibres of the genio-glossus, and the anterior belly of the digastric—of these the mylo-hyoid is functionally the most important. This group of muscles pulls the larynx upwards and forwards behind the base of the tongue, and also pulls the larynx forwards away from the posterior pharyngeal wall to open up the hypopharynx. It is apparent that, with impairment of this *forward* movement of the larynx, the valleculae can readily fill, and are not emptied by compression. Ledoux points out that in bulbar paralysis this anterior group of muscles is invariably the first to be affected; and Critchley and Kubik¹⁶ have drawn attention to the constant atrophic degeneration found microscopically in the mylo-hyoid muscle, which occurs in cases of bulbar paralysis in addition to the characteristic changes seen in the central nervous system. It may be, therefore, that there is

present some neuro-muscular inferiority of this group of muscles which will permit easily of its partial or complete dysfunction from non-obvious causes, and so give rise to vallecular filling and retention.

Goldmark and Scholz¹⁷ have reported two cases which they termed “diverticulum of the epiglottic valleculae.” A male, aged 60 years, and a female, aged 62 years, both exhibited filling and retention in the valleculae of barium paste. These two cases were characterized by discomfort on swallowing, occasional regurgitation of foul material, swelling in the submaxillary region, and local tenderness, with no laryngologic or systemic changes. They would be classified by the present writers as primary vallecular dysphagia. Goldmark and Scholz, basing their opinion on the erroneous theory that the epiglottis bends backwards during the passage of food through the pharynx, consider the pathology of the condition to be due to loss of elasticity of the epiglottis, whereby this epiglottic “bending-back” cannot take place. This theory is untenable. Holz knecht and Olbert,¹⁸ who found vallecular retention present in certain neuropathic persons, demonstrated an unusual relaxation of the entire oesophagus in such cases. They explain the condition as due to atony of the muscles of deglutition; and Paluguay¹⁹ has shown that atony may cause marked delay in any part of the path of deglutition. The marked dilatation of the lower pharynx shown radiologically in the present writers' cases of primary vallecular dysphagia would suggest atony of the middle and inferior pharyngeal constrictor muscles.

It has been shown almost conclusively that after food is propelled into the pharynx it is sucked down into the oesophagus by negative pressure. If the valleculae become filled owing to lack of compression the negative pressure in the pharynx is then insufficient to empty them. Barclay has instanced a similar difficulty experienced in dislodging mucus from the posterior pharyngeal wall by the action of suction alone. This leads to incompleteness of the act of swallowing. Although deglutition is commonly described as occurring in stages, it is one act only, and normally its initiation by stimulation of the fauces inevitably leads to completion. Incomplete deglutition will give rise to abnormal pharyngeal sensations, which are interpreted in various manners by different individuals.

DIAGNOSIS

Diagnosis can only be made by careful local and general examination, together with thorough radiological investigation. It is obvious from what has been said above that either of these examinations alone is valueless. In cases of primary vallecular dysphagia laryngoscopic and endoscopic examinations are negative, and the patient's symptoms are apt to be attributed to functional dysphagia. X-ray findings are inferential only without local and systemic investigations.

TREATMENT

In thirteen out of fourteen of the writers' cases of primary vallecular dysphagia troublesome symptoms disappeared after explanation and reassurance. One case was heard of occasionally going round from hospital to hospital for advice. Eventually all trace of him was lost. Goldmark and Scholz advise frequent gargling with plain water after meals, and one of their patients found alleviation by repeated drinking of small amounts of water after meals. It appears to the present writers that the relief derived from the above two measures can be attributed to the repeated movements of deglutition, whereby the valleculae would be “milked” of their abnormal contents. During the discussions which followed Dr. Wood's paper he stated that he had been attempting relief of the patient's symptoms by diathermy puncture of the

valleculae, and by section of the lateral pharyngo-epiglottic folds. Removal of the free portion of the epiglottis was suggested. This latter appears to be the rational treatment if the symptoms are sufficiently severe. The epiglottis in man is a vestigial structure, and it is well known that its removal, either by surgery or by disease, is devoid of any harmful after-effects, provided always that it is the free portion only that is removed.

In the secondary forms of vallecular dysphagia treatment naturally would be directed towards the primary lesion.

SUMMARY

1. Dysphagia, caused by filling and retention of food in the valleculae, is described.

2. This type of dysphagia may be a distinct entity of clinical importance—of unknown pathology—and present as primary vallecular dysphagia.

3. Alternatively, it may occur as a secondary phenomenon due to some manifest disturbance of the neuromuscular mechanism of swallowing, or to local disease in the pharynx or oesophagus. These cases are termed "secondary vallecular dysphagia," and they are not clinically important, as their symptoms are overshadowed by the existence of a more grave lesion.

4. Treatment of primary vallecular dysphagia has in most cases resolved itself into explanation and reassurance, with disappearance of symptoms. In more severe

cases, however, it might be necessary to amputate the free portion of the epiglottis.

5. Treatment of the secondary forms would be dictated by the nature of the causative lesion.

Our thanks are due to Dr. Walker and the staff of the x-ray department of the Royal Southern Hospital for their help in the examination of the cases on which this paper is based.

REFERENCES

- ¹ Woods, R. R.: *Lancet*, 1931, i, 1137.
- ² Barclay, A. E.: *Brit. Journ. Radiol.*, 1930, iii, 534.
- ³ Mosher, H. P.: *Laryngoscope*, 1927, xxxvii, 235.
- ⁴ Pancoast, H. K.: *Journ. Amer. Med. Assoc.*, 1930, xcvi, 1318.
- ⁵ Summary of conjoint meeting of the Sections of Radiology and Oto-Laryngology of the Royal Society of Medicine, *Lancet*, 1930, ii, 694.
- ⁶ Lexer, E.: *Münch. med. Woch.*, 1911, lviii, 1548.
- ⁷ Saint, J. H.: *Arch. of Surg.*, 1929, xix, 75.
- ⁸ Sampson, H. H.: *Brit. Journ. Surg.*, 1932-3, xx, 447.
- ⁹ Evans, Arthur: *Ibid.*, 1932-3, xx, 388.
- ¹⁰ Browne, Dennis: *Ibid.*, 1932-3, xx, 7.
- ¹¹ Assman, H.: *Die klinische Roentgendagnostik*, Leipzig, 1929, p. 210.
- ¹² Reiche, F.: *Fortschr. a. d. Geb. d. Roentgenstr.*, 1918, xxv, 353.
- ¹³ Weingaertner, M.: *Ibid.*, 1915, xxii, 397.
- ¹⁴ Desseker, C.: *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, 1924, xxxvii, 41.
- ¹⁵ Ledoux, D.: *Proc. Coll. Oto-Laryngol.*, 1929; abstract, *Journ. Laryngol. and Otol.*, 1931, xv, 899.
- ¹⁶ Critchley, M., and Kubik, C. S.: *Brain*, 1925, xlviii, 492.
- ¹⁷ Goldmark, C., and Scholz, T.: *Journ. Amer. Med. Assoc.*, 1932, xcvi, 1972.
- ¹⁸ Holzknecht, G., and Olbert, D.: *Zeit. f. klin. Med.*, 1910, lxxi, 91.
- ¹⁹ Paluguay, J.: *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, 1924, xxxvii, 107.

CERTAIN INTRACRANIAL TUMOURS THEIR VARIABILITY AND MULTIPLICITY*

BY

W. E. CARNEGIE DICKSON, M.D., B.Sc.,
F.R.C.P.Ed.

(From the Pathological Department, West End Hospital for
Nervous Diseases, London)

(With Special Plate)

An enormous amount of extremely valuable work on cerebral tumours has been published by Harvey Cushing, Percival Bailey, Wilder Penfield, Carmichael, and numerous other writers, not only within the past few years, but in the past few months, and even weeks. The classification of gliomas has especially been the subject of much recent research; and, unless one is prepared to offer some *simplification* of their present very elaborate classification, one had, in the meantime, better leave that subject alone.

Some twenty-three years ago, along with my colleague at the Royal Edinburgh Hospital for Sick Children, the late Dr. J. Stewart Fowler, I described one of the earlier cases of tuberous (tuberos) sclerosis,¹ in which, in addition to the typical cortical and ventricular nodules in the brain, there was a rhabdomyoma of the heart. We noted that: "The condition is characterized by the presence of congenital 'dislocation' tumours in all or any of the following sites: brain, kidney, heart, skin—more rarely, pancreas, duodenum, thyroid, breast. The condition in the brain is established probably at, or soon after, the seventh month of foetal life"—perhaps considerably earlier. Since investigating this case (the staining methods at our disposal were then very far behind those in modern use) cases of "neoplastic syndromes" have always appeared to me as being of especial interest and importance.

* Read, with naked-eye, microscopical, and lantern demonstration, in the Section of Pathology and Bacteriology at the Annual Meeting of the British Medical Association, Dublin, 1933. As it was not possible to reproduce all the naked-eye and microscopical photographs used to illustrate the points brought out in this paper a few of the latter only have been reproduced in miniature in the accompanying special plate, more as a general key-plan than to illustrate special details.

Lindau's disease is another good example of such multiple tumour syndrome, there being often, in addition to the specially characteristic haemangioblastomatous cyst of the cerebellum, an angiomas of the retina, and perhaps cavernous angioma of the liver, and cystic disease of the kidneys, pancreas, etc.⁵

In a recent case of von Recklinghausen's disease, under the care of my colleague Dr. C. Worster-Drought, the multiplicity and variety of the tumours found post mortem were so extraordinary that I thought it would make an excellent "text" for my present paper, especially if taken along with another case of haemangioma of the cerebral cortex, in which other and unsuspected tumours were also found—for example, a myxochondroadenoma invading the base of the skull, a pituitary cyst, etc.

HISTOLOGICAL VARIATIONS IN INDIVIDUAL TUMOURS

Before describing these cases of multiple tumour syndrome let me allude briefly to another aspect of the subject which may have some bearing upon it—namely, the wide histological variations which may be found in individual tumours, whether gliomatous or meningiomatous, etc. Variations and intermediate types occurring within the glioma group are too complicated for a short paper, but the study of one or two typical examples of the meningioma group may be helpful; and, for this purpose, I show you some specimens, naked-eye and microscopical, from a typical "dural" and a couple of typical "leptomeningeal meningiomas."* The dural tumour was one successfully removed by Mr. C. P. Wakeley from a male patient of Dr. Worster-Drought's, aged 39, and its chief interest for us lies in what one may term the "polyvalency" of its cells. In some parts the tumour cells show imperfect division, and tend to be arranged in groups and to reproduce the structure of Pacchionian bodies or arachnoidal villi. In others they tend to be arranged in "whorls," usually around a small blood vessel, and there is a

* Though often described as dural, these meningiomas probably arise from or in connexion with the leptomeninges—for example, from the arachnoidal villi.

JOHN E. G. MCGIBBON: VALLECULAR DYSPHAGIA

FIG. 1.—Oblique view of pharynx and upper oesophagus, showing both valleculae distended with opaque food, and dilatation of pharynx.

FIG. 2.—Antero-posterior view of same patient.

BRENNAN S. CRAN: FIBROSARCOMA OF THE SUPRARENAL GLAND

FIG. 1.—Fibrosarcoma of the suprarenal. (Spleen above, left kidney below.)

FIG. 2.—Microscopical structure. (Haemalum and eosin $\times 210$.)