valleculae, and by section of the lateral pharyngo-epiglottic folds. Removal of the free portion of the epiglottis was suggested. This latter appears to be the rational treatment if the symptoms are sufficiently severe. The epiglottis in man is a vestigial structure, and it is well known that its removal, either by surgery or by disease, is devoid of any harmful after-effects, provided always that it is the free portion only that is removed.

In the secondary forms of vallecular dysphagia treatment naturally would be directed towards the primary lesion.

SUMMARY

1. Dysphagia, caused by filling and retention of food in the valleculae, is described.

2. This type of dysphagia may be a distinct entity of clinical importance—of unknown pathology—and present as primary vallecular dysphagia.

3. Alternatively, it may occur as a secondary phenomenon due to some manifest disturbance of the neuromuscular mechanism of swallowing, or to local disease in the pharynx or oesophagus. These cases are termed "secondary vallecular dysphagia," and they are not clinically important, as their symptoms are overshadowed by the existence of a more grave lesion.

4. Treatment of primary vallecular dysphagia has in most cases resolved itself into explanation and reassurance, with disappearance of symptoms. In more severe

cases, however, it might be necessary to amputate the free portion of the epiglottis.

5. Treatment of the secondary forms would be dictated by the nature of the causative lesion.

Our thanks are due to Dr. Walker and the staff of the x-ray department of the Royal Southern Hospital for their help in the examination of the cases on which this paper is based.

REFERENCES

- ¹ Woods, R. R.: *Lancet*, 1931, i, 1137.
- ² Barclay, A. E.: *Brit. Journ. Radiol.*, 1930, iii, 534.
- ³ Mosher, H. P.: *Laryngoscope*, 1927, xxxvii, 235.
- ⁴ Pancoast, H. K.: *Journ. Amer. Med. Assoc.*, 1930, xcvi, 1318.
- ⁵ Summary of conjoint meeting of the Sections of Radiology and Oto-Laryngology of the Royal Society of Medicine, *Lancet*, 1930, ii, 694.
- ⁶ Lexer, E.: *Münch. med. Woch.*, 1911, lviii, 1548.
- ⁷ Saint, J. H.: *Arch. of Surg.*, 1929, xix, 75.
- ⁸ Sampson, H. H.: *Brit. Journ. Surg.*, 1932-3, xx, 447.
- ⁹ Evans, Arthur: *Ibid.*, 1932-3, xx, 388.
- ¹⁰ Browne, Dennis: *Ibid.*, 1932-3, xx, 7.
- ¹¹ Assman, H.: *Die klinische Roentgendagnostik*, Leipzig, 1929, p. 210.
- ¹² Reiche, F.: *Fortschr. a. d. Geb. d. Roentgenstr.*, 1918, xxv, 353.
- ¹³ Weingaertner, M.: *Ibid.*, 1915, xxii, 397.
- ¹⁴ Desseker, C.: *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, 1924, xxxvii, 41.
- ¹⁵ Ledoux, D.: *Proc. Coll. Oto-Laryngol.*, 1929; abstract, *Journ. Laryngol. and Otol.*, 1931, xv, 899.
- ¹⁶ Critchley, M., and Kubik, C. S.: *Brain*, 1925, xlviii, 492.
- ¹⁷ Goldmark, C., and Scholz, T.: *Journ. Amer. Med. Assoc.*, 1932, xcvi, 1972.
- ¹⁸ Holzknecht, G., and Olbert, D.: *Zeit. f. klin. Med.*, 1910, lxxi, 91.
- ¹⁹ Palugay, J.: *Mitt. a. d. Grenzgeb. d. Med. u. Chir.*, 1924, xxxvii, 107.

CERTAIN INTRACRANIAL TUMOURS THEIR VARIABILITY AND MULTIPLICITY*

BY

W. E. CARNEGIE DICKSON, M.D., B.Sc.,
F.R.C.P.Ed.

(From the Pathological Department, West End Hospital for
Nervous Diseases, London)

(With Special Plate)

An enormous amount of extremely valuable work on cerebral tumours has been published by Harvey Cushing, Percival Bailey, Wilder Penfield, Carmichael, and numerous other writers, not only within the past few years, but in the past few months, and even weeks. The classification of gliomas has especially been the subject of much recent research; and, unless one is prepared to offer some *simplification* of their present very elaborate classification, one had, in the meantime, better leave that subject alone.

Some twenty-three years ago, along with my colleague at the Royal Edinburgh Hospital for Sick Children, the late Dr. J. Stewart Fowler, I described one of the earlier cases of tuberous (tuberos) sclerosis,¹ in which, in addition to the typical cortical and ventricular nodules in the brain, there was a rhabdomyoma of the heart. We noted that: "The condition is characterized by the presence of congenital 'dislocation' tumours in all or any of the following sites: brain, kidney, heart, skin—more rarely, pancreas, duodenum, thyroid, breast. The condition in the brain is established probably at, or soon after, the seventh month of foetal life"—perhaps considerably earlier. Since investigating this case (the staining methods at our disposal were then very far behind those in modern use) cases of "neoplastic syndromes" have always appeared to me as being of especial interest and importance.

* Read, with naked-eye, microscopical, and lantern demonstration, in the Section of Pathology and Bacteriology at the Annual Meeting of the British Medical Association, Dublin, 1933. As it was not possible to reproduce all the naked-eye and microscopical photographs used to illustrate the points brought out in this paper a few of the latter only have been reproduced in miniature in the accompanying special plate, more as a general key-plan than to illustrate special details.

Lindau's disease is another good example of such multiple tumour syndrome, there being often, in addition to the specially characteristic haemangioblastomatous cyst of the cerebellum, an angiomas of the retina, and perhaps cavernous angioma of the liver, and cystic disease of the kidneys, pancreas, etc.⁵

In a recent case of von Recklinghausen's disease, under the care of my colleague Dr. C. Worster-Drought, the multiplicity and variety of the tumours found post mortem were so extraordinary that I thought it would make an excellent "text" for my present paper, especially if taken along with another case of haemangioma of the cerebral cortex, in which other and unsuspected tumours were also found—for example, a myxochondroadenoma invading the base of the skull, a pituitary cyst, etc.

HISTOLOGICAL VARIATIONS IN INDIVIDUAL TUMOURS

Before describing these cases of multiple tumour syndrome let me allude briefly to another aspect of the subject which may have some bearing upon it—namely, the wide histological variations which may be found in individual tumours, whether gliomatous or meningiomatous, etc. Variations and intermediate types occurring within the glioma group are too complicated for a short paper, but the study of one or two typical examples of the meningioma group may be helpful; and, for this purpose, I show you some specimens, naked-eye and microscopical, from a typical "dural" and a couple of typical "leptomeningeal meningiomas."* The dural tumour was one successfully removed by Mr. C. P. Wakeley from a male patient of Dr. Worster-Drought's, aged 39, and its chief interest for us lies in what one may term the "polyvalency" of its cells. In some parts the tumour cells show imperfect division, and tend to be arranged in groups and to reproduce the structure of Pacchionian bodies or arachnoidal villi. In others they tend to be arranged in "whorls," usually around a small blood vessel, and there is a

* Though often described as dural, these meningiomas probably arise from or in connexion with the leptomeninges—for example, from the arachnoidal villi.

marked tendency to hyaline degeneration and, later, calcification, with the formation of the so-called "brain-sand particles" or "psammomatous bodies" (Plate, Fig. 13). In other parts again, and especially in areas of degeneration and haemorrhage, the tumour cells have a marked tendency to form blood vessels, either capillary, with definite walls containing reticulin, or thin-walled cavernous spaces, sometimes showing little more than an endothelial lining formed directly by the tumour cells themselves (Fig. 14). All stages of this angioblastic process can be followed in the specimens from this case, but they can perhaps be even better illustrated from a leptomeningioma found post mortem in the right parietal region of a male patient of Dr. Macnamara's, aged 49. The naked-eye appearance of the cut surface of this tumour was very variegated, with patches of older and more recent haemorrhage and organizing blood clot, as well as soft tumour tissue, in some parts rather necrotic-looking. Here again all variations may be seen. There are actively growing, highly cellular, meningotheliomatous areas, many of the cells showing active mitotic division (Fig. 15). In other areas the formation can be observed of capillaries in very large numbers directly from the tumour cells themselves, vascularizing, as it were, any areas of haemorrhage, or even of degeneration, in the tumour, and closely simulating the appearance of organization of, say, a wound or a pleural or pericardial exudate (Fig. 16). One might quite well be justified in applying the term "capillary angioma" to these, but in other areas the vascular spaces are so thin-walled and dilated that they are "cavernous" in type. Or again, one finds the formation of "whorls" of tumour cells around the newly formed capillaries, and in some there is well-marked calcification and the formation of numerous psammomatous bodies, some of them coalescing to form rosette-like and other bizarre "compound" psammomata. As such leptomeningeal, in comparison with the so-called "dural," tumours are somewhat uncommon, I have put out in the museum a second specimen from a female patient, aged 64, in whose right occipital lobe a large leptomeningioma was found post mortem, in its naked-eye appearance very much simulating a glioma. Somewhat analogous meningiomatous (endotheliomatous) tumours showing varying admixtures of meningothelial, vascular, and other elements may also be found in connexion with the spinal cord.⁶

Having briefly studied what we have called the "polyvalency" of tumour cells in individual tumours of, especially, the meningiomatous group, let us now turn to the subject of multiple neoplastic syndromes.

MULTIPLE NEOPLASTIC SYNDROMES

I have already alluded to the condition known as tuberous sclerosis, now sometimes termed "epiloia." Closely allied with this is the syndrome usually known as von Recklinghausen's disease or multiple neurofibromatosis, and for the number and variety of tumours present the case which I have exhibited in the museum is perhaps unique, although somewhat similar cases have been described by Penfield and Young,² Hosoi,³ and Kernohan and Parker.⁴ As Dr. Worster-Drought, Dr. McMenemey, and I hope to publish this case in considerable detail presently I shall merely demonstrate to you the specimens and pictures of them and enumerate the extraordinary variety of tumours present. There are, for example, not only two large bilateral acoustic nerve tumours, compressing and displacing the pons, medulla, and cerebellum—so-called "cerebello-pontine angle tumours"—and showing a structure somewhat intermediate between a typical neurofibroma and a typical

meningioma (Fig. 1), but there are also innumerable other, though much smaller, tumours on practically all the cranial nerves. There are also several dozen—some fifty or so—meningiomas and meningiomatous plaques of various sizes, projecting inwards from the inner surface of the dura, the larger tumours, up to about the size of a pea or small bean, projecting into and indenting, but not adherent to, the subjacent surface of the hemispheres. These did not show any growth outwards through the dura and skull, except where a considerable number of small meningiomas, evidently originating from Pacchionian bodies or arachnoidal villi, were found projecting into the lumen of the sagittal sinus; and in the cribriform region the olfactory grooves were filled with a mass of slightly nodulated to cauliflower-like meningiomatous growth, invading the ethmoid sinuses and right orbit. There were also innumerable small tumours on practically all the posterior nerve roots of the spinal cord (Fig. 4), and especially on those of the cauda equina. There were also, in addition to these, numerous intramedullary tumours, some of them similar to the nerve-root tumours, but others again apparently fibrillary astrocytomas; whilst at many levels of the cord there were smaller or larger central ependymal proliferations and tumours (Fig. 7), as well as gliosis and the formation of syringomyelia-like cavities. There was a previous history of a single, subcutaneous neurofibroma having been removed from the left shoulder at another hospital, but no other subcutaneous tumours were found at our post-mortem, nor any naked-eye tumours on the subcutaneous nerves of the body and extremities nor on the main nerve trunks and their branches in the upper and lower limbs, or elsewhere; though a minute microscopical tumour, resembling those found in the posterior nerve roots, was found on the trunk of one of the sciatic nerves (Fig. 9). Tumours are also probably present in some of the sympathetic ganglia (Fig. 8) and in the suprarenal medulla, but these have not yet been fully worked out.

For comparison and contrast with this case of multiple neurofibromatosis, I have placed in the museum two cases of single cerebello-pontine-angle tumours from female patients, aged 33 and 32 respectively, the former being apparently a fibroblastoma definitely malignant and locally invasive, supervening upon a neurofibroma of the acoustic nerve (in this case there were also numerous enlarged, almost tumour-like, Pacchionian bodies, and a cavernous angioma of the liver), the second case being apparently not a neurofibroma but a meningioma, compressing the acoustic nerve and neighbouring structures. It also has become malignant and infiltrative.

In contrast to the foregoing, and for comparison with cases of Lindau's disease,⁵ the following case of multiple tumours is of interest. The case was that of a woman, aged 42, a patient of Dr. Yealland and Mr. Wakeley's. She gave an eleven years' history pointing to a right motor cortical lesion. A fronto-parietal decompression operation was performed, but twenty-four hours afterwards hyperpyrexia supervened and the patient died a week later. At the post-mortem this was accounted for by the rupture of an unsuspected suppurating pituitary cyst and a terminal acute streptococcal meningitis, curiously localized to the cerebral hemisphere *opposite* to the side decompressed. The right side of the brain had been sealed off and protected from immediate infection by a thin layer of blood clot, following upon the decompression. A cavernous angioma (Fig. 10) of considerable size was found towards the posterior part of the right parietal lobe, only about half an inch from the edge of the decompression window, and this was associated with multiple (presumably congenital) tumours at the base of

the skull, and in the walls of the pharynx, uvula, nose, etc., some of these containing numerous cavernous vascular spaces. In the roof of the pharynx and in the body of the sphenoid bone, etc., there was a very extensive myxochondroadenoma (Fig. 11) (resembling the congenital mixed tumour of the parotid), but in this case arising, apparently, from mucous glands; and there were also numerous adenoma-like masses of these in the uvula and neighbouring structures (Fig. 12). The pituitary fossa showed curious developmental abnormalities—for example, an opening into the right middle fossa under the thickened dura. The pituitary had evidently been the site of a glandular cystic dilatation, which had in some way become infected with streptococci, and it was obviously the rupture of this following upon the decompression and release of pressure which led to the acute streptococcal meningitis and death of the patient.

RELATION BETWEEN MULTIPLE TUMOUR SYNDROMES

Lindau,⁵ in his paper on cerebellar cysts and their relation to angiomatosis retinae, refers to the angiomata and gliomata often found in the walls of such cysts, and to the fact that in the same cases there may occur syringomyelia-like lesions in the spinal cord, as well as cystic disease of the pancreas and kidneys, hypernephroma, etc. He considers that angiomatosis of the central nervous system may be placed in the same group as tuberous sclerosis, and he believes that the multiple tumours which he describes are due to a loss of balance in the development of the mesoderm, commencing as early as the third month of embryonic life.

I am inclined to believe that there is also a similar relation between the multiple tumour syndromes of tuberous sclerosis and von Recklinghausen's disease, and also that there is a relation between the case of multiple tumours which I have just described and Lindau's disease or syndrome—thus bringing some of the "mixed cartilaginous and glandular tumours" into the same picture, and associating them, as in the case I have described, with certain of the haemangioblastic tumours of the central nervous system and elsewhere.

I put forward the suggestion that there is a distinct relation between all of these various groups of cases or types characterized by the presence of multiple tumours involving especially the central nervous system. The subject is one which, I venture to think, is worthy of discussion by the Section of Pathology, and may well form a basis of further research.

Finally, I wish particularly to acknowledge the skilled technical help of my laboratory assistants, especially Mr. John Ottewill, in the preparation of the material for the demonstration.

REFERENCES

- ¹ Fowler, J. S., and Carnegie Dickson, W. E.: *Quart. Journ. Med.*, 1910, iv, No. 13, 43.
- ² Penfield, Wilder, and Young, A.: "Observations on the Nature of von Recklinghausen's Disease and the Tumours Associated with it," *Arch. Neurol. and Psych.*, 1930, xxiii, 320.
- ³ Hosoi, K.: *Amer. Journ. Path.*, 1930, vi, 235 and 245.
- ⁴ Kernohan, J. W., and Parker, H. L.: *Journ. Nerv. and Ment. Dis.*, 1932, lxxvi, No. 4, 313.
- ⁵ Lindau, Arvid: *Acta Pathologica et Microbiologica Scandinavica*, 1926, Supplementum I; *Proc. Roy. Soc. Med.*, 1931, xxiv, No. 3, 363.
- ⁶ Wakeley, C. P. G., Worster-Drought, C., Allen, I. M., and Carnegie Dickson, W. E.: "Endothelioma of the Cauda Equina," *Brit. Journ. Surg.*, 1930, xviii, No. 69, 6.

FIBROSARCOMA OF THE SUPRARENAL GLAND

BY

BRENNAN S. CRAN, M.D.

DEPARTMENT OF PATHOLOGY, UNIVERSITY OF ABERDEEN

(With Special Plate)

The following case appears to be worthy of record on account of its comparative rarity. Within recent years certain tumours of the suprarenal gland, formerly regarded as of sarcomatous nature, more especially if round-celled, have been proved to be of sympathetic origin, and to arise from the medulla of the organ. This was first recognized by Marchand.¹ Suitable staining shows ganglion cells and an intercellular matrix, which is composed of neurofibrils. Actual sarcoma, however, may arise from the suprarenal gland, as the author's specimen shows, but in all cases it is a wise precaution to establish the nature of the neoplasm by suitable staining methods which are calculated to demonstrate the absence of nerve tissue elements. With this in mind sections from different parts of the tumour were prepared and stained according to the methods recommended (Bailey and Cushing²; Anderson³) for the demonstration of nerve cells and their processes—for example, Bielschowsky's and Cajal's methods. Haemalum and eosin, toluidin-blue, and van Gieson's and Mallory's stains were also employed. No nerve tissue elements were demonstrated in the tumour by any of the above methods.

CASE RECORD

A. C., a married woman, aged 38 years, complained for five months of a swelling situated in the left upper quadrant of the abdomen; it was steadily increasing in size. The swelling was accompanied by a great deal of discomfort on the left side, but not by actual pain. During the last few months she had lost weight rapidly, and had

a feeling of profound weakness. She had been troubled with flatulence for a number of years, but had no other gastro-intestinal symptoms except nausea at the time of menstruation. There was no history of frequency of micturition or of haematuria, and the examination of the urine was negative. She had not suffered from any previous illnesses of importance, and there was no record of abdominal injury. She had six children. She was distinctly cachectic; the temperature was normal. In the abdomen a large lobulated tumour mass was made out filling the epigastrium and extending from the tip of the ninth right costal cartilage far out into the left flank. The mass moved with respiration, was dull on percussion, and was easily distinguished from liver and spleen. The radiologist reported: "Stomach outline normal. Stomach appears to be pushed downwards and to the right side. Tumour appears to be extragastric." At operation a large retroperitoneal tumour was found apparently arising from the upper pole of the left kidney. Quite apart from the presence of several secondary deposits in the liver the tumour was hopelessly inoperable. The patient went rapidly downhill, and died three days after the operation. At no time had there been any clinical evidence of endocrine disturbance. Post-mortem examination revealed a large tumour of the left suprarenal gland in which the upper pole of the kidney was embedded, but did not show direct invasion. The omentum was firmly adherent to the peritoneal aspect of the growth, and was separated only with difficulty. Secondary deposits of growth were present in the liver and in the lungs. The viscera of the upper abdomen showed a remarkable alteration in their anatomical relationship, the stomach and colon, especially the splenic portion of the latter, being displaced by the tumour almost wholly across to the right side of the peritoneal cavity. The spleen was separated from the left kidney by the tumour for a distance corresponding fully to two hand-breadths. The tail and adjacent part of the body of the pancreas lay across the front of the tumour almost immediately underneath the anterior abdominal wall. The kidney was dislocated downwards. The right suprarenal gland appeared to be normal.

W. E. CARNEGIE DICKSON: INTRACRANIAL TUMOURS

MULTIPLE NEUROFIBROMATOSIS (VON RECKLINGHAUSEN'S DISEASE); FIGS. 1 TO 9 INCLUSIVE.

Right acoustic nerve tumour.

Meningiomatous plaque (psammoma).

Angioblastic invasion of cerebral cortex.

Lumbar nerve-root tumours (early).

Intramedullary tumour (dorsal cord).

Ditto: "palisading" of young fibroblastic cells.

Dorsal cord, central ependymal tumour.

Sympathetic ganglion (abdominal), small tumour nodule.

CASE OF MULTIPLE TUMOURS (BRAIN, PITUITARY, BASE OF SKULL, ETC.). FIGS. 10 TO 12.

Sclatic nerve trunk — minute tumour.

Cerebral cortex — cavernous angioma.

Base of Skull — chondroadenoma.

Uvula — congenital angioma-tous tumour.

MENINGIOMA: SHOWING VARIABILITY. (CASE I.) FIGS. 13 AND 14.

Parasagittal tumour — psammoma formation.

Ditto: Angioblastic reaction of tumour cells.

MENINGIOMA: SHOWING VARIABILITY. (CASE II.) FIGS. 15 AND 16.

Leptomeningioma — combination of new capillaries and psammoma-like whorls.

Ditto: active angioblastic reaction of tumour cells.