EUGENICS AND GENERAL PRACTICE

Until the doctor knows more about heredity he cannot properly advise young persons who consult him on the question of marriage because they have a growing consciousness that their duty to their offspring is no longer entirely covered by feeding, housing, and educating it, but that something can, and should, be done to protect it against tendencies to mental and physical defects, and make the stuff of the next generation better than their own. The problem of consanguinity, and its risks to children following marriage, comes more and more frequently before us, and is not an easy one to advise upon at any time. There is an increasing tendency to seek a medical overhaul by both parties prior to marriage, or even prior to engagement, and some knowledge of the laws of heredity is essential if helpful advice is to be given. It would be a sad thing if the growth of a eugenic conscience in the community anticipated sound knowledge on the part of the medical profession. Yet such a possibility is by no means a remote one. I am not sure that many of us have not been already caught with our lamps unlit in the matter of giving advice on contraception, when patients show a reasonable desire to exchange the casual, half-surreptitious, and rubber-shop affair that birth control has largely been

so far, for a really sound and scientific statement of the position, and for precise directions adapted to the particular indications of the case which is presented to us.

It is the duty of the doctor, as guardian of the health and happiness of the individual, to be well informed on all that makes for these desiderata. It is also his duty to the State to know what are the factors tending to racial deterioration, to point them out clearly, to issue the necessary warnings, and to indicate the remedies. He must be in the van in all these matters, for he has special opportunities in regard to them, and he is best fitted to make this contribution to the welfare of his fellows. True, as a servant of the community he must await the development in the community of the will to avail itself of the power of science to improve the stock. But the very fact that he has special knowledge and that he is himself a citizen—which both he and others sometimes forget—constitutes him a natural leader in the march of racial progress, and he cannot conscientiously escape this duty any more than others can conscientiously deny him this privilege. Though he may not force an issue, and though he must not speak before his knowledge is ripe, his position in eugenics is paramount; for without him this newest and most hopeful of the humane sciences must inevitably stand still.

THE LATE SURGICAL COMPLICATIONS
OF FRACTURE OF THE MANDIBLE*

BY

SIR HAROLD GILLIES, C.B.E., F.R.C.S.

PLASTIC SURGEON TO ST. BARTHOLOMEW'S HOSPITAL, LONDON COUNTY
COUNCIL, R.A.F., AND ST. ANDREW'S HOSPITAL, DOLLIS HILL
AND

A. H. McINDOE, M.Sc., M.S.MINN., F.R.C.S.

CHIEF ASSISTANT IN PLASTIC SURGERY, ST. BARTHOLOMEW'S HOSPITAL;
ASSISTANT SURGEON, PLASTIC UNIT, LONDON COUNTY COUNCIL;
SURGEON TO HOSPITAL FOR TROPICAL DISEASES

It is probably safe to say that 90 per cent. of all fractures of the mandible met with in civil life and resulting from riding, hunting, boxing, or motor car accidents will unite soundly in good position, with good dental occlusion, and without loss of external contour, provided they receive a minimum of interference from the surgeon and a maximum of expert attention from the dentist. The ordinary uncomplicated fracture of the lower jaw is, in fact, purely a dental problem, and one which does not often fall within the purview of the surgeon. In about 10 per cent. of patients in civil life, however, late complications ensue requiring operative interference. This figure was, of course, much higher during the war, when gunshot wounds and accidents, entailing much comminution and loss of bone substance, were the rule rather than the exception. The concentration of these patients at the Queen's Hospital, Sidcup, and in other centres during and after the war led to such an accumulation of valuable experience in their treatment that the care of cases in civil life has since then been much improved. Little, indeed, has been added to our knowledge of the subject in the past ten years, while it might even be said that some of the lessons learnt then have been forgotten. In this communication we have not dealt with the immediate treatment of the fractured mandible, but have confined our remarks to a discussion of the later complications of this condition as they have occurred in our experience since the war. These complications may be conveniently grouped according to whether they involve the bone or the surrounding soft parts.

COMPLICATIONS INVOLVING THE BONE

Mal-union

Union in bad position usually results from incorrect or inefficient fixation of the fragments, from unopposed muscular action or from the drag of contracting scar tissue. Except in the case of the edentulous jaw, it is a matter of considerable importance. Good dental occlusion may in fact be termed the prime requisite of successful immediate treatment. Its corollary, symmetrical external contour, is no less important, especially in woman. Mal-union is therefore particularly liable to take place where, with moderate loss of bone substance—that is, with more than 1½ cm.—efforts have been made to secure approximation of the fragments at the expense of dental occlusion. With fractures in the anterior half, the usual deformity is a lingual rotation of the fragments and a corresponding narrowing of the mandibular arch. If these cases are seen early it may be possible to separate the fragments by elastic traction, but if bony union is established and dental treatment unavailing, nothing short of osteotomy and bone grafting will suffice.

Two further deformities following fractures at the angle may result from inability to control a long posterior edentulous fragment. The posterior ramus has a strong tendency to be displaced forwards, upwards, and inwards by muscular action, where it unites with the anterior fragment to produce the well-known "flat angle." Conversely, it may be displaced forwards, upwards, and outwards, the resulting deformity being a marked prominence or outward bowing at the angle, which gives a peculiarly lop-sided appearance to the whole face. Provided good occlusion has been obtained and union is firm, it cannot be too strongly urged that these two deformities are of cosmetic importance only, and should be treated as such. A moderate degree of mal-union in fractures of this type is not incompatible with good function provided the teeth are in alignment. Attempts to control a long posterior edentulous fragment by means of intermaxillary blocks, or by direct fixation within the mouth at the time of the accident, are almost certainly doomed to failure. The only really satisfactory method is that devised by Darcissac and advocated by Ivy and Curtis. Here the angles of the jaw on one or

* Part of this paper was read to the London Medical Society, and was published in the Medical Society's *Transactions*, vol. lvi, 1933.

both sides well behind the fracture line are exposed through external incisions, and holes drilled in them through which wire hooks may be passed. Elastic traction in a downward and backward direction, taking purchase from rods embedded posteriorly in a plaster-of-Paris head-cap, can then be obtained, and good apposition of the bone ends effected.

Osteotomy may be necessary for severe malocclusion, or for marked irregularity of external contour such as occurs in the case of the too prominent angle. It is performed either by means of a Gigli saw introduced along a trocar through a tiny submandibular incision and carried round the inner surface of the mandible through the floor of the mouth, or, more directly, after exposure of the bone by means of the chisel. After division of the jaw the fragments are reset and resplinted or held firmly apart by a dental appliance of the cap-splint and rigid-bar type preparatory to bone grafting.

In the case of the "flat angle" (occurring after bony union) a simple compromise is effected by grafting fat beneath the skin over the depressed area until the contour is even on both sides.

Delayed Union

Delay in union of the fragments of a fractured mandible, as with any other bone, may be a matter of months or years. It is the most common surgical complication in civil practice, and, from the preventive standpoint, easily the most important. In our experience the factors responsible for delayed healing are usually multiple, and in any given instance the relative importance of each is difficult to assess. It is therefore essential that every possible cause be investigated and carefully eliminated.

Although most fractures of the mandible are compound and are necessarily infected from the buccal cavity, sepsis *per se* does not appear to be the deciding factor in delayed union. Everyone has seen sound bony union occur in the continued presence of sepsis, and repair become almost complete before the infected bone clears up or the last sequestrum is discharged. That it is of importance, however, cannot be denied, particularly if it is associated with an extreme degree of bone necrosis and comminution, or with surgical or other foreign bodies in the neighbourhood of the fracture line. Among the ordinary foreign bodies we include interpolated soft tissues, sequestra, and teeth or portions of teeth. The last we feel to be most important, and especially so if the x-rays show periapical absorption. Other infected teeth are also important foci of chronic sepsis. Surgical foreign bodies include anything which has been introduced at the time of the injury, but, more particularly, wires, plates, or screws used by the surgeon in an unwise effort to secure a precarious immobilization by direct means. We are strenuously opposed to any method of intra- or extra-oral fixation entailing foreign bodies which penetrate the medullary cavity of the bone ends and lie in the fracture line. These must soon cause necrosis and absorption. If none of these factors appears to be present poor approximation of the fragments or inadequate fixation must be held responsible.

The treatment of patients presenting themselves with delayed union must therefore include, in conjunction with a competent dentist, a careful review of the methods of previous fixation and of the reasons for their inefficiency. An investigation must then be made into the conditions present at the site of fracture, and a thorough search begun for all possible causes of continued sepsis. Good x-rays in many positions are, of course, essential. With removal of all foci of infection, drainage of abscesses, and interdental fixation in good position, sound union may be predicted in most cases. As an adjunct to treatment electro-vibro-massage and

other methods of increasing the blood supply at the bone ends are useful. The splints should be removed at the earliest possible moment, and simple massage and voluntary movements instituted. In a few instances, however, bone fails to be laid down in the fracture line, and the case becomes one of non-union.

Non-union

Absolute failure of union with the development of a pseudo-arthritis is rare in civil practice, but it may follow any of the causes enumerated above. Thus, while it may not be considered as established for many months in the case of preventable causes being allowed to persist, it may be determined immediately by the interposition of soft tissues, or by extensive loss of bone at the time of the accident. It was fairly common during the war in gunshot wounds with severe sepsis and much comminution, and is seen at present as a result of explosions and bad collision accidents. Strenuous efforts have usually been made to obtain union, but the most prolonged fixation and careful treatment are unsuccessful. Between the eburnated bone ends lies a mass of fibrous tissue, which effectually precludes the chance of osteoblastic activity. The amount of movement between the fragments varies considerably, and in certain cases where occlusion is reasonably good and bone loss small the fibrous union may be sufficiently firm to give a satisfactory functional result. These cases require no interference or, at the most, careful removal of the interposed tissue, freshening of the bone ends, and resplinting in good position. For losses over $1\frac{1}{2}$ cm., however, it is usually necessary to graft.

Loss of Bone Substance

As this condition constitutes the most serious of all the surgical complications of fractures of the mandible it warrants careful attention. It may be stated at once that whenever the loss of bone substance has been so extensive that reasonable approximation of the fragments and occlusion of the standing teeth are incompatible, bone grafting will become necessary.

In general, these conditions are fulfilled when the final gap exceeds 1.5 cm., although exceptional instances occur when bony union takes place with even wider separation. Approximation of the fragments in recent fractures with loss of bone would probably result in more certain bony union, but only at the cost of complete loss of dental occlusion and marked distortion of the jaw. Only when the mandible is completely, or almost completely, edentulous, or in old and enfeebled patients in whom grafting is impracticable, is it permissible to secure approximation at the expense of occlusion. On the other hand, replacement of the fragments in normal position, irrespective of the amount of bone loss, places the surrounding soft parts under normal tension, largely precludes the possibility of their interposition, and allows the gap to fill with blood clot in which osteogenesis may proceed normally. Torn strips of periosteum, the presence of which can only be guessed at, may cross the defect and hasten the process materially, thus accounting for those cases in which bony union has resulted in gaps of apparently over 1.5 cm. Ample opportunity must be given for union to take place spontaneously, and all the factors already discussed must be carefully considered before bone grafting is decided upon. Non-union with good occlusion, due to too great separation of the fragments, which can be corrected by bone grafting, is to be preferred to bony union by approximation obtained at the cost of loss of function.

It is of supreme importance to wait until all sepsis has subsided and all possible foci of infection and foreign bodies are removed. At least three months should elapse

from the last signs of sepsis before bone grafting is undertaken. In the meantime, the fragments are controlled by means of splints adapted to the individual case. Experience during the war has given us a wide variety of methods of bone grafting, each with its own indications. It is not proposed to discuss full details of operative technique, but rather to indicate the advantages and disadvantages of each.

BONE GRAFTING

Pedicle Graft.—This method, advocated by Cole, consists in the removal of a block of bone from the lower border of one of the fragments, and shifting it bodily, complete with its muscular attachments, into the gap. It has the advantage of being easy to perform, is not liable to sepsis, and may be used for small defects. Its disadvantages are that the union is liable to be springy, as it occurs only along the lower border of the mandible, and no regeneration of the bony tissue of the alveolar portion takes place. It is unsuitable for large gaps and, cosmetically, it is indifferent. On the whole, it offers little over the plain block graft from the ilium or tibia.

Block Tibial Graft.—The adoption of the simple method of cutting a suitable block of bone from the tibia and wiring it between the fragments in an end-to-end position has long been advocated by Albee. Lane first showed that the use of wire in the absence of sepsis is a perfectly safe procedure, and that it gives considerable inherent stability to the fragments during healing. The tibia as a source of bone, however, has certain disadvantages. The graft is difficult to shape and to manipulate, and its removal may be followed by a good deal of weakness in the leg. Furthermore, the osteoblastic activity of the graft is not great. For these reasons tibial grafts have been largely abandoned except for very small defects.

Osteoperiosteal Graft.—This graft, attributed to Delagénière, is obtained by removing a very thin strip of cortical bone from the tibia, complete with the overlying periosteum. It is divided into four pieces, which are then inserted into the prepared mandibular defect so that they entirely enclose the gap with the bone surfaces facing each other across it. The grafts contain all the elements necessary for osteogenesis, and at the same time are flexible and easily adjustable. No disability of the leg follows this operation. Consolidation, however, requires a much longer period, and during this time no dependence can be placed on the graft itself for maintenance of the mandibular fragments. It is useful for very small or for very large gaps, and, in particular, where an acute curve is required, such as in the region of the symphysis. It is also useful where the blood supply of the bone ends is poor, and in children where the crest of the ilium is still cartilaginous.

Osteochondral Rib Graft.—This method (Gillies, 1919) entails the use of the seventh or eighth rib of the opposite side, including the costochondral junction and the bone as far round as the point of maximum convexity. It is only indicated where the loss of bone includes the whole of the ascending ramus. The bony portion of the graft is wired to the refreshed anterior fragment, while the point of maximum convexity forms the new angle of the mandible, and the ascending ramus is represented by that portion of the costal cartilage which runs upward to the sternum. A false joint in the neighbourhood of the glenoid fossa is thus made, and a cosmetic and functional result accrues.

Iliac Graft.—This is so superior to all other types that it has almost entirely superseded grafts from other sources, except in children, where the crest of the ilium is cartilaginous. A curved piece of bone, easily bent

and manipulated, and consisting of a tough outer shell of compact bone and a considerable amount of cancellous tissue, may be cut with ease from the iliac crest behind the anterior superior spine without any subsequent disability. The osteogenetic power of this graft is high. As originally described by Lindemann, it consisted in a shaped iliac block wired between the fragments with end-to-end union. Chubb, who elaborated the method with extraordinary success at Sidcup, obtained perfect union in over 90 per cent. of 100 successive cases. He states that the aim in every case is to obtain a sound carpentering junction between the graft and the jaw fragments, the bones being so trimmed that they fit end-to-end with accurate apposition of the largest obtainable area of exposed vascular bone tissue. The method was evolved principally to control the edentulous fragment, and to facilitate union by early removal of the splints so that the junctions are exposed to the stresses of voluntary muscular action. Bone deposition is greatly stimulated by early stress, so that union of ivory hardness may be obtained at the end of three to four months.

A simpler method, and one not demanding so high a degree of craftsmanship, is to clear the ends of the fragments for 1 to 2 cm., to remove a shaving from the outer side of the bone ends, and to wire the graft to them in close overlapping apposition. We have obtained good results in this way. Kilner uses mattress sutures of wire introduced through appropriately placed holes drilled in the graft and the bone ends. This enables considerable dependence to be placed on the stability of the graft for the replacement and control of the posterior fragment. Figi utilizes chromic catgut by the same technique. Billington and Round omit all fixation and splints, and merely place the graft in a soft tissue cavity overlapping the bone ends, but in close contact with them. Two weeks later, while the union is soft, the jaw is manipulated into position and splinted. Experience shows, however, that some control of the posterior fragment is desirable.

The present position of bone grafting of the mandible may therefore be summarized as follows.

1. The ilium is the best source of supply, and the iliac graft will be found satisfactory in the vast majority of cases. It should be fixed by wire, and an overlapping junction probably results in earlier union and is certainly easier to perform than an end-to-end junction.
2. The pedicle graft is useful for small defects, but has no advantages over the plain block iliac graft.
3. The osteoperiosteal graft is indicated in very large gaps, in very small controlled gaps, and in tissues the blood supply of which is poor; it is also useful for bony losses in the symphysis region, where a marked curve is required.
4. The osteochondral graft is indicated where the condyle of the mandible is lost, together with the whole of the ascending ramus. It has a distinct field where half the mandible is removed surgically and a false joint is required.

Fixation

In any of the special circumstances which have been enumerated the method of fixation is particularly important. There are, in general, three types of intra-oral fixation: interdental and intermaxillary wiring; metal cap splints; circumferential wiring. The first depends upon sound standing teeth in both upper and lower jaws, though not necessarily in the posterior fragment. The second requires sound standing teeth in both upper and lower jaws, or standing teeth in the lower jaw only, provided they are in both fragments. Circumferential wiring is carried out by passing a wire round the fragment so that it may be hitched up to an intra-oral plate or splint bridging the gap. It is of particular use in edentulous patients

and where the central edentulous fragment has become depressed below the level of the two adjoining fragments. There is no doubt that for immediate fixation in the majority of cases intermaxillary or interdental wiring is the method of choice. It is quick, easily applied at short notice, and requires no elaborate dental fitting or apparatus. Metal cap splints, when properly applied, undoubtedly give the most accurate fixation, and should be used where intermaxillary wiring has failed, where delayed union makes the most perfect fixation necessary, or preparatory to bone grafting. They are particularly indicated where, with standing teeth in both fragments, rigid fixation may be obtained without immobilizing the temporo-mandibular joint.

COMPLICATIONS INVOLVING THE SOFT PARTS

Obliteration of the Buccal Sulcus

In nearly all cases which have been bone-grafted loss of the alveolar ridge obliterates the inferior buccal sulcus. In a few severe fractures of the lower jaw also the injury is so extensive that the sulcus is partly or completely destroyed. In riding, flying, or motoring accidents, or in those resulting from explosions, the lower lip or cheek may be so severely lacerated that the same effect is produced. Other causes are extensive destruction of the alveolar ridge and continued sepsis at the site of fracture. When the fracture heals and the sepsis clears up it is found that the buccal sulcus is so shallow as to make the fitting of a stable denture impossible. The treatment of this complication is exceedingly satisfactory. By the Sidcup modification of the method of Esser, called the "buccal inlay," the adhesions between the mandible and the lower lip or cheek are separated, and the resulting raw area grafted with a Thiersch graft cut from a hairless portion of the body, preferably the inner side of the upper arm. Pressure is maintained on the graft by means of a Stent mould of the defect held by a dental appliance previously fitted to the lower teeth. After ten days, when epithelialization is complete, the mould is removed and is replaced by a gutta-percha or vulcanite mould of similar shape. This is maintained *in situ* until all tendency for the cavity to contract has ceased. When this period is over the denture can be fitted.

Facial Deformities Resulting from Fracture of the Mandible

In dealing with mal-union mention was made of the treatment of irregularities of contour of the jaw, such as the "flat" or too prominent angle. Any such unevenness may be simply dealt with, on the one hand, by means of fat grafts introduced at appropriate sites, and on the other by the removal of excess bone, so that the jaw line is smoothed out and normal symmetry regained. In addition to concealing the defects of bony contour, it is frequently necessary to excise and resuture depressed scars of the chin or cheek caused by the original injury. These are frequently associated with the deep depressions adherent to the bone, which indicate the former presence of sinuses. Careful excision of all scar tissue and freeing of the skin from deep structures, together with local rearrangement of fat flaps or the introduction of fat from elsewhere, followed by resuture, will do much to obliterate all trace of what would otherwise remain as an unsightly disfigurement.

BIBLIOGRAPHY

- Chubb, G.: *Lancet*, 1920, ii, 9.
 Idem: *Ibid.*, 1921, i, 640.
 Billington, W., and Round, H.: *Brit. Journ. Surg.*, 1925-6, xiii, 497.
 Cole, P. P.: *Lancet*, 1917, i, 415.
 Fry, W. Kelsey: *Treatment of Injuries to the Jaw*, Cartwright Prize Essay, R.C.S., 1916-20.
 Idem: *Proc. Roy. Soc. Med.*, 1929, xxii, 663.
 Figi, F. A.: *Surg., Gynecol. and Obstet.*, 1932, lv, 762.

- Gillies, H. D.: *Plastic Surgery of the Face*, 1920, Chapter IV.
 Idem: *Carson's Modern Operative Surgery*, 1924, ii, p. 374.
 Gillies, H. D., and McIndoe, A. H.: *Med. Soc. Trans.*, 1933, lvi, 30.
 Gillies, H. D., and Kilner, T. P.: *Lancet*, 1932, ii, 1369.
 Lane, W. Arbuthnot: *The Operative Treatment of Fractures*, 1914.
 Ivy, R. H., and Curtis, L.: *Fractures of the Jaws*, Lea and Febiger, Philadelphia, 1931.

THE CRONIN LOWE SERUM REACTION FOR MALIGNANCY

BY

JOCELYN PATTERSON, M.Sc., Ph.D.

BIOCHEMIST TO CHARING CROSS HOSPITAL

AND

JOSEPH ADLER

(From the Biochemical Department, Charing Cross Hospital, Institute of Pathology)

Widespread interest has been aroused by the recent publication by Cronin Lowe¹ of a reaction claimed to be diagnostic of malignancy. This new test is a development of the vanadic acid flocculation reaction devised by Bendien² which, when coupled with a means of subjecting the precipitated protein to a spectrophotographic examination, was said to lead to results of diagnostic value. It has now been generally accepted, following upon the serious criticisms levelled against the value of the spectrophotographic examination by Smith, Holiday, and Marrack,³ that the serum reactions as described by Bendien do not provide any basis for arriving at a diagnosis of malignancy; hence the need of replacing them by others likely to prove more successful.

The test devised by Cronin Lowe is of a triple nature, the vanadic acid preparation of Bendien being performed on (1) untreated serum, (2) heat-inactivated serum, and (3) ether-extracted serum. In general, with a given vanadic acid mixture in the region of "first precipitation," if A, B, and C are used to denote the yields of protein deposit resulting respectively from each phase, then A is greater than B and less than C. If from these values a ratio be computed in the form of $\frac{C-A}{A-B}$, then a ratio greater than unity is observed for malignant cases, while other serums show a ratio less than unity. It is claimed that on this basis an accurate diagnosis as to the presence or absence of malignancy can be made in at least 95 per cent. of cases, and if certain grossly septic cases held to be responsible for introducing a complication into the reaction be left out of account the figure is raised to 98 per cent. These results are so striking that if the necessary confirmation be forthcoming the Cronin Lowe reaction would represent practically the most valuable diagnostic test known to clinical medicine. The present investigation was undertaken with a view to formulating an opinion with regard to the value of the test.

Our initial results fell lamentably short of the success which had been reported, and, after full investigation, and seeing no hope of reconciling our findings with his, one of us communicated with Dr. Cronin Lowe, who thereupon kindly provided us with very full technical details upon all the points of procedure which were regarded as being likely to cause discrepancies in results. Further work performed with the aid of this extra information still left us far short of achieving a diagnostic reaction, and we are thus obliged to conclude our investigation by presenting results which provide no confirmation of the usefulness of the triple reaction as a test for malignancy. It should be made clear that the present paper deals solely with results obtained subsequent to receiving the private communication from the original author.