the departure of the latter from the island, became shorn of its beams, and Hygeia was overshadowed by a cloud. Before the end of the year "fever of a severe bilious remittent type became prevalent," and three cases proved fatal. In 1874, there were 221 cases admitted under the head of fevers, and "most of these cases were complicated with enteric symptoms, in many cases severe."

Now, *why was the 29th regiment so exceptionally fortunate?* I think that anyone who has studied the medical history of European regiments in the West Indies, will admit that this question deserves to be italicised. For a more or less complete answer to it I must refer to papers of mine in the *Lancet* for 1880, volume 1, page 560 and 599. Here, I will notice only one point. Whilst other regiments had been sent to the West Indies from more or less debilitating climates, such as Gibraltar, Malta, or Bermuda, the 29th was sent from a very healthy and invigorating command, from Canada, where, for nearly two years and a half, it had been laying in a stock of disease-repelling vigour. Two years in such a climate must impart to the human economy a resisting power against morbid forces, very different from that to be gained by a residence of the same period in Malta.

I do not, of course, mean for a moment to insist that a regiment sent from Malta to the West Indies must suffer; or that a regiment sent from such a climate as Canada cannot suffer; but, I desire to strengthen the idea that a very important factor in estimating the effects of deteriorating or of lethal influences on any regiment, is a consideration of the hygienic characters of the place from which it comes; and, believing this, I hold there is much to be thought of before sending a regiment from Malta to Barbadoes to take the place of one removed therefrom in consequence of an outbreak of yellow fever.

Whether the few facts I have brought together, and the remarks which I have made, will convey to other minds the impression produced on my own, I cannot say; but the end which I have in view will certainly not be attained, if they are found wanting in suggestiveness.

POISONING BY CHROMATE OF LEAD IN WEAVERS.

By ROBERT C. SMITH, M.D.,

Bachelor of Science in Public Health, Univ. Edin., Manchester.

ON February 12th, 1881, Mrs. B., aged 30, a weaver in a cotton-mill, came to me for advice and treatment. She had been unable to follow her work for the previous six weeks, and was suffering from great weakness, wandering pains in her limbs, and anæmia. Her gums showed well-marked signs of lead-poisoning. There was no albumen in her urine. She attributed her illness to the inhalation of a yellow dust that was given up from the yarn in the process of weaving an orange-coloured cloth. She had always enjoyed good health before being employed at this class of goods. She gave, as a reason, that it was the coloured dust (lead dichromate) which had made her ill; that her fellow-workers, when engaged at the same kind of work, sooner or later sickened; and that there were many others at home ill from the same cause.

Mary C., also a weaver at the same mill, applied for advice a few days after Mrs. B.; and had, like her, been engaged as a weaver of orange yarn. The blue line was very distinct. Her breath was offensive, and the skin and conjunctivæ were of a distinctly yellow colour. There was no albumen nor bile in the urine; nor had she any tenderness over the liver. She suffered from tormina and wandering pains over the body, and was very costive. Under treatment, the yellowness of skin passed away; but the blue line remained long after other symptoms of lead-poisoning had disappeared.

Jane C., sister of the above, was visited by me at her own home on the same day. I found her also affected with lead-poisoning. She appeared at first sight to be suffering from an acute attack of jaundice: her skin and conjunctivæ were intensely yellow in colour. There was obstinate sickness, and purging of dark sap-green motions (probably stained with chromium oxide). The urine contained albumen and a trace of chromium, but no bile nor lead. This patient recovered with great difficulty—the staining of the tissues, sickness, and purging ceasing first, but the blue line is still yet present.

The above are types of cases that were of constant occurrence, about nine months ago, in the practices of medical men in the neighbourhood of the mill in question. I had eight patients at once from the same mill under treatment for this form of lead-poisoning. Another medical man has had twenty, and a third has had from thirty to forty similar cases in all under observation. One of the mill-hands, under the care of a fourth medical man, dying from lead-poisoning, an inquest was held, when, at the *post mortem* examination, this gentleman found distinct traces of lead in the liver. Public opinion having been drawn

to the dangerous nature of the work, pressure was brought to bear on the employers, who have since then caused the yarn to be more carefully dyed and prepared before being woven. They have also insisted on the winders and weavers wearing a muslin respirator over the mouth and nose, when at work; and the use of hooks to thread the shuttle, instead of sucking the web through its eye with the mouth, under a penalty. By these precautions having been rigorously carried out, the operatives, though still weaving chromes, no longer suffer from lead-poisoning.

In the cases which I have observed, the yellowness of the skin is the first symptom to disappear, and the blue gums the last; and chromium is found in the urine, and probably in the feces. I therefore conclude that, after absorption, the dichromate undergoes decomposition within the blood; the lead being fixed in the tissue, whilst the chromic acid combines most probably with soda—a compound which is intensely yellow, and stains the liquor sanguinis and skin for a time; and, finally, leaves the body by the liver and kidneys.

OVARIOTOMY IN GENERAL HOSPITALS.

By JAMES E. ADAMS, F.R.C.S.,

Surgeon to the London Hospital.

THE desirability of performing ovariectomy in general hospitals must be admitted to be a matter requiring some consideration, seeing that there are some surgeons, my colleague Mr. Hutchinson in particular, who think it better to send all patients requiring this operation to special hospitals; and this opinion is mainly based on the improving statistics of the practice of those who have made this operation a specialty, and through whose praiseworthy efforts the operation has become one of the most brilliant in surgery. There are, however, many reasons against declining to do this operation in ordinary hospitals, especially where there are large schools; and I hope to be able to show that patients operated on by us do just as well, and show as good an average as those dealt with by others. What reasons are there against it? Is it because we cannot have a sufficiently pure air, pure dressers, nurses, and materials? I aver that—by complete disinfection, isolation, Listerian methods of operating, careful selection of specially trained nurses, rigorous exclusion of all sponges and dressings not absolutely new and carbolicised—we can place our patients in conditions just as favourable as in any special hospital. If we could not, we ought to abstain from opening the peritoneal cavity at all, and decline all kinds of abdominal section. Do we lack the necessary manipulative skill or knowledge in dealing with difficulties? As to the former, the hands of any skilled operating surgeon should be good enough for the work; and to assume that we lack the latter, is paying a poor compliment to the lucidity of the various writers who have so carefully recorded and published such large masses of evidence as are now before us. My view is, that the specialists have given us the means of learning, and that it is the duty of surgeons in large hospital practice to acquire knowledge and to diffuse it. Nevertheless, I would not advise any patient to undergo any operation in my wards, if I thought it could be done with less risk in a special hospital; but this is the very thing which I do not believe; and I undertook my ovariectomies with the firm conviction that the preventable causes of danger or death could be as thoroughly eliminated in the London Hospital as elsewhere. I can only adduce the facts that occur in my own practice, and they are very few in number; but, so long as the belief holds that ovariectomy is not safe in a general hospital, the opportunities for extending our experience and enlarging our statistical tables are not likely to be frequent. Of five ovariectomies (properly so-called), including one woman aged sixty-one, where there were numerous adhesions, all recovered. There were two cases, besides, in which an operation was undertaken with a full knowledge of the probable difficulties and dangers, which terminated fatally; one being a soft solid malignant tumour involving the peritoneum, and the other a large succulent fibroid attached to the fundus of the uterus. The statistics, including the cases of my colleagues, extend back as far as 1878, and show a mortality of 23 per cent.; and even this, though including one or two very unfavourable cases, will bear comparison with some of the numerous series lately published. No one is more gratefully appreciative of the results obtained by, and the means of acquiring information given by, the masters and pioneers in this branch than myself, and I have no desire to acquire a special reputation as an ovariectomist; but I should like to show, if possible, that it is not only justifiable, but desirable, to perform these operations in those hospitals where patients seek relief in the first instance, instead of having to banish them to a distant institution. My argument, if I have made myself clear, is not in the direction of bold-