One surgical aid to medicine, gentlemen, I have reserved for the pleasure of making it my last word to night. This, although not mechanical, has done more than any one of those gifts of surgery to mankind to enlighten our darkness, to fortify our reason and to guide our handicraft—I mean the discovery and demonstration of that great agency which, dimly guessed at for many years, has but recently come fully into light—the work of germinal matter, or catalytic matter in the processes of death. These revelations we owe to many workers, perhaps to none more than to M. Pasteur, who is neither physician nor surgeon. But the working of these agents must be demonstrated in the simple and visible before it can be realised in the complex and invisible—on the surface of the body before the inward parts; and I am speaking to you as to one man when I say that which surgeons owe to the genius of Mr. Lister is but the beginning of the debt due to him from physicians. To him we owe that demonstration upon the outer surfaces of the body of the forces and changes which fail not to reach and disturb the innermost; and I would dwell upon the name of Mr. Lister as the last I will cite to-day of the surgical aids to medicine—an aid without whom many of the measures I have advocated to night would have been impossible.

SELECTIONS FROM CLINICAL LECTURES,

Delivered at the London Hospital.

By JONATHAN HUTCHINSON, F.R.C.S.,

Senior Surgeon to the Hospital; Consulting-Surgeon to Moorfields Ophthalmic Hospital; and Professor of Surgery and Pathology in the Royal College of Surgeons.

THE PRE-CANCEROUS STAGE OF CANCER, AND THE IMPORTANCE OF EARLY OPERATIONS.

GENTLEMEN,—The patient who has just left the theatre is the subject of cancer of the tongue in an advanced stage. As I demonstrated to you, the lymphatic glands are already enlarged. It is hopeless to think of an operation, and there is nothing before him but death, preceded and produced by a few months of great and continuous suffering. His case, I am sorry to say, is but an example of what is very common. Not a month passes but a case of cancer of the tongue presents itself in this condition. The cases which come whilst the disease is still restricted to the tongue itself are comparatively few; nor does this remark apply only to the tongue. "Too late! Too late!" is the sentence written but too legibly on three-fourths of the cases of external cancer concerning which the operating surgeon is consulted. It is a most lamentable pity that it should be so; and the bitterest reflection of all is, that usually a considerable part of the precious time which has been wasted has been passed under professional observation and illusory treatment. In the present instance, the poor fellow has been three months in a large hospital, and a month under private care. I feel free, gentlemen, to speak openly on this matter, because my conscience is clear that I have never failed when opportunity offered, both here and elsewhere, to enforce the doctrine of the local origin of most forms of external or surgical cancer, and the paramount importance of early operation. I have tried every form of phraseology that I could devise, as likely to impress this lesson. Nearly twenty years ago, I spoke to your predecessors in this theatre concerning the "successful cultivation of cancer"; telling them how, if they wished their patients to die miserably of this disease, they could easily bring it about. The suggestion was, that all suspicious sores should be considered to be syphilitic, and treated internally by iodide of potassium, and locally by caustics, until the diagnosis became clear. More recently, I have often explained and enforced the doctrine of a pre-cancerous stage of cancer, in the hope that, by its aid, a better comprehension of the importance of adequate and early treatment might be obtained. According to this doctrine, in most cases of cancer of the penis, lip, tongue, skin, etc., there is a stage—often a long one—during which a condition of chronic inflammation only is present, and upon this the cancerous process becomes engrafted. I feel quite sure that the fact is so. Phimosis and the consequent balanitis lead to cancer of the penis; the soot-wart becomes cancer of the scrotum; the pipe-sore passes into cancer of the lip; and the syphilitic leucoma of the tongue, which has existed in a quiet state for years, at length, in more advanced life, takes on cancerous growth. The frequency with which old syphilitic sores become cancerous is very remarkable; on the tongue, in par-

ticular, cancer is almost always preceded by syphilis, and hence one of the commonest causes of error in diagnosis and procrastination in treatment. The surgeon diagnoses syphilis, the patient admits the charge, and iodide of potassium seems to do good; and thus months are allowed to slip by in a state of fools' paradise. The diagnosis, which was right at first, becomes in the end a fatal blunder, for the disease which was its subject has changed its nature. I repeat that it is not possible to exaggerate the clinical and social importance of this doctrine. A general acceptance of the belief that cancer usually has a pre-cancerous stage, and that this stage is the one in which operations ought to be performed, would save many hundreds of lives every year. It would lead to the excision of all portions of epithelial or epidemic structure which have passed into a suspicious condition. Instead of looking on whilst the fire smouldered, and waiting till it blazed up, we should stamp it out on the first suspicion. What is a man the worse if you have cut away a warty sore on his lip, and, when you come to put sections under the microscope, you find no nested cells? If you have removed a painful, hard-based ulcer of the tongue, and with it perhaps an eighth part of the organ; and, when all is done, and the sore healed, a zealous pathological friend demonstrates to you that the ulcer is not cancerous, need your conscience be troubled? You have operated in the pre-cancerous stage, and you have probably effected a permanent cure of what would soon have become an incurable disease. I do not wish to offer any apology for carelessness, but I have not in this matter any fear of it.

EMPIRICISM AND SPECIFICS.

The patient whom we are about to discharge from Talbot ward, cured of severe pemphigus, was admitted for a special purpose. He was sent in by my friend and former pupil Dr. Tom Robinson, in order that he might be cured. You will say that the hope of cure is the motive which brings most of our patients to us. True; but in this instance there was something more than this. Dr. Robinson could easily have cured him himself, but he sent him here in order that I might do the miracle of cure under your eyes, and thus claim your belief in the efficacy of drugs. You will remember his state when admitted; he was covered from head to foot with bullæ; the trunk was less severely affected than his limbs, head, and genitals; on these, there was nowhere a space as large as the palm free from bullæ, and on the trunk also there were a considerable number. He was in a miserable condition from pain and irritation. The eruption had been out about ten days, and it affected the mucous membrane of his mouth as well as the skin. You may remember that we kept him in bed for a few days before we used the magician's wand, in order that all might see that there was no natural tendency to amelioration. More bullæ came out; then, without making the slightest change in diet, we ordered a few drops of a tasteless solution of arsenic to be swallowed three times a day. The result was that, at our next visit, most of the bullæ had dried, and there were no fresh ones. He continued to improve greatly for ten days, when suddenly a few fresh small bullæ seemed to threaten a relapse. We doubled the dose of our remedy, making the dose eight instead of four drops; and, from that day, with the most trifling exception, the recovery has been uninterrupted. With such a fact before you, let me beg of you, gentlemen, to believe in drugs, and to treat empiricism with respect. In the prescription which I ordered, I availed myself solely of empirical knowledge; I prescribed, just as any old woman might prescribe, that which I knew would do good. Concerning the nature of pemphigus, I knew nothing; of its cause, absolutely nothing; of its clinical relationship, but little; of the *modus operandi* of arsenic, I knew scarcely more; but this I did know as a fragment of assured conviction, that arsenic would cause the pemphigus eruption to disappear, and the patient to regain his health. Far be it from me to speak slightly of scientific work; let us by all means work as hard as we can in the laboratory and microscope-room, and penetrate as far as we possibly can into the mysteries of disease; let us never weary in our search after causes, or in our endeavour to find practical application for the facts of physiology. But, whilst doing this, let us remember that, as regards the relief of suffering, much of our usefulness must be based upon knowledge which is nowise scientific, but simply a matter of experience and memory. We have many specifics for many maladies, or rather for many symptoms, and he is the most successful practitioner who has stored in his memory the largest number of them. As years go on, we shall add many more to our list; and I doubt not that there are those who now listen to me who are destined to give help in their discovery; for discoveries in this direction are rarely made by single observers, but rather by the concurrent work of many experimenters, all keeping their eyes open, willing to try new things, and resolute to store faithfully the results of their observations. Iodide of potassium for tertiary syphilis, the bromide for epilepsy and as an anaphrodisiac, iodoform for phagedena and specific ulceration, balsam

of Peru for scabies; so silently have these invaluable specifics been introduced into practice, that it would puzzle most of us to say who first recommended them. I mention this fact, in order to show how important is the honest labour of all in the pursuit of therapeutics. We all prescribe, and we ought all, on system, to observe and record the results of our observation as to the effect of drugs. Five-and-twenty years ago, I believe that the case of pemphigus which you have seen cured would have been found incurable in all the medical institutions of the world, with one single exception. Much more recently than that, the disease was pronounced by Hebra to be invariably fatal. So, indeed, it would have been to this day, if we had not found out arsenic. I know of nothing else that will cure it. Our patient was already beginning to emaciate, and, in the course of a few months—possibly of a few weeks—he would have had to die, worn out by the constant discharge from his skin, had we not put the arsenic into his blood. Never shall I forget seeing a poor wretched child carried on a bed into Mr. Startin's out-patient room at the Blackfriars Hospital for Skin-Diseases. It had been brought straight from the wards of one of our largest hospitals, where, during three months, all had been done for its help that benevolence, aided by the science of the day, could suggest. Yet it was emaciated to skin and bone, and so covered with sores, that it was impossible to put its clothes on. A few minims of arsenic were prescribed, and in a few weeks the child was well. So much for empirical knowledge; so much for drug-specifics.

PROMPT AMPUTATION IN TRAUMATIC GANGRENE: IMPORTANCE OF AMPUTATION HIGH UP.

In cases of traumatic gangrene, ought amputation to be performed without waiting for a line of demarcation to be formed? I believe that the reply of most surgeons to this question will be an unhesitating affirmative. Such certainly would be my own. We have recently had a very instructive case. A man aged more than 50, but of good constitution, was admitted with a compound fracture of the lower third of the leg. We tried to save it, and the limb was put up in antiseptic dressings. The foot, however, became gangrenous, and, about the sixth day after admission, Mr. Tay amputated the limb below the knee, the man being at the time very ill. The amputation was done through perfectly sound parts, but it was presently followed by gangrene of the stump. The flaps became livid, and the man was in a most urgent condition. Mr. Tay and myself, in consultation, determined at once to perform a second amputation; and, within twenty-four hours of the first, this was done in the lower third of the thigh. The man did well, and the stump on the second occasion has made, as you saw the other day, a very good one. The main reason for prompt amputation in such cases is, that the gangrenous process is a very dangerous one. Whilst soft parts are dying, and the circulation still going on to some extent through them, the blood becomes poisoned by the absorption of gases and fluids from the putrescent parts, and a most dangerous condition of septicæmia results. Of this state, a rapid pulse, a sunken countenance, high temperature, and vomiting, are the most constant signs. It is remarkable how quickly they are sometimes relieved by the removal of the dying part. It may be that the process of mortification is also attended by shock to the nervous system, but I suspect that the chief part of the mischief is done through the blood. In the pyæmia which results from phlebitis, it is of no use to amputate after once the poisonous emboli have been shed from the inflamed vein into the blood. It is then too late, for the secondary abscesses will form, whether you remove the original focus or not. In the septicæmia from gangrene, however, the case is different. Here it seems to be easily possible for the blood to rid itself of the contamination. I well remember the case of a young soldier who was under treatment some years ago for a damaged foot, the consequence of a Canadian frost-bite. He had also obliteration of his femoral artery. My junior colleague at the time amputated through the tarsus. The stump never healed, and, some time later, I amputated in the upper third of the leg at a great distance from the disease, for the whole of his leg looked at the time as healthy as yours or mine. I went high up, because I knew that the femoral artery was occluded. The result, however, was, that the stump passed into gangrene, and very soon we had all the symptoms of the most severe form of that malady. The patient had frequent vomiting, a very rapid pulse, and was indeed in such a critical state when on the third day I decided to amputate again, that I did not dare to have him taken from his bed. The second amputation, performed high up in the thigh, saved his life. No ill symptoms occurred after it, and the stump healed well. I am inclined to believe that the usefulness of amputation in gangrene will become more widely appreciated, and that this measure will be resorted to, not exclusively in traumatic gangrene, but in all forms which are attended by serious constitutional symptoms. If a part be simply passing quietly into a

mummified condition, and the patient's health not suffering, then there is no reason for interfering until you see where nature is going to make the separation. There is, indeed, no reason for interfering at all, for you must let nature finish the work. If you amputate near to the line of demarcation, your stump is almost certain to slough, and all that you must dare do in the way of help in such cases is just to saw through the bones when they are laid bare. The explanation of disappointment in amputating for gangrene, whether traumatic or otherwise, is, I feel sure, almost always from amputating too near to the disease. In all such cases, we ought always to go high up. If the foot be concerned, go above the knee; if the upper extremity, near to the shoulder. You must think rather of the patient's life than of the length of his stump. Adopting this rule, I have of late years more than once amputated for severe forms of senile gangrene with very excellent results.

CAN A MAN HAVE SYPHILIS TWICE?

The man whom we have just seen offers a remarkable example of the occurrence of a second chancre soon after the first. His second sore has been, as I have repeatedly demonstrated, characteristically indurated. He is quite candid, and makes no doubt that this sore was the result of contagion. Yet it is barely a year since he had his first chancre, and this was followed by an eruption, of which he had scarcely got clear when this second sore occurred. The case is proof that a man may have an indurated sore on the penis within a year of a former one, but it is not proof that he may have syphilis twice, for this patient has not as yet had any constitutional symptoms as the result of the last chancre. If, however, you ask me for an answer to the general question, Can a man have true complete syphilis twice? then I must reply clearly that he can. Such cases are rare—as rare, perhaps, as examples of second attacks of small-pox—but they do occur. I am at present attending a gentleman who has a terrible phagedenic chancre and rupial eruption, and who unquestionably had complete syphilis, chancre, sore-throat, and rash, seven years ago. I have also a second case under care, very much milder, but illustrating exactly the same fact, with almost precisely similar dates. Second chancres are, however, far more common than second attacks of constitutional syphilis. Many of them are the result of fresh contagion, but seem to have no power to produce constitutional symptoms; but others are not from contagion at all, but form in connection with a taint still remaining from the first attack. It is a most important fact that indurations may form in the penis in every respect exactly like Hunterian chancres, not distinguishable in any way, and yet that they may be merely recurrent sores, and the products of constitutional taint. I have seen this over and over again; and M. Alfred Fournier of the St. Louis Hospital has written a very instructive paper on this form of sore. In the case of our patient, it is obviously impossible to say, after the statement which I have just made, whether or not his present sore is the result of fresh contagion. It may be simply a relapse, or it may be a gumma. He, however, confesses to exposure; and, as the sore followed in due course, it is probably true that he was afresh inoculated. Second attacks of syphilis are sometimes, as in the case just mentioned, very severe. The same has, I believe, been occasionally noted in recurrent attacks of variola. As a rule, however, they are mild, or even abortive. Third attacks may even occur; and so may, as we are told, third attacks of small-pox. We must explain such facts, I expect, by reference to individual peculiarity and idiosyncrasy, but it is important that they should be known. The belief that syphilis can occur but once in a lifetime is very widely spread amongst a certain class of the public. I have watched with amusement the change in expression in many a young gentleman's face when he got my reply to his smiling suggestion—"A man cannot, I suppose, have the disease a second time?"

TREATMENT OF LICHEN PSORIASIS (LICHEN RUBER).

We discharged recently from Sophia ward a middle-aged woman, who was the subject of lichen psoriasis. As I explained at her bedside, I much prefer this name to either of the others by which this disease is known. As you know, it has been named lichen ruber by some, and lichen planus by others. It is, however, essentially a form of psoriasis. It occurs to the same class of subjects, is curable by precisely the same means, and, like psoriasis, is liable to relapse or to recur after considerable periods of health. The case which we have just been studying was of much interest in reference to the points to which I have adverted. Although it certainly was an example of the malady known as lichen ruber, yet in parts the eruption was not distinguishable from common psoriasis. It conformed to the lichen type in that it began in little papules, which occurred in groups; and, when a patch was formed, it was by the coalescence of a number of small papules. This mode of spreading is, perhaps, the chief feature

of distinction between the malady in question and common psoriasis. The latter begins as a point, which, spreading at its edge, becomes a papule, which, again enlarged at its border, becomes a patch, possibly a very large one. Thus, psoriasis patches are always almost round, nummular, *i.e.*, like coins or rings, whilst those of lichen ruber are irregular, in lines or particles. In the case in question, most of the eruption was arranged in this manner, but some patches were not. On the elbow-tips and over the ulnæ were patches which, in mode of formation and in accumulation of scales, could not be distinguished from common psoriasis. Our treatment of the case was exactly that of the latter disease—tar externally, and arsenic internally. In nine cases out of ten, these remedies will cure lichen psoriasis pretty quickly. Some of you may remember a man whom we had under care six months, a splendid specimen of the disease. He had been sent to me by Mr. Forshall of Highgate. It was a first attack, and occurred to a healthy young man. I prescribed arsenic and tar. Through Mr. Forshall's kindness, I had an opportunity of seeing this man again last week. He told me that about six weeks' use of the remedies quite cured him, and that he has, during the last four months, remained without treatment quite well. In our last case, however, we have not been so fortunate. Our patient was of very peculiar nervous system, in fact almost insane, and the influence of arsenic appeared to be to excite her. Several times we had to suspend it on account of the irritable condition which it appeared to produce, and finally she was discharged from the hospital uncured, in consequence of the trouble which she gave in the ward. As a rule, I have found lichen psoriasis more easily influenced by treatment than common psoriasis. The cure is also usually more complete. The periods of immunity are also longer, often not less than several years; whereas psoriasis, however good the cure, usually relapses, I think, within the year.

CHRONIC SYNOVITIS, ARTHRITIS, OR STRUMA: IMPORTANCE OF THE DIAGNOSIS.

We have had lately a great many cases of synovitis of the knee-joint. I think you will have observed that, roughly, we may divide all the cases of chronic synovitis into two groups, those which are connected with struma, and those which are of an arthritic nature, in the conventional sense of that term. This division is of considerable practical value. Under the arthritic head, I comprise all that are associated with gout, rheumatism, or rheumatic gout, and all gonorrhœal rheumatism; and of all these, we may say that we expect them to get well. Sometimes there is stiffening, sometimes effusion is very long in disappearing; but still, in nearly all cases, in the end the patient again walks on the limb. It is very different with the strumous group. Here the tendency is to pulpy thickening of the synovial membrane, and to incurable conditions. It may be that destructive changes are ward off by long rest, but the patient is disabled, and the limb useless. We have half a dozen of this kind of knee now in our hands, not bad enough for amputation or excision, but still so bad as to prevent walking. In these cases, we are obliged to forbid walking, whereas in most of the arthritic cases, unless exercise causes pain, it may be permitted with impunity. A considerable variety of conditions is presented in this group, and especially is the arthritic process modified by the age of the patient. The older the patient, the more chronic and the less painful is rheumatism. You know that I am in the habit of insisting upon the importance of the patient's diathesis, even in cases of synovitis which is called traumatic. We admit a great many cases in which free synovial effusion has followed a sprain or contusion. In these cases, if the effusion lasts long, or if it is in excess of what its supposed cause will account for, you must suspect the arthritic diathesis. The patient is rheumatic or gouty. We have had numberless illustrations of this. Sometimes it is difficult to get at the exact facts. In the case of a man who has just left us, the synovitis persisted in spite of treatment, and relapsed after an apparent cure. It appeared likely that the case might end as hydrops articuli. I had repeatedly taxed the man with being gouty, but we could get but little evidence. Last week, his employer called on me; I then learned that the man had been for thirty years employed as a bottler in wine vaults, and that his habits of free wine drinking had often nearly cost him his place. I was told that no objection was made to a bottler drinking as much wine as was good for him, and that complaint only resulted when so much was taken as to interfere with his efficiency as a workman. It is not easy to imagine a position more likely to produce a gouty state of system. We have since let this patient leave the hospital, supplied with a knee-cap. He still has some fluid in the joint, but he can walk without any pain. Exercise, which would of course be most injurious if the disease were strumous, will not hurt him.

Dr. BEAMISH, late medical officer of the City Prisons, Cork, has been granted a retiring allowance of £110 *per annum*.

A FEW REMARKS ON YELLOW FEVER, AND ON THE MOVEMENT OF TROOPS.

BY SURGEON-MAJOR J. P. H. BOILEAU, B.A., M.D.,
Diplomate in State Medicine, University, Dublin, etc.

THE return of the 1st Royal Lancaster Regiment (1st 4th Foot) from the West Indies, has recently been made the subject of a short paragraph in a contemporary. The regiment arrived in England in November last, having been ordered home from Barbadoes, on account of the prevalence of yellow fever at that station, and in consequence of the officers and men of the corps having suffered very severely from the disease. The paragraph states, that it was anticipated that a regiment from Malta would be ordered to Barbadoes as soon as the local authorities considered the epidemic to have passed off, and after the barracks had been thoroughly disinfected.

It seems to me, judging from the lessons of the past, that the authorities will not hastily order a regiment from Malta to Barbadoes under present circumstances. I will narrate a few cases which seem to justify me in forming this opinion; and first, that of the 84th foot, in 1867.

I served in the Malta command during the years the 84th was stationed in the island—1865-67. The first year was that of the great cholera epidemic there, and in that year sickness and mortality were very high in the 84th; during the year 1866 continued fevers were much more prevalent and fatal than usual in the command. The 84th left Malta early in 1867, and landed in Jamaica in April; it was immediately marched up to the hill station of Newcastle; in May, 18 cases of fever were admitted to hospital; in June suspicion arose as to the true nature of the disease, and before the end of the month a case proved fatal. The disease—yellow fever—spread, and the regiment lost 13 men in July, 13 in August, and subsequently 5 more. Further particulars will be found in the 9th Volume of the *Army Medical Department Reports*.

The 1st-4th Foot, which has returned from Barbadoes, arrived there in January, 1879, and like the 84th, it came from the Mediterranean, from Gibraltar, where it had been stationed upwards of four years. The sad experience of this regiment in Barbadoes is too fresh in the minds of your readers, to need dwelling on here.

The 2nd-3rd Foot arrived in the West Indies in July 1864, also from Gibraltar. Soon after its arrival it suffered from an epidemic of remittent fever, which was the cause of 182 admissions into hospital in November, and 133 in December. Officers, men, women, and children were alike subject to it. The Royal Artillery also were affected by the epidemic, "but by no means to so great an extent" as the 3rd regiment.—Vide *Army Medical Department Reports*, Volume 6.

The 1st-21st Foot went from Malta to the West Indies in April, 1860. During the nine months of 1860 that it was in Barbadoes, 84 cases of remittent fever were admitted, and two died; and it is worthy of notice that the year before the regiment left Malta, it was exceptionally healthy; for whilst the admissions for fever were "exceedingly numerous" in other regiments, the 1-21st was "the only corps which appeared to have enjoyed a marked exemption from the disease."

Now, contrast the above cases with that of the 29th Foot, which after two years in Malta, went to Canada in July, 1867, and thence, in December, 1869, to the West Indies, (relieving the 84th). I was then the assistant-surgeon of the 29th, and served with it continuously for more than three years in Jamaica and in Barbadoes. I well remember that, as we lay off Port Royal, in view of the dreaded palisades, we contemplated our future abode on the distant hills, between three and four thousand feet above us, with many and with mingled feelings, for *inter alia*, had we not the unfortunate experience of the regiment we were about to relieve as a subject for our reflections? Yet, as the sequel proved, we need not have troubled ourselves with a single gloomy foreboding; for the fates were singularly propitious to us. During the three years and three months that I served with the 29th in the West Indies, I attended altogether but two fatal cases (mitral valve disease, the sequel of rheumatic fever, and a case of acute hæmorrhagic phthisis in a young recruit). We lost not a single man, woman, or child from any endemic or epidemic disease; nor indeed from any zymotic disease whatever, either at Newcastle, Jamaica, or at Barbadoes, nor did we invalid any, to the best of my recollection.

The 98th Foot relieved the 29th in the West Indies, in March 1873. It came from Bermuda, and its medical history, in the West Indian command may be found in the *Army Medical Department Reports*, volume 15, page 77, and volume 16 page 75, etc. Suffice it to say here, that the sun of health which had shone so brightly over the 29th, with