amongst them one or two women who had signed antivivisection petitions. I did not stop to inquire if they were willing to be saved by practice based upon vivisection; but I should hope to be forgiven. The means by which the necessary knowledge was obtained may, in their particular cases, be condoned by the end to which that knowledge was applied.

I have had the rare opportunity of treating a case of strychnine poisoning. I was called to a gentleman who had inadvertently swallowed a poisonous dose. I saw him within a short time of the accident. He was in the most violent tetanic spasms; opisthotonos was so marked that his body was arched back, and respiration was nearly suspended. Pending the fetching of the nitrite of amyl, the fits recurred at short intervals with unabated energy. I obtained the assistance of a young medical friend, who sat by the bedside, diligently watching, and making the patient inhale nitrite of amyl the moment the premonitory twitchings or facial expression appeared, and always with the effect of averting or greatly moderating the fits; and, to make the evidence complete, when the warning was not seized in time, the fit appeared in nearly its original intensity. This treatment, continued during sixteen hours, resulted in the recovery of the patient. It was impossible to determine with precision the dose taken. We had only the physiological test to satisfy us that he had taken enough to destroy life.

The spasm-subduing virtue of nitrite of amyl has lately been most happily illustrated, in a case of *enclitonnement* of the placenta by hour-glass contraction, by Dr. Fancourt Barnes. This case was published in the JOURNAL for March 18th, page 377.

ACUTE TRAUMATIC MALIGNANCY.

By HENRY B. WALKER, L.R.C.P. & S.ED., Lowestoft.

SOME cases seeming to confirm the theory implied in the heading of Mr. Barwell's paper in a recent number of the JOURNAL, have fallen under my notice, and may be interesting.

CASE I. Mr. H., a master mariner, aged 53, was always healthy until fourteen weeks before death, when he was struck violently in the right hypochondrium by the shaft of a cart. The blow felled him, rendering him insensible, and it was some time before he could be removed to his own house. Here he was laid up for two days, and, though feeling "very sore and ill", he then started in charge of his ship on a trip to Sunderland. During the voyage, he was in much pain, and on his return to Lowestoft was again laid up at home; feeling better after some rest, he took another voyage, but was so ill that he had practically to give over the command to his chief officer, and when he reached port, to place himself under a doctor's hands. His symptoms were frequent vomiting, complete loss of appetite, and intense pain in the region of the liver, shooting across the epigastrium. He was much bruised, and very tender. I saw him on his return to Lowestoft, and he was then very ill, with high temperature, thirst, constant vomiting, and wasting. On examining the abdomen, I found there was some remaining sign of the bruise—discoloration, etc.—and he was extremely tender all over the region of the liver in front, the tenderness extending to the left hypochondrium and down nearly to the navel. I could not detect any sign of enlarged liver, but there was some jaundice, with pale stools and constipation. A fortnight later, his condition becoming gradually worse, a well defined hard and tender bossy mass with rounded edges, under which the fingers could be passed, was found projecting below the ribs (at a point which would be intersected by a line drawn perpendicularly from the anterior superior spinous process of the ilium), and extending well into the left hypochondrium. This mass, which was taken to be liver, extended downwards and to the left side, till it filled the abdomen in front to a level with the umbilicus. Its extreme hardness, and the constant presence of sharp lancinating pain, together with the cachexia accompanying it, led to a diagnosis of malignancy, which was confirmed by an eminent consultant. Towards the last week of his life, the patient vomited a quantity of blood, containing pyriform and oval multinucleated cells. After the last hæmorrhage, the patient died exhausted.

I could get no *post mortem* examination, but there was no doubt in the minds of three medical men as to the nature of the disease, and it was believed to be carcinoma of the liver, invading the stomach secondarily. There was no history of malignancy in the family, though several members had died about the same age as my patient. No history of dysentery or syphilis was attainable. I may state that both the patient and his friends assigned the accident as the cause of his illness, and certainly facts seem to warrant them in doing so. He certainly never recovered the effects of his bruise, had not complained before, and died fourteen weeks after he received it.

CASE II. Mr. F., a farmer, was out shooting; and, while he was trying to force his way through a hedge, a thorn penetrated his trousers and wounded his prepuce, causing blood to flow. He took little heed of it at the time; but a few days later, feeling some heat and uneasiness, he found swelling and irritation about the wound. This rapidly increased, and he then consulted his medical man, who advised certain remedies. The penis grew worse, and the whole glans presented a mass of carbuncular hardness in a short time, and was very painful. An incision into it giving no relief, and suspicion being confirmed on consultation, the penis was amputated, and the growth proved to be an epithelioma. This was only a few weeks from the date of injury.

CASE III. A gentleman rode to hounds, mounted on a very broad-backed horse, and in the evening felt the inner side of his left thigh stiff and sore. The stiffness persisted, and was followed by a slight fulness, and finally by a well defined tumour about the size of an orange, nestling amongst the deep muscles of the thigh. This was removed by operation, and proved to be a small spindle-celled sarcoma.

CASE IV was related to me by a lady whom I was attending in labour. Her brother was playing football, when he received a kick on the shin. This led to severe inflammation, which laid him up for some time. On apparent recovery, a hard swelling remained on the tibia, which soon began to cause pain and enlarge. As it increased and bore suspicious characters, the limb was amputated at the knee-joint, but the growth recurred, and the patient lost his life. I asked the lady what the growth was, and she said, "The doctors did not call it a cancer, but by a queer name—'sarcoma'".

In all the foregoing cases, the patients confidently referred to their injury as the cause of their trouble; and in all except the last, I can state that minute inquiry failed to trace a family history of malignancy. In the last case, I have no note on this subject, and therefore cannot speak with certainty, but I believe I was assured that such was the case in this one also.

A DISPLACED LEUKÆMIC SPLEEN, CAUSING OBSTRUCTION AND DEATH.

By M. COLLINS, M.D., Scarborough.

ON February 15th, I received a message to "come at once and attend K.P., who was in labour." I found the patient, aged 36, very much emaciated, and presenting a peculiar but very suggestive waxy appearance. A quick and feeble pulse, dyspnoea and urgent fecal vomiting, had set in already. The os was patulous, so as to admit the examining finger readily: the posterior lip was hard and nodulated. Otherwise the uterus was in its normal unimpregnated condition.

On inspection, the abdomen appeared to be as large as at the sixth month of utero-gestation; but by manipulation, I could detect a solid mass in the left inguinal, and half the left lumbar regions. Its upper margin was sharp and well defined, corresponding to a line drawn from the umbilicus to the crest of the ilium. The right lumbar and inguinal regions were tympanitic. The liver was visibly enlarged, projecting below the ribs, and filling up the left hypochondriac region.

That there was neither labour nor pregnancy, was manifest; and it was equally evident that the patient was *in articulo mortis*. As the hand of death was upon her, I deemed all operative interference imprudent. Enemata, draughts, etc., were administered, but in vain. She died within forty-eight hours.

The patient died, as she had lived, in the belief that she was *enccinte*; for the patulous condition of the os, pain, a gradually increasing abdominal tumour, etc., led a certified nurse to an erroneous diagnosis. I could not dissuade either from their mistake. The nurse stoutly and rudely maintained that the patient was *de facto* in labour.

Post Mortem Examination.—On opening into the abdomen, I found the spleen filling up the left inguinal and half the left lumbar regions. Its internal surface was uppermost, and its external undermost. The gastro-splenic and suspensory ligaments were elongated. The spleen weighed two pounds and eleven ounces. Its external surface was traversed from above downwards by a well marked fissure, which gave the organ a bilobed appearance. On section, considerable resistance was offered to the knife by a thickened capsule and trabeculae: it presented a bloodless, homogeneous, reddish-brown appearance. The intestines and omentum were pushed to the right side, and those lying contiguous to the spleen were dark and considerably congested. Part of the descending colon and sigmoid flexure were pressed upon; and the latter was atrophied to about the size and thickness of the finger of a glove. There were some seybala in the colon. The uterus was normal, with the exception of a patulous os and nodulated posterior lip. A small abscess, nearly as large as a pigeon's egg, was found in the left broad ligament. It contained about a drachm of horribly offen-