LABOUR IN A PRIMIPARA COMPLICATED BY A LARGE URETHRAL CYST AND BY PROLAPSUS UTERI :

DELIVERY BY CEPHALOTRIPSY: SUBSEQUENT OPERATION AND
RECOVERY.

By J. HICKINBOTHAM, M.D., M.R.C.P., F.R.C.S. Ed.,
Physician to the Birmingham and Midland Hospital for Women.

ON November 12th, 1880, in consultation with Dr. Young of Erdington, I saw Mrs. S., aged 21, who was daily expecting her first confinement. An examination revealed a huge mass protruding from the vulva, which consisted of a greatly hypertrophied cervix; and in front of it a large soft tumour, which I took for a cystocele. Behind the prolapsed mass, and lying in a flaccid condition against the posterior vaginal wall, was a perfect hymen of the typical sickle shape. The pelvis seemed of average size, and there was nothing in the general state of the patient to cause anxiety. The os uteri easily admitted a finger; and with some trouble (from the extreme length of the cervix) I was able to recognise the head presenting. The membranes were entire. After consideration, it was determined to wait and watch until labour should come on, in the hope that gradual dilatation would occur, and the delivery end naturally. Labour began on the evening of the 21st, and, although the pains were strong and regular, no advance was made; therefore Dr. Young asked me to see her again at 11 a.m. the next day.

At this time the os was but little larger than above described, and presented a very peculiar appearance, being perfectly annular, and surrounded by a ring of white, shining, gristly-looking fibres. Immediately above this, the cervix was encircled by large veins, with dilations almost hæmorrhoidal in their character. The membranes had ruptured some hours previously, and the head could be felt at the brim, but was not affected by the pains, which were of moderate length and force.

As the patient showed signs of exhaustion, it was decided to attempt delivery without further delay; she was, therefore, chloroformed, the bladder emptied, and with great caution I endeavoured to overcome the rigidly contracted os. After considerable effort, I was able to get in three fingers, when a smart gush of blood warned us that the vascularity of the cervix had not been over-estimated.

Two courses were now open to us: the first and easiest was to make two or three small incisions, and then deliver by forceps; the second was to treat the constriction as being insusceptible of dilatation, and to deliver by lessening the volume of the foetal head. After earnest consideration, we decided upon the latter alternative; because, although in ordinary cases the incisions through the os contract so much as to be unimportant, and the risk from hæmorrhage is, therefore, small, yet in the present instance, from the altered condition of the tissues, and especially on account of the great vascular turgescence, we felt that the danger would be very considerable. The cranium was, therefore, perforated and crushed by a small cephalotribe, after which there was no difficulty. She went on well; but the tumour anterior to the cervix began to increase, and there was some pain. On the tenth day it ruptured, and there was a free discharge of puriform fluid, after which convalescence was uninterrupted. Neither during pregnancy nor after labour was there any incontinence of urine, or other trouble connected with the bladder.

The prolapse of the uterus had occurred quite gradually, so gradually, indeed, as to push aside the hymen without rupturing it, and so entirely without inconvenience that the poor woman did not know until her marriage that "she was different from other women." Her menstruation was always regular, and her health had always been robust. The cause of the procidentia was, no doubt, lifting and carrying heavy weights whilst very young. Six weeks after her confinement she called upon me, at Dr. Young's request. I found the womb still prolapsed, but the condition of the os and cervix was nearly natural. The tumour which I have described lay in a flaccid state upon the anterior vaginal wall, or rather hung from it, and a large hole, admitting my finger, still remained. A sound passed into the bladder showed nothing abnormal, and she could hold her urine from morning to night.

When the cyst was itself examined, by the finger or sound, I could merely make out a large empty cavity between the bladder and uterus, and apparently communicating with nothing.

She was afterwards admitted into the Women's Hospital, under my colleague, Mr. Lawson Tait, who has kindly given me the following report of her then condition, and the treatment adopted.

"On examining the patient, there appeared, protruding from the vulva, what looked exactly like a cystocele, but which had the singularity of having a hole, about the size of a shilling, punched (as it were) into its most prominent part. The finger was admitted into a large pear-shaped cavity, having its apex directed up towards the urethra; and it was ascertained that it opened into this canal by a small aperture, not larger than a hemp-seed. When the bladder was injected full of milk, and the external meatus forcibly closed, some of the milk found its way into the sac. The cyst ran backwards as far as the cervix uteri, and seemed to double up on each side of this structure into small pockets. I laid the whole cavity open from the urethra to the cervix, dissected out the mucous membrane over the whole surface, and shortened the remaining flaps of the vaginal surface, so that they met correctly in the middle line, when united by suture.

"The removal of the cyst-wall from the cervix was difficult, and the hæmorrhage was profuse and troublesome. The patient made a perfectly uninterrupted recovery, and all traces of the abnormal cyst have disappeared, and the protrusion is cured.

"I have no doubt this case is an illustration of the rare form of congenital malformation, which, I believe, I was the first to describe in the *Lancet* of January 9th, 1875. I have seen three such cases, and in two of them the distress of the patients was very great, on account of the decomposition of the urine escaping into the sac, and there becoming ammoniacal and purulent. In the present instance, this was obviated by the hole in the depending part of the cyst; but it is perfectly certain that urine in considerable quantity must have passed through the cyst.

"The size of the sac in this case was probably as great as that of the bladder, and it was probably five times as large as in either of the other cases upon which I have operated. In all three cases I dissected the cyst out, and found the aperture into the urethra to be very small, being, as I have said, about the size of a hemp-seed.

"I think the relations of the cyst, in Dr. Hickinbotham's case, prove its origin to have been congenital, as it is perfectly impossible to imagine that a cyst could develop itself from any structure lying between the urethra and the cervix uteri, so as to have relations so intimate with both, as existed in this case. In all three cases the cure by operation has been complete."

I may just remark, upon this note of Mr. Lawson Tait, that his hypothesis that the absence of symptoms was due to the hole in the cyst, could only be correct after the labour, since before that there was no hole. I think a more likely explanation is, that the opening into the urethra was protected in some way by a valvular fold of the mucous membrane, which was sufficient, until there was obstruction in front, and more pressure than usual from the bladder. These conditions no doubt obtained during the labour, and then for the first time urine found its way into the cyst, whose contents had previously been unirritating and harmless.

Since the above was written, the patient presented herself at the hospital, complaining of a "lump in her stomach which gave her great pain." Her temperature was 102°, and pulse quick. She was therefore admitted, and examination showed a tender tumour above and to the right of the umbilicus, which I believed to be a movable kidney. A day or two of entire rest in bed seemed to set matters right; the tenderness and fever disappeared, and she was again sent home.

A CASE OF LABIO-GLOSSO-LARYNGEAL PARALYSIS.

By J. M. HOBSON, M.D., Croydon.

IN the JOURNAL for December 3rd, 1881, a case of "Brain-Disease simulating Bulbar Paralysis" is reported as brought before the Manchester Medical Society by Dr. Ross of that city. The primary disease does not appear to have been in the medulla oblongata, but in the central ganglia, and on the left side, "a streak of secondary degeneration" was traced from the diseased corpus striatum (the right was also affected), "through the crus, pons, and anterior pyramid of the medulla oblongata." Moreover, "these degenerated fibres occupied the internal margin of the anterior pyramid of the medulla, and appeared to bend back in the median raphe to reach the bulbar nuclei of the opposite side." It is to be noted that this man had an attack of hemiplegia and aphasia eighteen months previously.

This striking and, as far as I know, unique case, set me thinking whether one which came under my observation in the early part of last year, and which till now I have associated in my mind with those cases described by Dr. Wilks (*Diseases of the Nervous System*), some of which had hæmorrhage into the medulla oblongata, ought, perhaps, after all,

to be classed with Dr. Ross's case. I therefore bring my case before the readers of the JOURNAL for comparison with this other one, although, in one sense, they cannot be compared, as my case is still living, and, unless she gets another "stroke", is likely to live for years to come.

Mrs. —, aged about 40, travelled, on February 6th, a distance of over 130 miles to stay with a relative in my neighbourhood. In the middle of that night, I was called to see her, and found her in the following condition: quite speechless, though fully conscious; no marked hemiplegia; lower jaw clenched; no cardiac symptoms; radial artery thickened. Next day, the jaw was more opened, and it was then discovered that the tongue was almost powerless, and that deglutition was difficult. She could scrawl her name with her right hand, but she held the pencil badly. There was indication of slight facial palsy on the left side of the mouth. The optic discs were normal. On inquiry from her friends, and on writing to her medical man at home, I ascertained the following history.

She had good health up to a year before, then had slight and transient loss of speech. Her next attack was on one night in January 1881, when she lost her speech for a very short time. Again, a third time, on January 16th, she had partial aphasia, accompanied now with partial left hemiplegia and impaired deglutition; these all coming on gradually in the course of the day. In about a week, she got better of this, and when she arrived here was very nearly recovered. She could talk and walk quite well, and could use her arms, only the left was a little impaired in its movements.

On the third day, February 8th, she could swallow better, and could speak a little. Albumen was found in the urine. On the 9th, she was not so well. On February 11th, Dr. Hilton Fagge saw the patient with me in consultation. At this time, complete facial paralysis was noted. This paralysis remained, as far as voluntary movements were concerned, during the remainder of the time the patient was under my care. On the other hand, *blinking* occurred; and, when amused, the face would be drawn up into an expression of laughter, but never into a smile. When an effort was made to close the eyes, a wide staring only was produced. I noted on this day that the *right* hand had been weaker for a day or two. It was the *left* which had been affected before. On the 19th, she had remained about the same for the last week. On the 24th, she had some power in the right arm, and swallowed better. She had been complaining for some time past of pain in the masseters—probably spasm. When tested with carved letters, she could use them well to form words. On March 21st, the right hand had gained considerable power. She was now able to dress herself. Swallowing had improved, the jaw could be more widely opened, and the pain in the masseters was gone; but there was no marked improvement in the movements of the tongue, and the voice was still absent, as it had been throughout, except on February 8th. There was, as far as I remember, one inarticulate sound for everything, and a slight sound, not sustained, on laughing. It was not clear that speech by letters was quite perfect, for, on trying to spell "cannot" on the letters (before several persons), she spelt "connot". Cutaneous sensibility had been found perfect in the face, but diminished in the right fingers.

Mrs. — now returned to her home, fortunately without any mishap this time, and passed at once under the care of her usual medical attendant, by whose courtesy I am enabled to publish these notes. This gentleman has kindly informed me of the progress of the case. On September 19th, he wrote to me: "Her general health has much improved, and she is able to engage in some of the lighter duties of her home, but she cannot utter a word;" also, "she is able to put the tongue out, which she could not move at will before, and she swallows with less difficulty." I have learned still later that swallowing is now quite easy; there is no power of speech, but some voice.

Now, the symptoms of this case much resemble those of Dr. Ross's; only his had less pronounced symptoms at the time—"loss of facial expression, imperfect articulation, and slight difficulty of deglutition"—while the previous attacks in my case were lighter. Dr. Wilks mentions a case in his book (*op. cit.*) which I well remember in the wards of Guy's Hospital, and which now reminds me of Mrs. —'s case. This woman could not utter a word; but when, apparently, pleased at something said to her, her mouth opened, her face was opened out too into a broad laugh, while one slow inspiratory noise was made. I think this woman eventually went home, and I do not know what became of her. Her case was "cerebral" for a long time before she settled into this state.

Till I saw Dr. Ross's case, I had thought of my own as one of hæmorrhage into the neighbourhood of the medulla. Nevertheless, I was at a loss how to explain the remarkable facts of the facial muscles being quite cut off from the direction of the will, but responding to the stimu-

lus of a humorous idea, and of the blinking. Now, however, I think I have a clue to the solution of that difficulty. If the lesion were of the central ganglia, the facial nuclei would certainly be cut off from the ordinary motor impulses descending from the hemispheres, but would be free to respond to impulses reaching them through the optic nerves and corpora quadrigemina. Thus the blinking would be accounted for; but how about the laughing? Though this is a respiratory act, and is probably directly regulated by the medulla, a stimulus originating above must reach the medulla first. How is this to be accomplished, if the motor tracts be interrupted? Meynert's notion (Stricker's *Histology*, vol. ii) is of a double motor tract, represented by the crusta, formed of fibres from the corpora striata, on the one hand, and by the tegmentum (or rather a portion of it), formed of fibres from the optic thalami, etc., on the other hand. Now, according to the same writer, the corpora striata receive fibres mainly from the anterior part of the brain, while the optic thalami, corpora quadrigemina, and corpora geniculata receive a portion of their fibres at least from the occipital and temporal lobes. Thus, a humorous idea,* taking its origin in the occipital convolutions as the seat of the emotions, may be reflected into this latter group of ganglia, and thence again into the various nuclei of the medulla oblongata, including those of the seventh pair, concerned in the act of laughing. It seems that the central connections of these nuclei are through both the crusta and tegmentum; those connecting the common nucleus of the sixth and seventh passing through the crusta, while those connecting the inferior nucleus of the seventh, as well as some direct central roots of the nerve, pass through the tegmentum.

One has been accustomed to think of deglutition as so perfectly reflex an act, that surprise is, at first sight, felt at having to associate its impairment with cerebral lesion; but we must remember that, in neither Dr. Ross's case nor my own, was the act deeply involved; and it was probably affected only through paralysis of the tongue, which would hamper only the first or voluntary stage of deglutition. In Dr. Ross's case, the streak of degenerated fibres was traced into the inner border of the pyramid, and then "appeared to bend back in the median raphe to reach the bulbar nuclei of the opposite side". This is the course taken by the central fibres of the hypoglossal nerve. Then the paralysis of the spinal accessory nerve, causing the aphonia in my case, could just as well be accounted for by an unusual central lesion—and we are now contemplating unusual lesions—as the paralysis of the hypoglossal nerve. Lastly, the previous attacks which I have described would receive a more probable explanation—more in accordance with experience—from the idea of central lesions than from that of repeated accidents in the medullary region.

On the other hand, if one tried to associate the symptoms in Mrs. —'s case with a lesion, or rather a series of lesions, in the neighbourhood of the medulla oblongata, the amount of disorganisation in that all-important part of the nervous fabric would be so great as to be incompatible, one would think, with the maintenance of life.

Thus, to my mind, at present, the balance of evidence is in favour of central lesions in this remarkable case. But the subject stands in need of elucidation.

ON AN "EPILEPTIC" CONDITION OF THE CENTRES WHICH CONTROL THE NUTRITION OF THE BODY.

By W. MUSSEN YOUNG, M.D.,

Assistant Medical Officer, Suffolk County Asylum.

A GREAT extension has been made of late of the class of cases which are recognised as entitled to a place under the generic heading "epilepsy". Formerly this designation was applied only to those cases in which there had existed a train of symptoms so marked and severe in character as to justify the popular appellation of "falling sickness". In order to constitute an epileptic fit, it was deemed necessary that there should be observed distinct loss of consciousness, foaming at the mouth, tonic, followed by clonic, spasms of the entire voluntary muscular system; in fact, all the symptoms which, taken collectively, are now known as *haut mal*. But it has been for some time generally admitted that in epilepsy, as in everything else, there are degrees of severity, and that it could no more be expected that every attack should be of this severe character, than that all cases of small-pox should be of the confluent form. Accordingly, attention has been directed to a class which had previously attracted little notice, owing to the slight disturbance which they cause when compared with the struggle for life which is witnessed in a seizure of the more severe type. In this class—*petit mal*—would be included cases in which the symptoms are so slight as

* One should, perhaps, rather say, the cerebral process of a humorous idea.