such ulceration could not be imagined, and few hospitals would entertain the idea of receiving him. He was, however, admitted at the request of a very valued friend of the institution, who represented that he had come a long way from home in the hope of relief. Strapping being plainly out of the question, from the size of the ulcers and the low vitality of the surrounding skin, I grafted the entire surfaces with layers of sponge. This process went on in the usual manner, and left a healthy surface; the granulations of which, however, soon died away, and could not be kept up. Skin-grafting failed utterly.

We had now come nearly to the end of April, and our failure in this case was as complete as that of our sister hospital. It appeared to me that now was the time to try the galium aparine, which was beginning to peep out in all the hedgerows about Dublin. Here I must tender my acknowledgments of the zeal and energy of the members of my clinical class, who were untiring in their efforts to collect this herb, which was not to be had of any of the herbalists. An ample supply for this and other less severe cases has since been kept up, and it has been used with the most marked success in the following manner.

Grasping in the left hand a bundle of ten or twelve stalks, with a scissors held in the right hand, the bundle is cut into junks about half an inch long. These are thrown into a mortar, and pounded into a paste. This paste, which has an acrid taste and slightly acrid smell, is made up into a large poultice, applied to the ulcer, and secured with a bandage. It is renewed three times a day. Its action appears to be a slight steady stimulant, and powerful promoter of healthy granulation. Its effect in this most unhelpful case was decisive and plain to all. Healthy action ensued, and has since steadily continued; and, after a month of treatment, both ulcers have been reduced to considerably less than half their original size. If this action continue, which I have no reason to doubt, the cure will be accomplished within a measurable and short period. The patient is in the ward, and anyone can see the great amount of new dermatisation which has been effected during the month.

I could give several other cases not so striking as the above, but it would be mere repetition. Of one, however, I would wish to make brief mention.

Mary G., aged 34, was in a very advanced stage of pulmonary consumption, with great wasting and emaciation. Severe lesions of the left lung compelled her to lie always on her right side; and, as a result, she got bed-sores on the right shoulder and on the right trochanter. In addition to a large water-bed, I dressed both bed-sores with the galium pulp. The application was found most grateful and soothing, and rapidly healed the bed-sores. She is still alive, though passing rapidly away. She does not, however, suffer from the sores; and I have had some other instances of the same kind. As far as chronic ulcers are concerned, the application suits best in the indolent or in the healthy states. In irritable ulcers, its stimulating property causes pain. In such cases, it is necessary to reduce the irritability by poultices, iodoform, or other well-known remedies, before beginning the galium.

My reason for putting forward this remedy so soon is, that now is the time to try it. It is growing freely in almost every hedge, and can be got in any quantity during the rest of the season.

A difficulty at once suggests itself as to its general employment; viz., that in winter and spring it is not to be had at all. It appears to me that this difficulty can be effectually met by the method of ensilage, by means of which green food for cattle has for the last few years been kept perfectly sweet and fresh by burying it in silos under the ground. This plan is generally known, but all particulars about it can be learned in the pamphlet of Mr. Thomas Christy, F.L.S. (Christy and Co., 155, Fenchurch Street, London, E.C.). In the case of the galium, the process would consist of cutting the herb very fine, ramming it down by screw-pressure into a glazed earthenware jar with an air-tight cover, and burying it in the ground. Thus secured from air, moisture, and heat, it would be likely to keep through the winter. One of my pupils, Mr. M. Pierce, has already laid it thus down, and will report the result to me.* This plan, if successful, might be extended to other pharmaceutical herbs; for I have always had the idea that green herbs are more powerful than dried ones. Indeed, the late Mr. Donovan of this city used to maintain that, to make tincture of digitalis properly, the alcohol should be brought to where the foxglove was growing, and the live plant plunged into it.

Many virtues are attributed to the galium by old writers; but I

* The first jar was taken up on June 7th, after being buried for one month. The pulp was perfectly fresh and sweet.

have not been able to find any allusion to its employment in the treatment of ulcers. Linnæus mentions that in Sweden the shepherds use its stalks for straining milk—a fact also stated by Dioscorides. Its roots form a nice red dye. These are the only uses that I have been able to ascertain in regard to it in the books at my disposal.

CASE OF PERICARDITIS, WITH EFFUSION.

By JOHN W. MARTIN, M.D., Sheffield.

THE following case is a good example of the insidious and unsuspected onset of pericarditis, accompanied by effusions into the pericardial sac.

CASE. Insidious Onset: Slight Preceding Muscular Rheumatism in the Right Arm and Shoulder: Dyspnoea: Orthopnoea: Assigned Cause, the House Cellar-Floor constantly covered with Water: Patient not Intemperate, but accustomed to take Five or Six Glasses of Beer, and One or Two of Spirits, a day: Treatment: Result.—H. G. B., aged 29, residing in Sheffield, formerly a wine and spirit merchant, and, for the three months previously to my seeing him, occupied as traveller for a brewery, came under my care on January 5th, 1883. He was a large, well made, well developed, and well nourished man. He had always been accustomed to live well, and to take beer and spirits, but never to actual excess, as far as intemperance was concerned. For the time he had been engaged in travelling, partly in the way of business, he had been taking from five to six glasses of beer, and one or two glasses of spirits, on an average, a day. There was no family history of rheumatism, as far as the patient knew. He had always been strong and healthy. He changed his residence about three weeks previously to my visit. His former house was very damp, the floor of the cellar being almost always three or four inches under water, no drains existing to carry the water off. Before leaving it, he felt the present attack commence, the only symptoms being a dry troublesome cough and dyspnoea. Before the onset of these symptoms, he had slight pains in the muscles of the right shoulder and arm, but only very slight indeed, and attributed by him to holding his whip when driving. These pains were intermittent. He never had any joint-affection. The dyspnoea gradually increased to orthopnoea. For the ten days before my seeing him, he had not been able to lie down; and, for the last four or five nights, he could only get snatches of sleep of from twenty minutes to half an hour at the time.

When seen, his face presented a dusky, bloated, and congested appearance; the conjunctivæ were injected, causing the eyes to look red and ferreted. The veins of the neck were full and prominent, not extraordinarily so, but to a noticeable degree. The expression of the face was anxious and distressed. He complained chiefly of the shortness of his breath, and the troublesomeness of his cough, but did not complain of any actual pain. He seemed to think that his attack was one of bronchitis and congestion of the lungs, and had been diligently taking a mixture from a herbalist in Sheffield, each dose of which sickened him considerably, and gave him considerable distress, without proving in the slightest degree beneficial to the symptoms in the way of giving relief. Careful percussion and auscultation of the lungs failed to elicit any positive symptom of their being engaged. At the same time, so well nourished was the patient, and so abundant was his covering of adipose tissue, that I would not like to be positive as to there being no deeply seated congestion; but I could obtain no exact evidence of it.

Examining into the condition of the heart, I found slight tenderness, increased by percussion over it. The impulse was very weak, but there was no displacement of the apex-beat. The area of cardiac dullness was largely increased, measuring five inches in the vertical direction from the upper border of the fourth rib on the left side, and six and a half inches in the horizontal, from the midsternum, over towards the left side, on a level with the sixth rib. There was no thrill felt on placing the hand over the heart. The first sound could not be heard; the second was heard very faintly towards the base of the heart, on a level with the second intercostal space.

Over the whole surface of the heart, and well round into the axilla, a well marked double friction-murmur was heard, faint towards the apex, loud and rasping at the base. I could not detect any endocardial murmur. The pulse was 104, weak, with a slight thrill in it; no irregularity or intermittency. Temperature 98.4°; tongue clean; liver normal; bowels regular; felt a desire for food, but had an uncomfortable sense of fullness after meals. He was passing thirty ounces of urine daily, highly coloured, at present clear, but a little time before depositing lithates freely. It contained no sugar nor albumen. No casts

epithelium, or blood-cells, could be found. He complained of pain in the lumbar region. The treatment first ordered was the application of a blister six by six inches over the heart, for six to seven hours, and the following mixture: R. Potassæ bicarbon. ʒij; potassii iodidi ʒj; tincturæ digitalis ʒijss; spiritûs ammon. aromat. ʒss; syrupi zingib. ʒj; aquæ chloroformi q.s. ad ʒviij. M. An ounce to be taken four times a day.

January 6th. The blister rose well. He was much relieved, and able to lie down. He slept fairly, as well as the pain of the blister would allow. Breathing was much easier. The pulse was stronger, 104°. The expression of the eyes and face was much improved. The conjunctivæ were clear. The congestion of the face was gone. The blistered surface would not permit a close examination into the condition of the heart. The treatment was continued.

January 8th. Improvement continued. The blistered surface was healing, but not sufficiently healed to permit percussion. The heart-sounds were reappearing, and the friction-murmurs were decidedly diminished in intensity. Tongue clean. Pulse stronger, 100; no thrill was felt. Breathing was easier. The treatment was continued, and he was ordered to take a Seidlitz powder to regulate the bowels.

January 9th. He slept well through the night, and felt thoroughly rested this morning; he was much improved. The cough was less troublesome; some slight expectoration accompanied it. The area of dulness was decidedly diminished. The friction-murmurs were not now heard in the axilla. They were confined to a space between the sternum and left nipple. These murmurs were heard very loud and rasping, as high as the upper border of the second rib. The first sound of the heart was again plainly, though faintly, heard. The second sound was stronger. I could not, owing to the friction-sounds, determine the existence or absence of endocardial murmurs. He was passing an abundance of urine, high coloured, and depositing lithates freely; specific gravity 1030; no albumen. His appetite was improving rapidly.

January 11th. I happened to call at dinner-time, and found him eating stewed rabbit with the heartiest of appetites, and relishing it thoroughly. His face and eyes and look were clearer and better than I had seen them for a long time. The veins of the neck were no longer visible. He expressed himself as feeling better than he had felt for a long time. The cough improved. There was no sense of dyspnoea nor pain. He slept well. He was passing an abundance of urine, and the bowels were acting regularly. There was no tenderness over the heart. Dulness was reduced to three inches in the vertical, and four and a half in the horizontal direction. The superior margin of dulness was now on a level with the middle of the fifth rib. The external margin of dulness was a little outside the left nipple-line. The heart's impulse was strong and diffused. The first sound was distinct, but not so distinct as in health. A single light brushing friction-murmur was heard over the apex, not confined to any single place or direction. Towards the base, the double friction-murmur noticed before was heard diffused over a very large area. These murmurs were greatly diminished in intensity all over, and seem inclined to disappear. They were heard faintly in the axilla. The pulse was taken in the standing position, and after the examination, which may account for its being 104; it was full and strong. The treatment was continued. His wife to-day told me she thought this attack had been stealing on him for some time, and that she had noticed him not to be quite himself for the past three months.

January 13th. The patient looked well, and expressed himself as quite ready for work. The eyes and the expression of the face were clear and healthy. He was free from all pain and tenderness. He slept and ate well. The bowels were regular. He passed plenty of urine, clear, and free from deposit. The dulness over the pericardial region remained unchanged. The heart's impulse was stronger, but much diffused. The friction-murmur remained as last described, but was heard round into the axilla, and faintly in the upper portion of the interscapular space on the back. The pulse was 104, unchanged. No sense of dyspnoea was left.

REMARKS.—The foregoing, though by no means an uncommon case, is, I think, not altogether devoid of interest, or unworthy of notice, on account of the insidious character of the onset of the heart-affection; so much so, that the patient's attention was never once directed to the heart as the seat of mischief. If there had been any pyrexia, it had disappeared when I saw him. The case affords another good example of what seems to be pretty generally admitted, that the serous linings to the visceral cavities, and the envelopes to their contents, must be regarded in the light of lining membranes of joints; and that, in persons affected with any of the

leading cachectic taints—gout, rheumatism, tubercle, or struma—these membranes may become affected primarily, and be the first, and indeed the sole manifestation of the constitutional diathesis. This view reduces the term "idiopathic" pericarditis, pleuritis, and peritonitis to very small limits; since, in the vast majority of cases, when we exclude extension of inflammation from neighbouring organisms, and injury, as causes, some one or other of the taints I have mentioned will be found present. That pericarditis, unaccompanied by any marked symptoms of muscular rheumatism, or by any joint-affection whatsoever, is a matter of fairly common occurrence, I have had many instances during the last fifteen years' experience in general practice. Indeed, this is one of the facts in clinical observation that fall peculiarly within the field of the general and family practitioner. Hospital physicians, and men who have attained a high standing in their profession as consultants, usually are called upon to examine and treat the grosser lesions, the results of pre-existing attacks, which possibly had not been observed or treated at the onset, or the effects of which probably had resisted all efforts to cure them.

When reporting a case of heart-disease to the *Medical Press and Circular*, March 31st, 1875, I then expressed my conviction that many cases of heart-affection take their origin in what is termed "a simple cold," where a tendency exists on the part of the patient to the rheumatic or gouty dyscrasia—attacks which I now regard as "rheumatic pyrexia," in which, when the heart becomes affected, it does so as the local affection. I dwell on the necessity for attention to the state of the heart in such cases, and for the utmost vigilance being exercised when there is morbidly excited action, accompanied by roughening and "buffing" of the first sound. I also pointed out how valuable a knowledge of family tendencies to inheritable diseases would be in guiding the physician in his treatment of what otherwise might be passed over as unimportant, and so happily enabling him to prevent such apparently simple cases from terminating in consequences the most disastrous. Since then, increasing experience has impressed upon my mind more and more the deep importance of never lightly passing over the condition of the heart when called upon to treat these so-called attacks of "feverish cold." Truly, in such cases, an over-caution is far preferable to carelessness. Rapid action, weakening even to a slight degree of the first sound, and the occurrence of "humming" or "burring" noises, tending to conceal or give the idea of prolongation to the first sound, are always signs that excite my attention and give rise to anxiety. Of course, just as we may have light and transient affections of the joints, which pass off and leave no trace behind, so, in perhaps the majority of instances, these symptoms may subside and leave no *souvenirs* to mark their occurrence; but, on the other hand I have too frequently watched them develop steadily into the full-blown and well-marked characteristics of advanced pericarditis, to permit of my treating them with contempt or indifference. In the case before us, I think it is a point of interest to note the occurrence of double friction-murmur at the apex, and in the axilla, when the loss of the first sound of the heart, weakening of impulse, and extent of dulness, would all point to effusion; in which case, we expect not to have contact and consequent friction. I can only explain it, first, by chest-wall conduction of the sound from the loud and rasping basal murmurs; secondly from the extent of dulness that still exists, together with the marked improvement in the local and general symptoms; also bearing in mind the full habits and free consumption of malt and spirituous liquors, which may have produced their effects on the kidneys and the circulatory system generally, I am inclined to think that possibly there may be some dilatation and hypertrophy present; in which case, a small amount of fluid only may have been present. This would explain also the absence of displacement of the apex-beat. Another point to be borne in mind is, that the heart itself may have shared in the unusual development of the body generally, and have been unusually large.

The rapid relief afforded by the treatment and the subsidence of the congested appearance of the face and eyes are also to my mind points of interest. No one in this case can well question the benefits derived from the measures adopted; the symptoms had existed long, and had been going steadily from bad to worse, and the relief followed immediately after the use of the remedies.

In conclusion, I would express the satisfaction I feel, at the new, and I think every one must admit, most important step taken by the Association in the collective investigation of disease. A vast amount of, so to speak, outlying and hitherto inaccessible clinical experience will be gathered in, tabulated, and made useful to the profession generally. It will give men fixed points on which to work,

and encourage them to closer and more exact observation; and to the record of such, men who would shrink from the labour of extended note-taking, with a view to publication, will not grudge the lesser labour of answering set questions on given points. I would also say one word to the junior members of the profession in favour of shorthand. Nothing can possibly be of greater use to them. The knowledge of its principles is easily acquired; with a good teacher it is simplicity itself. It only requires half an hour's daily practice, for from two to three months, from dictation, to attain a speed in writing of 80 to 100 words a minute. When written, it is as legible as the present style of writing, or more so, if only care be taken to practise and acquire the power of reading what has been written. It saves a vast amount of labour, and enables those who know it to keep most valuable records of their observations and experience. I owe many a valuable note of cases observed, to its labour-saving properties.

CLINICAL MEMORANDA.

A CASE OF ULCER OF THE STOMACH.

IN connection with the two interesting cases recently published in the JOURNAL, by Mr. Battams, the following notes of one which came under my care at Ranibleet, East Indies, in November 1880, may not be uninteresting. No. 2,517, Pte. J. B., aged 30 according to his records, but evidently an old soldier, in India with the 30th Regiment nine months, of very intemperate habits, came a prisoner to hospital on the morning of November 3rd, and was detained for gonorrhoea. On the 20th, when on the point of being discharged, he complained of feeling "done up, and unable to do anything," also of dyspeptic symptoms, with irritability and uneasiness of the stomach, not relieved by sinapisms, effervescing salines and bismuth powders. Jaundice quickly followed, and he could retain no food. On the 24th he had several attacks of hæmatemesis, and passed a quantity of altered blood by stool. There was marked decrease in the size of the liver, as evidenced by percussion and palpation. Temperature was normal. The symptoms were slightly relieved by treatment. On the 26th, he was not quite so jaundiced. The hæmatemesis continued unchecked by gallic acid and opium, and other remedies. He was somewhat better. On the 27th, there was no vomiting or purging, and he retained a little food, but in the evening he passed a dark tarry stool, and suffered from collapse and great exhaustion: he was relieved by stimulants, but continued very weak from the excessive loss of blood. On the 28th he vomited a pint of altered blood, and complained of much pain in the epigastrium; he could retain no food; and was supported by nutritive enemata. On the 29th, there was no return of the hæmorrhage, but he was in great pain all night, extending up towards the right side, shoulder, and back; he had also symptoms of collapse, and was constantly bending forward to relieve the intense pain. He died quietly at midnight, after having asked a patient in an adjoining bed, what o'clock it was. The *post mortem* examination revealed congestion of the right lung; the heart was small and healthy; there was no atheroma of the valves or of the coats of the vessels. The liver was contracted, small, hard, and of a light nutmeg colour; the gall-bladder distended; the kidneys were enlarged and softened. The spleen was very small, contracted, and weighing only four ounces. The rugæ of the stomach were prominent, and the mucous membrane thickened, especially towards the pylorus, in the immediate vicinity of which was a small ulcer with thickened margins, and base congested where in contact with the liver. The intestines were healthy. No perforation was found. The ulcer was about the size of a four-anna piece.

ALBERT A. GORE, M.D., F.R.C.S.I., Surgeon-Major Army Medical Department; Female Hospital, Dublin.

OBSTETRIC MEMORANDA.

BREECH-PRESENTATION COMPLICATED BY SPINA BIFIDA. I HAVE read with much interest a case of the above in the BRITISH MEDICAL JOURNAL of May 12th, by Mr. C. Penruddock, reminding me, as it does, of a similar labour which has recently occurred in my own practice. I copy the following notes from my case-book.

Mrs. C., aged 19, primipara, sent for me on April 21st at 8 A.M. Labour, I learnt, commenced the previous evening at eight o'clock. On examination, I found the os fully dilated, the pains strong and bearing-down in character, the presenting part of the child just en-

tering the brim of the pelvis, the membranes being entire. First, my finger came upon what I supposed to be the tuber ischii of the child, and farther back a soft roundish swelling which, on firm pressure, felt like several vertebral spines, suggesting to me that, instead of a breech presenting as I supposed, possibly I might have a shoulder to deal with. This induced me to further examine for the fetal clavicle, which I failed to find; so being a little puzzled as to what the presentation might be, I decided to insert my whole hand into the vagina, in order to more thoroughly explore the contents of the uterus. I then discovered that I had a breech complicated with spina bifida to deal with. As the os was fully dilated, I ruptured the membranes, and left the termination of the case to nature.

The child, a male, was born at 9.30 A.M. It was of average size, well nourished, and apparently perfectly healthy, with the exception of this arrest of development in the three lower lumbar vertebrae. The tumour was of the size of a large walnut, spherical in shape, and covered with a thin bluish membrane.

The child remained in fair health for the first fortnight, when ulceration of the sac commenced, and convulsions supervened at the third week, terminating fatally. The mother made an excellent recovery. She attributed the malformation of her offspring to a fright she had had from a dog when six months advanced in her pregnancy, stating soon afterwards to her husband that she felt sure the child was not right.

HAROLD THOMPSON, M.R.C.S., L.S.A., Oxford.

SURGICAL MEMORANDA.

OBSTRUCTION OF THE BOWELS: FÆCAL VOMITING: RECOVERY.

HAVING read Mr. Alder Smith's "Successful Case of Gastrostomy for Intestinal Obstruction," which appeared in the JOURNAL for May 26th, I am induced to communicate the following case, which is, perhaps, worth recording.

On April 11th, at 10 P.M., I was hurriedly sent for to visit a lady, aged about 45, who was said to be suffering from "cramps of the stomach." She was in bed, vomiting frequently, and complained of intense pain of the stomach and bowels. Her pulse was little affected, her tongue clean, her temperature normal, and her bowels had been freely moved twenty-four hours previously, after the use of aperient medicine. I prescribed bismuth with hydrocyanic acid, and also a full dose of tincture of opium, under the impression of having to deal with a case of acute gastralgia. The treatment had no marked effect; for, upon visiting her five hours after, I found she had passed a restless and sleepless night. The pain was sometimes acute, and the nausea and vomiting recurred frequently. I was shown a hand-basin containing upwards of a pint of distinctly faecal material which she had just vomited, and her breath had also a strongly faecal odour. The real nature of the case was now apparent. On careful examination, I could ascertain no cause of strangulation; no external hernia, nothing abnormal within reach by the rectum, and no abdominal tumour existed, and faecal impaction could not be looked upon as probable. Copious injections failed to bring a trace of faecal matter from the bowels, and only served to show that obstruction was complete. The abdomen was distended, and the pain, as already noticed, often most severe. The early appearance of faecal vomit was remarkable. In all the circumstances I ascribed the symptoms to a twist, or to an intussusception at some point in the course of the small intestines. If due to intussusception, might not the purgative taken by the patient have had something to do with its production? We know that invagination is apt to arise from causes that produce increased irritability of the bowel. The stercoraceous vomit enabled me to form an early diagnosis, a point of the greatest moment in these cases, as it enables us to adopt a rational course of treatment. Better leave such cases entirely to nature, than administer a single dose of drastic medicine. No time was lost in placing the patient under the influence of opium. The drug was given as tincture, but generally in the form of powder, frequently repeated and continued throughout the attack; and no food of any kind was taken, for which, indeed, the patient expressed no desire. Ice was not procurable, but cold spring-water and soda-water were enjoyed in small quantities, frequently repeated to allay thirst. The effect of the opiate was soon apparent. Vomiting became less frequent, no doubt from the influence of the drug in controlling intestinal peristalsis; and the patient became comparatively easy, and had some