Where great inflammation or shrinking exists, extraction is out of the question, for sympathetic ophthalmia is thereby favoured, without any compensatory improvement of vision. Enucleation only remains for such cases.

A few items of a statistical nature may be added. Cysticercus in the eyeball is rarer in England than on the Continent, and especially few are the cases from English sources where the deeper parts of the eye were concerned. On the other hand, where the anterior parts were the seat, the number is relatively large. In North Germany, in von Gräfe's time, the proportion was 1 per 1,000; but, in the last few years, a decline has been observed by Hirschberg, who met with only three cases in 8,000 new eye-patients during fourteen months—1882-83. This is probably the result of the compulsory examination of flesh-meats, both in respect of trichina and of cysticercus, which obtains in Germany. The occurrence of the parasite is much rarer in southern Germany, Austria, Italy, etc. In Vienna, for example, *Tænia mediocanellata* is the rule, *T. solium* the exception. As *Cysticercus mediocanellata* infests oxen, and cannot thrive in the human subject, it is never met with in the human eye. The only scolex found in the human eye is *C. cellulosa*, and hitherto only one entozoon has been found in a given eye, though Becker (*Mauthner's Ophthalm.*) says he has seen two in the same eye. The greatest size and age on record in the human eye are 15 millimètres and two years respectively.

The number of cases reported where an anatomical examination of the eye took place is about twenty, and of these, ten were sub-retinal, ten in the vitreous body. Von Gräfe thought that the original seat of the parasite was twice as often beneath the retina as in the vitreous body, the entozoon in a number of cases subsequently breaking through into the latter.

ON THE COMBINED USE OF THOMAS'S AND FURNEAUX JORDAN'S LAMINATED SPLINTS.

By E. STANMORE BISHOP, M.R.C.S. Eng.,

Surgeon to the Ancoats Hospital and Ardwick and Ancoats Dispensary, Manchester.

IN the BRITISH MEDICAL JOURNAL (July 15th, 1882), the use of laminated plaster splints is recommended and very clearly described by Mr. Furneaux Jordan. He advises their use in Pott's disease of the spine, in hip-disease, fractured femur, contracted knee, etc.; and he points out the fact, which must be at once patent to every one, that their application requires only one movement of the patient—namely, that on to the surface of the splint; and does away entirely with the very great annoyance and source of non-success in cases of fracture of the neck of the femur, that of constant relifting of the patient as each successive turn of the bandage has to be carried around the back.

In this communication, I wish to deal entirely with cases of hip-disease and fracture of the neck of the femur. There are good reasons for bracketing these two classes of cases together. First, in both there exists a necessity for some method of dressing which will allow the patient quickly and safely to get up. In hip-disease, the patients are generally already marasmic, and fresh air and a modicum of exercise are of immense importance to them. In fracture of the femoral neck, the accident usually occurs in elderly people, who bear confinement badly. Secondly, in both there exists a necessity for absolute immobility at the hip-joint.

If these reasons be taken together, it will be seen that a method of treatment which will suit the one, cannot be far wrong if applied to the other. This being granted, two methods naturally occur to our minds as being most likely to fulfil the indications presented; and both, at first sight, and on reading the papers of their introducers, appear the thing required. I refer, of course, to Thomas's splint, and to some form of plaster apparatus, of which, without doubt, Jordan's laminated splint affords the best example.

I suppose that no method has ever perfectly fulfilled the hopes and predictions indulged in by their authors; no practice of any plan proposed is so entirely *coulour de rose* as the original theory; and I wish to indicate certain points in which both these methods are likely to disappoint those who trust to either entirely—points in which, when used conjointly, each supplies the other's defect; then, as in some combinations of drugs, powers reside which are sought for in vain when these are separate—I will show that this conjunction renders possible a control over the limb, and, through it, over the joint, which cannot be obtained when either is used singly.

From my own observation, I should say that most surgeons pay but little attention to the actual measurements given by Thomas as to the thickness of his splints; or surely the weight of them would have been objected to long since. Probably, in the majority of cases, the size and weight are left to the instrument-maker, who proceeds on common-sense principles. In other cases, especially in the country, the surgeon will choose his iron in the most convenient blacksmith's shop, and be guided by his own ideas as to the weight his patient can carry, having it afterwards covered and padded by the local saddler; but, if any one who has been doing this, with comfort and ease to his patient and satisfaction to himself, come, like myself, to have to dictate actual thicknesses and widths of iron to a professional instrument-maker, and take them direct from Mr. Thomas's work, he will be amazed to find that he will have a splint which is so heavy that, in the majority of cases, instead of thereby enabling his patient to get up, he has still more certainly condemned him to bed, as the only place in which its weight can be supported.

When we consider that a child with hip-disease is already weakened and emaciated by the pain it is suffering, or has suffered, that its muscles are more or less degenerated by the state of tonic contraction in which they are kept by the irritation of the branches of their own nerves supplied to the joint, as demonstrated by Hilton, it is perfectly evident that we have not much to fear from any sudden or powerful contraction of them; and that the minimum amount of weight, compatible with the requisite rigidity, will be most conducive to the second object of the splint, that of allowing the patient fresh air and exercise.

The amount of weight in a splint made for me, according to Thomas's directions, for a woman aged 26, was 4 lbs. 5 oz., and she could only bear it on when in bed, or, if standing, for a few minutes.

Rigidity, too, may be equally obtained by making the hinder rod still thinner, and supplementing its action by the side-rod recommended by Morris, in the last edition of Holmes's *Surgery*. He, however, merely carries the side-rod to the thigh clip; whilst, for a reason which will be presently evident, I prefer to carry it to the lower or leg clip.

But, apart from the weight, Thomas's splint has another drawback—its readiness to slip round; and the bad effects of this are far more evident when we have cases of fractured neck of the femur to deal with than in hip-disease. It does not fit closely to the body—closely, that is to say, in the sense of having any particular grip upon it. As I read Mr. Thomas, he seems to think that the way in which the upper or body clip is fixed or bent on the opposite side to the disease will suffice to prevent this. I can only say that I have never met an instrument-maker who could give it this knowing set or bend; and that, in one or two cases in which this was supposed to be done, the patient complained so forcibly of the impediment caused to the free action of his ribs, that the cunning bend had to be unbent, as the lesser of two evils, and the irritating tendency to swing round remained as before.

This, I take it, is practically the great drawback in the use of Thomas's splint; the weight can be regulated, but how is the tendency to slip round to be remedied?

Before answering this question, I should wish to examine the working of Jordan's laminated plaster. Will this take the place of the other? Are there no faults to find with this?

The laminated plaster, as I began by saying, has a great advantage over all other methods, except perhaps the Bavarian, of applying plaster-of-Paris, in that the position of the limb can be fixed once for all, and not interfered with again until the plaster has set. It is light, can be well padded, and, when dry, fits closely and firmly to pelvis and limb. Can anything further be required?

Well, practice does not bear out its perfect rigidity. It is extremely difficult to keep that part of it which is most important, the flexure of the hip, firm and uncracked; indeed, I may say I have never seen one that fully bore out its theoretical advantages. This, no doubt, may be due to several things: bad plaster, unskilful application, etc.; but, in ordinary practice, the fact remains that these splints frequently fail to give satisfaction from this sole fact, that they, as ordinarily applied, have not sufficient rigidity.

Still, they have a good grip of the body and the limb; they have no tendency to slip round, are light, and add no appreciable weight to the burden the patient has to carry.

Thomas's splint, then, has for *advantages*: perfect rigidity; no possibility of flexion or extension at the hip-joint; for *disadvantages*: weight, if made according to the original directions; want of actual grip; and tendency to rotation around an axis represented by a line drawn upwards through the main axis of the limb.

Jordan's splint has for *advantages*: lightness; actual grip of the pelvis and limb; and no tendency to rotation; for *disadvantages*: want of rigidity; danger of flexibility over the most important point.

As tabulated, each shows strength exactly where the other is weak. This alone might appear to point to the advisability of their combination; but I would wish to point out that such combination has one or two definite advantages. And first, as to the way in which they may be used together. It is not enough that the one should be merely superimposed upon the other; not so are the best effects of their united action to be obtained.

In making the Thomas's splint use iron of No. 15 gauge, for the uprights; No. 21 for the clips; and, in fixing the uprights, let the hinder one be placed as Thomas recommends, following his line. In addition to this, let a second be fixed so as to lie in the mid-line of the side of the body and limb, hollowing it out in a curve over the trochanter, and riveting it to each of the three clips. It requires no padding; but a simple covering with leather, for the sake of appearance; straps and buckles as originally directed.

As to the laminated plaster, cut out five or six sheets of the stiff muslin used; the inner two require nothing further beyond saturating with plaster; but the outer three or four should have slits cut in them, the width of half-inch tape, and directly opposite the iron uprights. Two slits, *i.e.*, one opposite each upright, should be cut in three places: *a*, midway between the crest of the ilium and the trochanter; *b*, midway between hip and knee; *c*, two inches below knee. The *a* tape should also have another point of exit in front, by means of which it should be attached to a ring on the front strap of the upper clip. Now, lay the tape between the inner two and outer four sheets, allowing a good loop to pass out at each slit; and having saturated the sheets with plaster, and properly moistened them, lay the patient upon the whole.

Fix the plaster splint carefully and allow it to dry, carefully drawing out each loop as far as it is free, and dividing.

Thus, the upper tape will have three points of exit from the plaster splint: one behind, opposite the upright representing Thomas's original upright; one on the outside of the limb affected, opposite the additional upright; and one in front, opposite a ring attached to the strap connecting the upper or chest clip. The free ends of tape projecting from these openings are to be made fast to the points of Thomas's splint indicated; and thus that splint can be rendered perfectly firm and immovable.

The two lower tapes have each two points of exit, both opposite the respective uprights. Now it can easily be seen that, if the position of the limb with reference to the trunk be correct, by fastening these tapes such position is maintained; but, should there be too great eversion, by drawing the outermost tape up and over the anterior surface of the leg, until, by passing round the inner side, it reaches the hinder upright, passing around this and drawing upon it, any amount of traction can be exerted and maintained by fixing the tape in this position; such tape cannot injuriously compress the tissues of the leg, as it is warded off them by the stiff plaster casing. Inversion can be treated in the reverse manner.

Thus, then, the combination of the two splints permits of definite regulation of the amount of eversion or inversion of the limb. The upper part of the outer instrument is fixed in a definite position, from which it is impossible for it to slide; whilst the lower is thus made available as a rigid point or points, to and from which the leg can be drawn and fixed.

Such control over the position of the limb, as to rotation, is impossible with either of these appliances separately; combined, it is at once practicable and easy. To sum up: the advantages to be gained by the combination of the two splints are these.

1. It becomes possible so to lighten the weight of Thomas's splint that it can easily be borne by the weakest patient; and the weight of the laminated plaster is so inconsiderable, and so diffused over a large area, that the combined splints weigh about half the amount of an ordinary Thomas's splint, made after the original directions.

2. Whilst the Thomas's splint supplies the requisite impediment to flexion and extension of the hip, the laminated plaster supplies the firm grip of pelvis and limb, and forms an admirable *point d'appui* for any attempt at adjustment, in cases of fracture.

3. The combination permit of a more certain fixation of Thomas's splint itself, in any desired position; and renders it not merely a safeguard against flexion and extension, but a very certain appliance for remedying excessive rotation, either external or internal.

Of course, the patten on the opposite foot, the crutches, etc., are equally necessary, as they would be with the Thomas's splint alone.

INFLUENCE OF BENZOATES OF ALKALIES ON THE EXCRETION OF URIC ACID.

By EDMUND ALLEYNE COOK, L.R.C.P.Ed., F.C.S., etc.

DR. GARROD, in his third Lumleian lecture, as reported in the BRITISH MEDICAL JOURNAL, April 21st, 1883, gives a summary of some interesting experiments on the influence of hippuric and benzoic acids on the elimination of uric acid, the conclusion he arrives at being that these two former, when in dilute solution, and alkaline, are capable of destroying the latter as a chemical experiment, and, when administered medically, of materially lessening the excretion. He gives some sets of experiments, which show that benzoates, when administered in set doses, have the effect of diminishing the uric acid. I have repeated his experiments so far as regards the administration of benzoates, and do not get results similar to his. On May 13th, the urine was collected, the diet having been regulated exactly for seven or eight days previously. The collections of 14th and succeeding days are as given in the following table.

Date.	Urine.	Specific Gravity.	Uric Acid.
May 14th...	38 ozs.	1020	13 grains.
" 15th...	36 "	1022	12.3 "
" 16th...	26 "	1030	13.0 "
" 17th...	29 "	Not analysed	
" 18th...	20 "	1030	13.2 "
" 19th...	32 "	1023	13.0 "
" 20th...	38 "	Not analysed	
" 21st...	40 "	1023	12.4 "
" 22nd...	48 "	1020	13.7 "
" 23rd...	40 "	1026	14.0 "
" 24th...	32 "	1025	13.2 "
" 25th...	28 "	1025	Not analysed.

On the 20th, 30 grains benzoate of soda in solution were taken at night in one dose. On the 21st, three doses, of 20 grains each, of benzoate of soda, 6 ounces extra water. On the 22nd, two doses of 20 grains. On the 23rd, three doses of 20 grains. On the 24th, none.

The benzoate acted somewhat as a diuretic, although very little extra water was taken. The estimations of uric acid given were made by the zinc sulphate precipitation process; and to corroborate, quantities of the urine were precipitated by the zinc sulphate solution after removal of the phosphates, and the precipitate, washed and decomposed by dilute sulphuric acid, gave abundant crystals of uric acid, as proved by microscope and murexide test.

Results so divergent as these are from those of Dr. Garrod, certainly surprised me; for although I have not seen his paper to the Royal Society, I have followed his directions closely, as given in his lecture-report. His process for estimation, as given in his first lecture, is the acidulation process, and I sought in this some explanation of our difference.

A portion of the urine collection of the 21st, and also of the 22nd, 23rd, and 24th, were set aside after acidulation. These urines gave very little crystallisation of uric acid—a scarcely perceptible trace appearing even after seventy-two hours' standing, with occasional stirring. Of the urine collection of the 21st, 20 ounces were acidulated; and, on the 24th, the crystallisation from the whole 20 ounces was scarcely ponderable, but some feathery crystals were evident, probably benzoic acid.

From this, it is evident that the presence of a benzoate, or even of benzoic acid, can prevent the crystallisation of uric acid from urine; but it is also evident that it does not destroy it; and this result is in agreement with those of Kerner, and of Wöhler and Keller, as quoted by Parkes.

But Dr. Garrod states in his lecture that the uric acid is destroyed, and states that the murexide test failed to answer, after a mixture of a benzoate and a urate in alkaline solution had stood some time at a temperature of the body. He states that a dilute solution is necessary, and from twenty-five to fifty times as much benzoate as urate. In his lecture, he gives nothing more definite than a dilute solution, but I took uric acid, 2 grains, benzoic acid, 60 grains, water, 4 ounces, and dissolved till clear with caustic soda, the solution being alkaline to test-paper; the whole was put into a four-ounce bottle, and kept for nine hours at a temperature of 98° Fahr. One-half of this mixture, diluted and precipitated by zinc sulphate solution, gave nearly one grain of uric acid. The other half, diluted again to 4 ounces, was kept for twelve hours at a temperature of