Jordan's splint has for *advantages*: lightness; actual grip of the pelvis and limb; and no tendency to rotation; for *disadvantages*: want of rigidity; danger of flexibility over the most important point.

As tabulated, each shows strength exactly where the other is weak. This alone might appear to point to the advisability of their combination; but I would wish to point out that such combination has one or two definite advantages. And first, as to the way in which they may be used together. It is not enough that the one should be merely superimposed upon the other; not so are the best effects of their united action to be obtained.

In making the Thomas's splint use iron of No. 15 gauge, for the uprights; No. 21 for the clips; and, in fixing the uprights, let the hinder one be placed as Thomas recommends, following his line. In addition to this, let a second be fixed so as to lie in the mid-line of the side of the body and limb, hollowing it out in a curve over the trochanter, and riveting it to each of the three clips. It requires no padding; but a simple covering with leather, for the sake of appearance; straps and buckles as originally directed.

As to the laminated plaster, cut out five or six sheets of the stiff muslin used; the inner two require nothing further beyond saturating with plaster; but the outer three or four should have slits cut in them, the width of half-inch tape, and directly opposite the iron uprights. Two slits, *i.e.*, one opposite each upright, should be cut in three places: *a*, midway between the crest of the ilium and the trochanter; *b*, midway between hip and knee; *c*, two inches below knee. The *a* tape should also have another point of exit in front, by means of which it should be attached to a ring on the front strap of the upper clip. Now, lay the tape between the inner two and outer four sheets, allowing a good loop to pass out at each slit; and having saturated the sheets with plaster, and properly moistened them, lay the patient upon the whole.

Fix the plaster splint carefully and allow it to dry, carefully drawing out each loop as far as it is free, and dividing.

Thus, the upper tape will have three points of exit from the plaster splint: one behind, opposite the upright representing Thomas's original upright; one on the outside of the limb affected, opposite the additional upright; and one in front, opposite a ring attached to the strap connecting the upper or chest clip. The free ends of tape projecting from these openings are to be made fast to the points of Thomas's splint indicated; and thus that splint can be rendered perfectly firm and immovable.

The two lower tapes have each two points of exit, both opposite the respective uprights. Now it can easily be seen that, if the position of the limb with reference to the trunk be correct, by fastening these tapes such position is maintained; but, should there be too great eversion, by drawing the outermost tape up and over the anterior surface of the leg, until, by passing round the inner side, it reaches the hinder upright, passing around this and drawing upon it, any amount of traction can be exerted and maintained by fixing the tape in this position; such tape cannot injuriously compress the tissues of the leg, as it is warded off them by the stiff plaster casing. Inversion can be treated in the reverse manner.

Thus, then, the combination of the two splints permits of definite regulation of the amount of eversion or inversion of the limb. The upper part of the outer instrument is fixed in a definite position, from which it is impossible for it to slide; whilst the lower is thus made available as a rigid point or points, to and from which the leg can be drawn and fixed.

Such control over the position of the limb, as to rotation, is impossible with either of these appliances separately; combined, it is at once practicable and easy. To sum up: the advantages to be gained by the combination of the two splints are these.

1. It becomes possible so to lighten the weight of Thomas's splint that it can easily be borne by the weakest patient; and the weight of the laminated plaster is so inconsiderable, and so diffused over a large area, that the combined splints weigh about half the amount of an ordinary Thomas's splint, made after the original directions.

2. Whilst the Thomas's splint supplies the requisite impediment to flexion and extension of the hip, the laminated plaster supplies the firm grip of pelvis and limb, and forms an admirable *point d'appui* for any attempt at adjustment, in cases of fracture.

3. The combination permit of a more certain fixation of Thomas's splint itself, in any desired position; and renders it not merely a safeguard against flexion and extension, but a very certain appliance for remedying excessive rotation, either external or internal.

Of course, the patten on the opposite foot, the crutches, etc., are equally necessary, as they would be with the Thomas's splint alone.

INFLUENCE OF BENZOATES OF ALKALIES ON THE EXCRETION OF URIC ACID.

By EDMUND ALLEYNE COOK, L.R.C.P.Ed., F.C.S., etc.

DR. GARROD, in his third Lumleian lecture, as reported in the BRITISH MEDICAL JOURNAL, April 21st, 1883, gives a summary of some interesting experiments on the influence of hippuric and benzoic acids on the elimination of uric acid, the conclusion he arrives at being that these two former, when in dilute solution, and alkaline, are capable of destroying the latter as a chemical experiment, and, when administered medically, of materially lessening the excretion. He gives some sets of experiments, which show that benzoates, when administered in set doses, have the effect of diminishing the uric acid. I have repeated his experiments so far as regards the administration of benzoates, and do not get results similar to his. On May 13th, the urine was collected, the diet having been regulated exactly for seven or eight days previously. The collections of 14th and succeeding days are as given in the following table.

Date.	Urine.	Specific Gravity.	Uric Acid.
May 14th...	38 ozs.	1020	13 grains.
" 15th...	36 "	1022	12.3 "
" 16th...	26 "	1030	13.0 "
" 17th...	29 "	Not analysed	
" 18th...	20 "	1030	13.2 "
" 19th...	32 "	1023	13.0 "
" 20th...	38 "	Not analysed	
" 21st...	40 "	1023	12.4 "
" 22nd...	48 "	1020	13.7 "
" 23rd...	40 "	1026	14.0 "
" 24th...	32 "	1025	13.2 "
" 25th...	28 "	1025	Not analysed.

On the 20th, 30 grains benzoate of soda in solution were taken at night in one dose. On the 21st, three doses, of 20 grains each, of benzoate of soda, 6 ounces extra water. On the 22nd, two doses of 20 grains. On the 23rd, three doses of 20 grains. On the 24th, none.

The benzoate acted somewhat as a diuretic, although very little extra water was taken. The estimations of uric acid given were made by the zinc sulphate precipitation process; and to corroborate, quantities of the urine were precipitated by the zinc sulphate solution after removal of the phosphates, and the precipitate, washed and decomposed by dilute sulphuric acid, gave abundant crystals of uric acid, as proved by microscope and murexide test.

Results so divergent as these are from those of Dr. Garrod, certainly surprised me; for although I have not seen his paper to the Royal Society, I have followed his directions closely, as given in his lecture-report. His process for estimation, as given in his first lecture, is the acidulation process, and I sought in this some explanation of our difference.

A portion of the urine collection of the 21st, and also of the 22nd, 23rd, and 24th, were set aside after acidulation. These urines gave very little crystallisation of uric acid—a scarcely perceptible trace appearing even after seventy-two hours' standing, with occasional stirring. Of the urine collection of the 21st, 20 ounces were acidulated; and, on the 24th, the crystallisation from the whole 20 ounces was scarcely ponderable, but some feathery crystals were evident, probably benzoic acid.

From this, it is evident that the presence of a benzoate, or even of benzoic acid, can prevent the crystallisation of uric acid from urine; but it is also evident that it does not destroy it; and this result is in agreement with those of Kerner, and of Wöhler and Keller, as quoted by Parkes.

But Dr. Garrod states in his lecture that the uric acid is destroyed, and states that the murexide test failed to answer, after a mixture of a benzoate and a urate in alkaline solution had stood some time at a temperature of the body. He states that a dilute solution is necessary, and from twenty-five to fifty times as much benzoate as urate. In his lecture, he gives nothing more definite than a dilute solution, but I took uric acid, 2 grains, benzoic acid, 60 grains, water, 4 ounces, and dissolved till clear with caustic soda, the solution being alkaline to test-paper; the whole was put into a four-ounce bottle, and kept for nine hours at a temperature of 98° Fahr. One-half of this mixture, diluted and precipitated by zinc sulphate solution, gave nearly one grain of uric acid. The other half, diluted again to 4 ounces, was kept for twelve hours at a temperature of

110° Fahr., and then gave very markedly the murexide test in a single evaporated drop. It was left twelve hours more at 100° Fahr., and again tested, no change being apparent. It was then rendered more alkaline with a mixture of potassic and sodic carbonate, and left twelve hours at 100°, at the end of which time, one drop, evaporated as before, gave faint evidence of uric acid by the murexide test.

Again, a weighed quantity of 1 grain of uric acid and 60 grains of benzoic acid were dissolved in 6 ounces of water with soda to a clear solution, and left 12 hours at a temperature of 100°. One drop gave clearly the murexide test. A little mixed sodic and potassic carbonates were added, and the mixture again left 12 hours at 100°. There was now no evidence of uric acid, by the murexide test; but on precipitation with zinc sulphate solution, and solution of the precipitate in dilute sulphuric acid, abundant crystals deposited, of which a minute quantity gave very markedly the murexide test.

Again, a weighed quantity of one grain of uric acid was dissolved in 6 ounces of water by means of sodic and potassic carbonates in slight excess; and one drop of the solution evaporated rapidly (as in the other testings), gave a far fainter test spot of murexide than the drop from the same strength solution of the preceding experiment prior to digestion. This solution was then set aside, at 100° Fahr., for ten hours. One drop then evaporated gave very slight evidence of uric acid by murexide test. After twelve hours, no evidence of uric acid could be obtained by this test.

The results of the above experiments seem to point to the fact that the uric acid is not destroyed by benzoic acid, but exists in a state of combination (it may be as a compound acid), and when this combination is evaporated with nitric acid it gives no murexide with alkalis, and when it is precipitated by zinc sulphate solution, the uric acid combines with the zinc oxide. And the last experiment seems to demonstrate that an alkaline solution containing soda and potassa is capable of really destroying uric acid if kept at a temperature 98° Fahr., even without the presence of benzoic or hippuric acids, probably by oxydation. This was a discovery of Schunk, who found the products of decomposition to be oxaluric acid, and finally urea and oxalic acid; and Dr. Basham has noticed that this takes place in the animal economy when potash salts are administered, coincidentally with the diminution of uric acid (*Practitioner*, vol. v, 1870).

It is well known that there are many substances which, when present in a solution of uric acid, are capable of increasing its power of remaining dissolved, and benzoic acid certainly appears to be one of these.

I have not tried the effect of the administration of hippuric acid by mouth, because it is well known that if benzoic acid be administered, hippuric acid appears in the urine; and if this hippuric acid, formed in the animal, did not diminish the uric acid (though it did prevent its precipitation by acid), neither would hippuric acid administered by the mouth be likely to do so. Moreover, we have no guarantee whatever that hippuric acid, administered by the mouth, will appear unchanged by the urine; and, reasoning from the behaviour of uric acid when so administered or injected, it would certainly not do so.

It is tolerably certain that variations in diet will cause variations in the excretion of uric acid; and, if a strong vegetable variation in the diet of mixed feeders will cause it to be replaced by hippuric acid, it is reasonable to suppose that the same cause, acting in a less degree, will produce the same effect in a less degree; therefore, we may conclude that, in the urine of mixed feeders, we have more or less of a substance which will prevent the precipitation of free uric acid; that the quantity, and therefore the effect, varies with the diet. It must be evident, therefore, that a process of estimation, capable of being affected in this manner, cannot be reliable.

SANITARY INSTITUTE OF GREAT BRITAIN.—At an examination of local surveyors and inspectors of nuisances, held June 7th and 8th, the Institute's certificate of competency to discharge the duties of local surveyors was awarded to P. F. Comber and E. G. Mawbey; and the certificate of competency to discharge the duties of inspectors of nuisances was awarded to W. Dent, S. Edmondson, W. Hearne, G. Johnson, A. W. Merrie, and G. F. Wells.

LUMINOUS PAINT.—An ingenious use of luminous paint now provides, at the Lyceum, a reassuring safeguard against panic. The directions in the corridors and passages as to the "way out" are now as legible in darkness as in light, and thus if from fire or another accident the gas were suddenly cut off, the audience need experience no difficulty in finding their way to the several exits.

ON THE PATHOGENESIS OF PNEUMONIA.

By GEORGE M. GILES, M.B., F.R.C.S.,
Surgeon to the Indian Medical Department, 2nd Punjab Cavalry.

THE publication of a leading article on the above subject in the *JOURNAL* of March 3rd, 1883, leads me to now make this communication, which I have been delaying until I should have cleared up in my mind certain points in connection with an organism I have been observing in the blood and tissues of cases of pneumonia.

In February 1882, I was sent to do duty with the 4th Punjab Infantry at Dera Ismail Khan, on the Punjab frontier, and found that there was then so great an amount of pneumonia among the troops, that it might be well called epidemic. Outbursts of pneumonia of epidemic characteristics are not uncommon among native troops in this part of India, during the cold season, and there is a very generally diffused opinion among the medical officers of the Punjab frontier force, that the disease is, at any rate occasionally, communicated by contagion.

At that time, I had not seen any notice of the observations alluded to in the *JOURNAL*, but so many diseases had been recently demonstrated to be characterised by the presence of micro-organisms in the blood, that I was led to carefully examine the blood of my patients.

On submitting the blood of a case to a power of about 800 diameters (Ross, $\frac{1}{2}$ immersion objective), I was surprised to find that it contained immense numbers of minute more or less spherical foreign particles; and further careful observation of cases brought out the following facts. These particles are present from the very commencement to the end of the attack, and often persist in scanty numbers, far into convalescence.

If the blood be examined in a case of pneumonia as soon as the earliest symptoms declare themselves, there will be observed a number of micrococcus-like bodies about one-eighth to one-tenth of the diameter of a blood-corpuscle, i.e., about one micromillimetre. They appear spherical or nearly so, unless great care be taken in the adjustment of the lens. If examined in a warm, moist chamber, they will be found to possess two motions; first, the ordinary Brunonian vibration, and secondly, an independent locomotion of their own. On occasions when the adjustment of the screw-collar of the objective and the light have been favourable, I have made out, as they twist about, a pyriform outline, and have once or twice had a glimpse of an extremely fine lash springing from the smaller end. After two or three days, a second form appears, in the shape of a nucleated cell from two to three micromillimetres in diameter, the nucleus of which, in size and appearance, is much like one of the first described form. In these, no signs of locomotive power have as yet been observed. The smaller kind are still found in great numbers, and by this time will be often found joined into chains of from two to ten or more individuals. When this is the case, they too appear to have lost the power of locomotion.

Coincidentally with the appearance of this nucleated form (or stage, as I take it, of the micrococcus), a third appearance manifests itself, in the shape of large colonies of the smaller sort, united together by a hyaline structureless material, forming masses as large as fifteen to twenty micromillimetres. These collections are probably zoogloea, and are of a consistence of some firmness; for, in specimens examined warm, I have not unfrequently observed the red corpuscles to run against them, the corpuscle becoming altered in form, while the zoogloea-mass yielded scarcely at all. If the blood be dried on a cover and stained by "Koch's method," these micrococci will be found to stain well with most anilin dyes. That best suited is one I obtained some years ago under the name of "anilin blue." I am unacquainted with its exact name, but it is a different article from methyl blue.

During the progress of the case, these zoogloea-masses commonly disappear, and a more or less distinct repetition of the same cycle may often be observed; but as a general rule the process appears to become confused.

The sputa, too, may, by Koch's method, be demonstrated to absolutely swarm with micrococci, and often, in fact, to be largely made up of them.

After death, both in man and, as will be described below, in rabbits, I have made out their presence in the tissues of all parts of the body, but especially in the capillaries of the lung, which are often found blocked with clot containing immense numbers.

In severe and fatal cases, I have frequently observed a curious circumstance. When the patient is at his worst, the drop of blood