SEA-SICKNESS.

By R. VACY ASH, M.B.Aber., L.R.C.P.Lond.

I WAS glad to see a short communication in the JOURNAL of July 7th on sea-sickness, and hope all who have the chance will contribute something to our at present small stock of knowledge on this distressing complaint. To me it is particularly interesting, for it is my lot to be painfully sensitive to the movements of a ship, and to have passed more than a year of my service on the sea. I have an idea that the sympathetic nervous system is the culprit, for the following reasons.

1. Flushing of the face is a common sign of the approach of nausea, and we all know that irritation of that nerve will cause this, as well as an extra secretion in a gland.

2. There is an increase in the quantity of fluid ejected from the stomach after it has lain there for a short time. In my own case I frequently noticed, and I subsequently verified it in many others, that, if I took half a cup of beef-tea, and lay in the horizontal position for a time, so as to avoid vomiting, when I did again vomit, when the exhausted muscles had regained their tone and were ready for another attack, the quantity ejected was greatly in excess of that taken in. For instance, if four ounces had been drunk, about twenty ounces would be ejected, of a sour beef-tea liquid. Now, whence did the surplus come? That it was gastric juice, may, I think, be taken for granted; for, although I had not the means of chemically examining its component parts, it certainly partook outwardly of the character of that juice, inasmuch as it would dissolve meat and had an acid reaction, and it did not contain any special features that would lead to the supposition that it came from other gastric organs.

Granting then that it was gastric juice, it follows that secretion, induced by the presence of the beef-tea, was in action, while the balancing power of absorption was held in abeyance. Now, if we follow this out we shall see that the sympathetic nerve-power was acting regularly; for secretion of gastric juice is governed in the follicles by the latter, while absorption of fluids direct by the veins which are governed by the former is held in abeyance, or, in other words, paralysed. I do not say that it is so, I only throw these facts out for others to corroborate, or not, as the case may be. Whence could the increase in the quantity of fluid have come? It must have been taken in some way from the blood; and what so ready to do so as the gastric follicles, stimulated into action by the presence of the small quantity of beef-tea?

Again, I noticed a rise in temperature of from one to two degrees in almost all cases on which I made observations. This may have been induced by the excited action of the sympathetic, allowing an increased flow of blood through the capillaries. By the term "excited," I mean "confused" action of that system, corresponding in effect to paralysis in the cerebral system.

Now, as to remedies. If my observations be correct, any drug or remedy acting on the sympathetic nervous system would cure this tiresome complaint; ice to the spine may so act, as well as the remedies mentioned by Mr. Kendall, in a more direct way. The tea-spoonful of Worcester sauce, which, by the by, I have also found useful, may owe its efficacy to the hot condiments contained therein, and I imagine it to be possible that they act through the sympathetic in the coats of the stomach. I need not follow out the other remedies, or name the thousand and one new ones that are constantly being poured down the throats of the unhappy traveller; but I do know that the majority of the quack remedies for sea-sickness contain a mixture of nearly all the carminatives and condiments under the sun, with the hope that one out of the lot will hit, and they do hit, or rather temporarily relieve; as cayenne pepper and Worcester sauce will do. There is one mode of applying remedies that I should like to see tried by someone who would honestly take the matter in hand; and that is, the introduction of certain remedies by subcutaneous injection, for it necessarily follows that, if my idea be correct and absorption be held in abeyance in the stomach, it is of little use to pour any medicine into that viscus when it is impossible to be taken up by the blood. Why the sympathetic nerve is the culprit, I will not attempt to prove. It may be that the motion of the vessel is foreign to the understanding of any cerebral nerves or organs governed by them, and hence is thrown back, so to speak, on the more animal nerves, lower in the organisation for appreciation, and that this, the sympathetic, unable to cope with such a strange motion or phenomenon, gives up in despair and refuses to act naturally. This may be far-fetched, but, at least, it is open to argument in the absence of a better reason.

VESICO-VAGINAL FISTULÆ.

By PERCY BOULTON, M.D., M.R.C.P.Lond.

Physician to the Samaritan Free Hospital.

It is unnecessary for me to give a vivid description of the terrible state to which a woman is reduced who suffers from a severe form of this injury. Probably, nothing except cancer renders her so unfit for every duty or pleasure in life. When such a fistula is favourably situated, that is to say, some little distance from the neck of the bladder, and where there is ample tissue to bring the parts easily together, any neat operator, with ordinary manipulative skill, may succeed. In cases the reverse of this, he will require a considerable fund of resources and patience. Such cases often become plastic operations, surrounded by every possible difficulty, requiring the greatest dexterity and mechanical inventiveness, not only as regards the tissues to be dealt with, but also with the instruments to be used for the purpose of repair.

When the operation seems to be successfully completed, we have to deal with a viscus that is never at rest, and that is the receptacle of a fluid peculiarly penetrating and irritating to raw tissues; and, although our operation may be well timed, so irregular and uncertain are some women, that the catamenia may come on unexpectedly at the very time when it is ruination to all the work.

The two following cases have just left my ward, at the Samaritan Hospital, cured, and a short account of each will be instructive and, I think of interest.

CASE I.—M. E., aged 65; married, had suffered for twelve years from prolapsus uteri, for which she had worn supports. The last one, inserted by her medical attendant in the country, was a "Blackbee," after wearing which for about three months she found that she was constantly damp. After noticing this for four months more, she was seen by my colleague, Dr. Day, who found a fistulous opening into the bladder about an inch from the cervix uteri, through which a five-shilling piece could have been passed. The vesical mucous membrane was everted, and very vascular, like a large caruncle. Dr. Day transferred the case to me for operation.

On January 26th, I carefully pared the edges, using for this purpose a small tenaculum and a pair of curved scissors, the patient being on her back, the parts exposed, with the Sims' speculum in the vagina, and a straight metallic sound in the bladder. I brought the edges together with eight fine wire sutures; and the patient, who refused to be anaesthetised, never uttered a sound. The urine was drawn every six hours, and, on the tenth day, I removed all the sutures, and found the opening completely closed and water-tight. I kept her in hospital another fortnight under observation, and she returned home on February 18th, well.

The age of the patient and the very vascular condition of the vesical mucous membrane suggested some degenerative change as a cause of want of vitality in the parts, leading to such extensive injury, from what must have been slight pressure in so lax a vagina. Otherwise, this was a favourable case for operation, as the result proved.

CASE II.—M. B., aged 37, deserted by her husband, came up from Cornwall to see me at the Samaritan Hospital in November 1881. Her last confinement had been seven years before, and she had never been able to retain any urine since. The labour was a very severe one; for five hours, the patient had been in convulsions, and was unconscious at the time of delivery. She was told that the forceps was applied, and that a strong pain suddenly expelled forceps and child.

On examination, I found the whole anterior vaginal wall torn through from the cervix uteri to within half an inch of the urethral orifice. I could pass four fingers upwards into the bladder, which, from long disuse, was small, and the edges of the cicatrix were hard and unhealthy.

I determined to try to bring the whole together at one sitting, as the number of sutures required would enable me to keep up a slight and even tension that would bring the edges sufficiently in apposition. On November 22nd, 1881, I did this; and, on December 2nd, I removed the sutures, and found all united, except the three anterior ones. This is the part which always gives trouble. Dr. Marion Sims has said he "regards cases in which the neck of the bladder is destroyed as all but incurable."

On December 4th, the catamenia returned after fourteen months' cessation, so that I did nothing more till December 14th, on which day I pared the edges of the opening remaining, and brought it together with three sutures. On December 24th, I found that two out of three had healed, but there was still an opening of the size of a