CASE IV. A private patient, aged 27, had hysterotomy performed by Dr. Bantock. The heart and lungs were normal. She was rather nervous; pulse, 120. Full anaesthesia was produced after inhaling one drachm of the mixture in five minutes, without excitement or retching; pupils contracted, strabismus divergens; pulse, 108. After eighteen minutes, pulse, 78; pupils normal. After twenty-three minutes, pulse, 72. After fifty-six minutes, 92; when the narcosis was stopped, the patient having inhaled four and a half drachms of the mixture. No sickness; no nausea.

CASE V. In a case of exploratory laparotomy by Dr. Bantock, in the Samaritan Hospital, the patient, aged 47, whose heart and lungs were healthy, was in deep narcosis in six minutes, after inhaling one drachm of the mixture, and had altogether three drachms in twenty-two minutes. There was no muscular excitement nor retching. The pulse at the commencement was 120; sunk after fifteen minutes to 94; and after twenty-two minutes, when the narcosis was stopped, to 80.

CASE VI. Dr. Marion Sims removed several small fibrous growths from the uterine cavity of a patient, aged 48, highly anæmic from repeated hæmorrhages, in consequence of which also great nervousity and partial aphasia. Thoracic viscera normal; pulse, 108. She was much afraid of any narcotic, but when on the operating table, quietly inhaled the mixture given in my apparatus very slowly; so that half a drachm brought her in deep quiet sleep in seven minutes. Pulse, 98; after fifteen minutes, 84; after twenty minutes, 87, when the narcosis was stopped, only three drachms having been given. The patient was quietly sleeping during the operation; about half an hour after, a momentary slight retching of froth.

CASE VII. Another patient of Dr. Marion Sims, aged 40, suffering from a very high degree of vaginismus. Pulse, 100. Full narcosis in ten minutes, after inhaling two drachms of the mixture. Had seven drachms in thirty minutes, when the anaesthesia was stopped; pulse, 88. The patient was quietly breathing, and kept perfectly quiet, but the slightest touch of the genitals brought on a start like an electric shock. During the whole time of the operation (thirty minutes), the narcotic mixture was given without interruption, whilst in the other cases it was occasionally discontinued for five minutes and longer. When awaking, the patient said that she did not feel anything of the operation, but complained of local pain. She had no retching during the operation, but whilst removing her into her bed, she brought up about a mouthful of water. All these patients recovered full consciousness almost immediately after the operation, and looked as well as after a natural sleep. There was no struggling or muscular excitement, except very slightly in Case III, and the abnormal reflex movements in Case VI, which were due to the peculiar complaint of the patient.

Dr. Bantock had this mixture given since in two other cases in the Samaritan Hospital, in one of abdominal section, and in another of extirpation of a cyst from the leg, in neither of which sickness or nausea were caused by the narcotic.

STATISTICS OF ANÆSTHETICS IN RELATION TO AFTER-SICKNESS AND DEATH-RATE.

By R. MARCUS GUNN, M.B., F.R.C.S.,

Ophthalmic Surgeon to the Hospital for Sick Children, Great Ormond Street;
Clinical Assistant (late House-Surgeon) to the Royal London
Ophthalmic Hospital, Moorfields.

The following statistics are based on my experience of anæsthetics at Moorfields Ophthalmic Hospital from November 1st, 1876, till January 17th, 1880. Notes were kept of the cases as to sex, age, the anæsthetic employed, and the hour when food was last taken, in reference to the occurrence of sickness. The condition of the weather as to barometric pressure, temperature, etc., was also noted, the weekly chart of the *Times* being used for this purpose.

Whenever any retching occurred during the operation or within the next two hours, the patient was described as having been sick. Sickness while on the operating-table was comparatively rare; but the proportion of cases in which it occurred afterwards seems large.¹ This latter fact is probably accounted for, partly by the profound anaesthesia necessary for eye-operations, and partly by the large number of young out-patients, who, notwithstanding all directions to the contrary, had eaten shortly before. What the usual percentage of sickness at other hospitals is would be interesting to know.

¹ In the Report on the Action of Anæsthetics by McKendrick, Coats, and Newman (BRITISH MEDICAL JOURNAL, 1880, II, p. 351), a still higher percentage is given—viz., 48 per cent.—in the case of ethidene dichloride, and chloroform.

Anæsthetics were given altogether 4,138 times; there was sickness in 1,902 cases, or in 45 per cent.

Sex.—From the following table, it appears that females have a decidedly greater tendency to this form of sickness than males.

Sex.	Total.	Sick.	Percentage of Sick.
Males	2,130	870	40
Females	1,982	1,027	51
Not stated	26	5	—

Age.—	Total.	Sick.	Percentage of Sick.
Under 1 year	119	5	4
Above 1 year and under 5 years	448	137	31
„ 5 years and under 10 years	558	349	63
„ 10 „ „	15	331	72
„ 15 „ „	20	445	60
„ 20 „ „	25	285	56
„ 25 „ „	30	182	52
„ 30 „ „	40	289	39
„ 40 „ „	50	319	31
„ 50 „ „	60	363	32
„ 60 „ „	70	433	31
„ 70 „ „	80	183	27
„ 80 „ „	90	17	29
Age not stated	40	23	—

The liability to sickness is thus at a maximum about the commencement of puberty, and decreases gradually towards each end of life. Under one year of age, there is very little sickness. There is a second slight increase in the percentage of sick in extreme old age; but the total number of cases observed at this period of life (above eighty years) was too small to draw any positive conclusion. The youngest patient I anæsthetised was aged fourteen days, and the oldest was eighty-six years. I consider *age* the most important factor in relation to after-sickness.

Anæsthetic.—Chloroform was frequently used for children under three, and for adults over sixty. Between these ages, ether, a mixture of ether and chloroform vapour, or the ACE mixture was employed. Until the introduction of Clover's inhaler, ether was always given in a hollow conical sponge, about four ounces being necessary for each case. The objections to this method were its extravagance, the choking sensation and struggling of the patient, and the disagreeable smell of ether that impregnated the air and the clothes of every one around. In March 1877, Clover's small ether-inhaler was first employed. The original form was found inconvenient, on account of the frequent necessity of removing the whole apparatus during the operation, whenever the bag became too highly charged with carbonic acid gas. To obviate this difficulty, I had a valve made on the brass tube leading to the bag, so that, by pressure on this, the bag could be emptied and fresh atmospheric air supplied without disturbing the face-piece. By placing over this valve a layer of lint, on which about half a drachm of chloroform had been poured, I found that the choking sensation and struggling during the early stages of ether inhalation were in great measure prevented, and the time necessary for anaesthesia considerably shortened. One great advantage of this form of apparatus is the ease with which the breathing can be watched, from its effect on the bag. The amount of ether can be graduated, too, and more or less chloroform or ether given, as symptoms may indicate. About half a drachm of chloroform and an ounce of ether were sufficient for anaesthesia, and the average time was about four minutes. This combination of the vapours corresponds to the heading "Ether and Chloroform" used below.

By "ACE" is meant the usual mixture of absolute alcohol, chloroform, and ether, in the respective proportions of one, two, and three parts.

	Total.	Sick.	Percentage.
Ether (by sponge)	263	128	48
Ether (Clover's inhaler)	1,072	574	53
Ether and Chloroform	2,306	1,089	47
Chloroform	320	61	19
ACE mixture	58	17	29
Not stated	83	33	—

One is at first struck with the small percentage of sickness after chloroform. But it must be borne in mind, as mentioned above, that it was chiefly given at the extremes of life, at periods which are naturally less liable to sickness after anæsthetics. It may be argued that these tables show a small percentage of sickness in the very young and old, because chloroform was used in such cases. As a fact, however, the ether and chloroform mixture was employed in the majority of them, and the result as to sickness was the same.

One of the arguments used in favour of ether, especially in cases of ovariectomy, is less liability to sickness than with chloroform. My impression is that, ages being equal, the tendency to after-sickness is about the same with each, but the nature of the sickness is different in the two cases. Ether-sickness seldom lasts long after

the stomach is emptied, though there are complaints for some time of a "nasty taste in the throat." Chloroform-sickness often continues for several hours, and leads to great exhaustion.

Influence of Food.—From a careful comparison of the cases I draw two conclusions.

1. No breakfast² (or too long a fast) is a disadvantage, especially in the very young and the aged.

2. Speaking generally, seven to eight o'clock seems to have been the most favourable time for breakfast for patients of all ages, *i.e.*, the last meal should be given about four hours before operation.

Influence of Weather.—It must have been remarked by every one in the habit of giving anesthetics, that on certain days nearly every case is sick, while on other days sickness is quite exceptional or even entirely absent. These meteorological notes were kept (mainly at the suggestion of my predecessor in office, Mr. Morton) to see how far the weather had an effect in this way.

During the entire period of three years, there were fifty days on which the tendency to sickness was exceptionally great. The temperature of the theatre was kept fairly uniform, except in very warm summer weather, and neither this nor the barometric pressure seems to have had any influence on the liability to sickness. The only peculiar coincidence, which I simply mention without laying any stress upon it, is in reference to the direction of the wind. Of the forty-two sick days, on one only was the wind easterly, while it was from that exact quarter on eleven of the fifty exceptionally exempt days. The moisture of the air, as measured by the wet and dry bulb thermometers, varied much in both series.

Several of these exceptional days could be accounted for by the ages of the patients that were anesthetised. Thus, if nearly all were in early adult life, or about the age of puberty, the rate of sickness was high; if, on the contrary, there were many very young children, or aged adults, the rate was low. Still, however, there remained many days on which the peculiar liability to, or immunity from sickness, could not thus or in any other apparent way be explained.

On comparing the corresponding months of the three years, the greatest percentage of sickness for the entire month occurred twice in January; the lowest percentage always in March.

Mortality.—One death occurred, the details of which were published at the time in this JOURNAL (1877, vol. ii, pp. 266 and 318). The patient was a very stout woman, aged 46. Before going into the operating-theatre, she told the nurse that she would die on the table. She inhaled a mixture of ether and chloroform vapour for about one minute, when the pulse became very feeble, and the respirations shallow. The anæsthetic was then at once removed, and stimulants tried (flicking with wet towel, etc.). The pulse and respiration persisted for fully four minutes after the discontinuance of the anæsthetic, but then ceased, notwithstanding artificial respiration and other measures. At the necropsy, the muscular walls of the heart were found to be fattily degenerated, and there was much fatty growth around the organ. The mitral orifice was contracted, and the aortic valves incompetent. Considering all the circumstances of the case, I am inclined to think that the death occurred more from shock than from any poisonous dose of the anæsthetic.

How far the following case is to be considered one of death due to anæsthesia, it is now impossible to say. The patient was a thin, delicate-looking girl, aged 8, with light hair and fair complexion. She took the anæsthetic (ether and chloroform) well, seemed to recover in the usual way, and spoke once to the nurse in reply to a question. She continued to look pale, however, and became sick for the first time more than an hour after the operation. Being an out-patient, she was shortly afterwards taken home by her parents, travelling in a tramway-car. Next morning, the father called, and told me that she never spoke after leaving the hospital, and died in the evening. There was no necropsy, and I was not summoned to attend the inquest.

Ether was introduced as the principal anæsthetic at Moorfields about the beginning of 1874. Calculating that anæsthetics have been given from that date until now at the same rate as during my term of office³—a perfectly just calculation—we have a total of about 13,000 administrations. The only fatal result that can fairly be attributed to the anæsthetic is that of the woman mentioned above.

Next to carelessness in administration, nothing is more calculated

to increase the death-rate of anæsthesia than wide-spread dread of its danger. An article by Mr. Roger Williams, recently published in this JOURNAL, is, I consider, calculated to do harm in this direction. His position as surgical registrar to an important hospital gives an official weight to his remarks, and lay readers might therefore be unduly influenced by them. His experience—a mortality of one in 350 inhalations—is certainly exceptional, and I cannot believe that "the majority of fatal cases are hushed up." Even this frightful mortality does not, according to Mr. Williams, "represent the whole of the danger; for, in a still larger proportion of cases, alarming symptoms supervened, and fatal results were only narrowly avoided." Now, alarming symptoms will, we all know, unavoidably occur occasionally during anæsthesia; but the greater the proportion of such cases to the actual number of deaths, the smaller the *real* danger of the anæsthetic. In other words, the danger is estimated by an appeal to the death-rate; and the more frequently we can avoid a fatal termination after the occurrence of alarming symptoms, the more we are encouraged. The hypothetically perfect anæsthetic should neither cause alarm nor death; but I fear it will never be found.

Again, the anæsthetic must not be blamed for all cases of death during its administration. Chloroform vapour, without a due admixture of air, will paralyse a strong heart, not to mention a fatty one. (See Case I, described by Mr. Williams.) The shock of a severe operation may kill without any anæsthetic, and none the less when perfect anæsthesia has not been produced. (See Cases IV and VI.) In both these cases, too, blood was found at the necropsy even in the smaller bronchi; and, in Case VI, the "lungs were firmly bound to the chest-walls by old adhesions."

A moribund patient may die from the mere shock of removal to the theatre, and the dread of the impending operation; nor is this risk removed till perfect anæsthesia has been produced. If chloroform be given to such an one while the head is raised, and if, under such circumstances, death occur, we should not lay much blame on the anæsthetic. (See Case II.)

The vomiting of acute intestinal obstruction and peritonitis has been justly attributed to peripheral irritation of the vagus; the great and early collapse in such cases may fairly be held to be due to a reflex inhibition of the heart's action, through the influence of the same peripheral irritation on this nerve. The dilatation of the cardiac orifice of the stomach, which occurs as a preliminary to the act of vomiting, is, like the inhibition of the heart's action, here caused in a reflex manner, by impulses descending the vagi. Great dilatation of the orifice, manifested by a sudden large gush of vomited matter, is then a sign of a powerful descending impulse; a sudden stoppage of the heart's action in diastole is also produced by a powerful descending impulse. It cannot, then, be a matter for surprise that death should so often occur, in this manner, in cases of acute intestinal obstruction; and we may surely expect to meet with this association of vomiting and sudden heart-failure, occasionally, during, as well as before and after, the induction of anæsthesia. Three of Mr. Williams's cases (III and V, and an additional one brought forward by him in the JOURNAL of May 5) are of this nature. In all of them, vomiting occurred at or about the last moment of life; in two of them, the heart was found, at the necropsy, in the condition of diastole. In the remaining one, I think death might have been at least as fairly attributed to the presence of foreign material in the air-passages as to the anæsthetic. "The air-passages, even to the smallest bronchi, contained fluid fecal matter, which was probably forced into them during the process of artificial respiration."

On the whole, I think that the cases (seven in number) brought forward by Mr. Williams are unfortunate, and most unsuitable for the purpose of founding a death-rate upon them. A rate of one in 350 inhalations, due really to the anæsthetic, is, I hold, much too high. A mortality of one (or at most two) in 13,000 cases, as at Moorfields, corresponds more nearly, I think, with general experience. I can truly say that not one of our cases was hushed up.

I suppose I need hardly say that, on the subject of anæsthetics, I do not speak as a specialist, but only with the object of making my past experience of some service. My aim has been a double one—viz.: first, to show how far my statistics explain the liability to after-sickness; secondly, and more especially, to prevent an unwarranted popular dread of anæsthetics—a dread, I am convinced, in itself most dangerous.

MEDICAL MAGISTRATE.—Edward Lister, L.R.C.P.Ed., M.R.C.S. Eng., L.S.A. (retired), of Swarthdale, Ulverston, has been placed on the Commission of the Peace for the County of Lancaster.

² The operations at Moorfields are performed from about 10.30 to 1.30. No breakfast was frequently recommended.

³ I gave no anæsthetics during about two months of this period, on account of the hospital being closed for repairs. This must be considered in making the calculation.