but where is the individual life in disease? In my humble opinion, the germ will not serve our purpose, for, according to the great Darwinian analogy, the germ is always a part of the individual, and always presupposes the individual. The germ, or the sperm, is no doubt a peculiarly important part, and it is charged with the most marvellous representative powers; but it is always representative of the individual, and it derives its powers from the individual. Such, at least, is our only analogy.

If, then, we must have the individual to start with, before we can apply the "origin of species" to disease, we come back to the old question, how a morbid state of the body can become a semi-independent thing, how it can exist, not in absolute independence of the body, but autonomous within it, an *imperium in imperio*. This is the perplexing question which I stated at the outset with reference to cancer, a disease which, in the popular estimation is almost synonymous with a semi-independent life, and I quoted the words of Paracelsus: "In such a disease, a man is himself and another; he has two bodies at one time, enclosed the one in the other, and yet he is one man." I have spoken to several metaphysical friends of the difficulty of conceiving how a mere state of the body, a complex or integrated morbid state, can become an individual existence with the power of propagating itself; but I cannot say that I have got anything satisfactory out of them. We shall probably have to settle this question within our own science as a part of the subject-matter of pathology. Pathology is indeed varied enough in its subject-matter. It studies disease on many sides, as the founder of this Association planned that it should be studied—the side of its anatomy and its physiology, of its geography and its ethnology, of its history and its natural history. Its aspects are as various and attractive as the fruit trees in the Mohammedan paradise. But there is nothing in all this boundless field of inquiry that is more likely to occupy the mind of the profession for years to come, as it has occupied it in times past, than the difficulty for conceiving how a state or condition of the body can become a species of disease.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF OBSTETRIC MEDICINE,

At the Annual Meeting of the British Medical Association in Liverpool, August, 1883.

By GRAILY HEWITT, M.D., F.R.C.P.,

Professor of Obstetric Medicine, University College; President of the Section.

THE QUESTION OF FOOD IN OBSTETRIC AND GYNÆCOLOGICAL PRACTICE.

GENTLEMEN,—Permit me, in the first place, to express the deep sense which I entertain of the honour conferred upon me by the Council of the Association in requesting me to preside over the meetings of this section. I trust that I shall be favoured with your kind indulgence in endeavouring, to the best of my ability, to fulfil the duties entrusted to me.

It is with much pleasure that I congratulate the section, and our esteemed colleague, Sir Spencer Wells, on the public recognition of his important services in the cause of humanity which is embodied in the honour which he has recently received. The extended application of surgical skill to the relief of a class of diseases which were formerly considered incurable, and in which Sir Spencer Wells has taken so considerable a part, constitutes a great claim by gynæcological practitioners on the gratitude of the public, and I am quite sure that you will join with me in felicitating our distinguished colleague on his well-merited reward. None the less is the profession to be congratulated on the further recent developments in the application of surgical skill to the relief of suffering which have been made by gynæcologists, as evidenced by the fact that novel and bold operations, which were hardly thought of as possible before ovariectomy had become an established success, have of late been performed with a large amount of success also.

While, however, we congratulate ourselves on the brilliant achievements of gynæcological surgery, it is well to be reminded that there are other kinds of work to be done, on the due carrying out of which the lives and health of vast numbers of the community depend, and in which, as gynæcologists, we are deeply interested.

I take advantage of this opportunity to present to you a few ob-

servations on a matter which has for some years deeply interested me, and concerning which I have arrived at conclusions which, to me at all events, appear to possess a certain value, viz., The Question of Food in Obstetric and Gynæcological Practice.

Now, it may appear unnecessary at this period of medical history, and in the present advanced state, as it is considered to be, of medical knowledge, to insist on the importance of food in maintaining a healthy activity of the vital processes. But a lengthened experience has convinced me that the public at large have no practical knowledge of this, and that the most disastrous effects result in multitudes of cases from this ignorance of simple physiological laws.

Engaged as I have been more particularly in obstetric and gynæcological practice, the non-observance of these important physiological laws in connection with the occurrence of diseases of the uterus, disturbances of the functions of the uterus and ovaries, and the diseases incidental to child-bed, has forced itself on my notice. It is pretty certain that the generalisation which applies to these classes of disease extends, or would be found to extend, to diseases of other organs of the body, for it is the merit of a true generalisation that it is of wide application.

There are certain terms in general use by the public and by the profession which are often employed in a vague, unmeaning, and indefinite manner—such as "weakness," "delicacy," etc.—whereby it is intended to designate a condition very frequently met with, but the essential nature of which is, as my experience has shown me, very imperfectly apprehended. It is well recognised that this weakness, delicacy, or what not, is very commonly observed not only in young women coming under the notice of the gynæcologist, but also in women who have arrived at adult age, and suffering from disorders in connection with pregnancy, parturition, and child-bed. My observation has in a particular degree been attracted to the presence of this condition. I have taken every opportunity in my power for analysing its nature and ascertaining its essential cause. One conclusion arrived at is, that this general weakness is very far more common than is supposed as an antecedent in cases coming under our notice as gynæcologists, and it may even be said that it is almost invariably present in a marked form in such cases. Another conclusion is, that the "weakness" is associated with and caused by a long-continued inadequate dietary, and that it is essentially what may be termed "chronic starvation." The rational view of the matter is, that this "weakness" is, in the very large majority of cases, preventable; that it should be regarded as the first stage of a serious possible disease; that it is mostly the result of chronic starvation; that it is curable, to a greater or less degree, in most cases; and that it is of the highest importance that it should be formally and carefully treated, as other diseases are, instead of being allowed to smoulder on until the constitution is thoroughly undermined, or until some ailment distinctly classified in our nosology makes its appearance.

It may be asserted that, as yet, neither the public nor the professional mind has seized the importance of preventive measures in regard to disease. It is, indeed, true that the sanitary engineer is at work in our houses; but it cannot be said, as yet, that anything like systematic attention is devoted to the maintenance of healthy and vigorous growth, and the prevention of those diseases which are ultimately so deadly in their effects in many cases; I mean what may be termed the nutritional diseases incident to the human frame.

The analogy between the growth of a plant and of the human organism is one which has been often pointed out. The various chemical changes involved in a continuance of healthy growth in the plant and in the animal, are well capable of definition. We know that the plant requires air, moisture, and a supply of nutritive material regularly, and in sufficient quantity. And we should know, for physiologists inform us of the fact, that the vital processes in the human organism require, for their continuance in a vigorous state, material supplies of a suitable kind, as well as hygienic surroundings, fresh air, and the like. Again, it is well known by good gardeners that nothing so effectually protects the plant from external injurious influences, from blight, from the attacks of parasites, from insects, and from the diseases to which plants are liable, as the maintenance of the plant in a state of active growth. Once let the growth fall off in its vigour, the plant becomes forthwith liable to fall a victim to canker, to the pestiferous invasion of insects, to deterioration, to weakness, and often to decline and death.

It is a remarkable circumstance that so little as yet seems to have been done in the investigation of the effects of an insufficient dietary, looking at it from the quantitative point of view, although the effects of complete deprivation of food are well known. Thus, in the last edition of Parkes' well known work on hygiene, it is stated

that little is known on the subject of the effects of insufficiency of food; and the space devoted to the consideration of this matter is less than one page.

The valuable facts recorded by a series of careful inquiries into the changes occurring in the human body in a state of health do not seem as yet to have been applied at all extensively to the treatment or prevention of disease. It should be known that each day the human body undergoes, in a state of health, a certain degree of waste and loss; and that, experimentally, it has been found that health is maintained by daily administration of certain quantities of material in the shape of food. The obvious inference is that, when the supply is continuously below the amount of daily waste, deterioration must occur, and a state of weakness will result.

It is a curious circumstance that the physiological law and the popular impression (shall I say, in too many instances, the "medical" impression) are found widely different. Physiology teaches the necessity for a continuous supply of a certain quantity of food. The popular impression is, that some people do not require so much food as others; and, consequently, important quantitative diminution in the supply often escapes attention. I must confess that, not very many years ago, I shared in the popular impression. What induced me to form the opposite conclusion was that, in the first place, I was struck with the fact that, in almost every case coming under my notice, a state of what was termed "weakness" by the patient had been notably present; and, secondly, that, on inquiry, this weakness was almost always found to be associated with a notably deficient dietary. For the last six or seven years, I have tested the accuracy and applicability of these generalisations, by carefully inquiring into the past history of patients—mostly suffering from some uterine or ovarian disease, or some affection incidental to child-bed; and these conclusions have stood the test of this long-extended inquiry. I have to state the important conclusion, that a continuous insufficiency of food, or what may be termed a "chronic starvation," more or less intense in different cases, was found to have existed almost universally. Consequently, I have naturally been led to consider chronic starvation as a most important factor in disease, certainly in those classes of cases which have come more particularly under my notice.

It will be well to cite a few typical instances:

1. A young married lady, aged 24, consulted me for a severe retroflexion of the uterus, which appeared to have set in shortly after her first labour, some months previously. She underwent treatment for this affection; and, in the course of it, one day her mother came with her, and inquired of me why it was that her daughter had become liable to this affection. I replied that, in all probability, it was due to a previous condition of weakness, and insufficiency of meat as an article of food. "Yes," said the patient, "that is very likely true; for when I married, and could please myself, being very fond of sweets, I took little ordinary food during the first year, but lived chiefly on sweets, and ate little or no meat at all."

2. A young lady, aged 18, suffered much from menorrhagia, and her condition became finally one of complete inability for exertion. It was ascertained that this patient had lived regularly as regards her diet, but that her diet consisted principally of bread and puddings; and, having a marked aversion to meat, she had taken very little for two or three years previously; and it is the fact that I was afterwards consulted by two of her sisters presenting uterine symptoms, and with a corresponding history as regards the previous dietary.

3. Again, a young lady of very active habits, but who hardly ever took meat, became very ill after an unusual exertion, was affected with severe sickness, which went on unchecked for nearly a year, and was finally so ill that her life was despaired of. Here, the previous dietetic deficiency was most marked. The patient was finally completely cured, the sickness being found to be a reflex symptom dependent on an uterine displacement.

These cases might be multiplied *ad infinitum*.

I have been much interested in observing, also, the effects of previous insufficiency of food in apparently predisposing patients to attacks of puerperal septicæmia. In the cases of this disease which I have seen in consultation, I hardly recollect having seen a case where the patient so affected had, during the pregnancy, lived fairly well; and the worst cases have been those in which the patients fed badly and insufficiently during the pregnancy, and had been fed on a gruel diet after the labour was over. In cases where severe sickness during the early part of the pregnancy prevents the proper nutrition of the patient, the system is liable to become much impoverished, and an insufficient dietary may be, and often is, the preliminary to a dangerous childbed.

With very few exceptions, and those exceptions only tending to prove the rule, it is, I hold, impossible to find patients suffering from chronic uterine disease, who have not undergone, at some former period, what may be termed a starvation process; and careful inquiry generally elicits the fact that the quantitative deficiency in the diet extended over a considerable period. In many cases, the patients are found to be still under the influence of a deficiency in this direction, and to be "eating," as the expression is, "next to nothing."

The period of life during which quantitative deficiencies in the dietary are most common, is the two or three years immediately following the arrival of puberty. The girl is at school probably, her appetite is bad, or the food is not palatable, or is deficient in important particulars, or, as I have found in some cases, she eats little in order to keep thin; the strength fails, the appetite diminishes, and a habit of taking little is formed—particularly little animal food is taken. Three or four years of the most critical stages of life are thus passed, a time at which the body should be growing fast, and to maintain this growth in adequate vigour, large supplies of nutritious material are required; instead of which, the supply is far below the normal standard. The result is a general impairment of vigour and of vital action. On the generative organs, the effect is, as I have observed in numbers of cases, most decidedly injurious; and the effect, in most instances, is of this kind, that the tissues of the uterus lose their normal firm healthy consistence. The further result is, that the pelvic organs as a whole, but particularly the uterus, undergo a subsidence in the pelvis; the uterus becomes liable to change of shape, and other alterations, more or less marked, in different cases. Slight exertions, slight accidents, or even moderate exercise, are, under these circumstances, liable to act most prejudicially on the softened and weakened contents of the pelvis. This is a faithful description of what I have observed to occur in very many cases. I refrain from pursuing the further history of patients so affected, as not falling within the scope of these remarks.

It is generally admitted by authorities on the subject of diet that nitrogen is the most essential of all foods, and that a certain amount—about three hundred grains—should be taken daily. In cases of chronic insufficiency of food, it appears that the diminution in quantity of food most frequently affects the nitrogen. Meat is the article of diet which, as a rule, is the source of the greater part of the needed amount of nitrogen, for, in England, at all events, meat is the popular article of food; and, in cases of chronic starvation, we mostly find that the quantity of meat taken is exceedingly small. "Never a meat-eater," "Do not like meat," "Have got out of the habit of taking meat"—such are common replies made to interrogations of patients under these circumstances.

No doubt, meat can be replaced dietetically by other foods containing nitrogen in sufficient quantity; but practically, owing to the habits of families, good substitutes for meat are not easy to find. The weakly one of the family is too often allowed to take her own course, and, if she does not take meat, often gets nothing sufficiently nitrogenous to answer the same purpose. Of all the nitrogenous foods, meat is admitted by all authorities to be the most easily digested, most easily assimilable, and most rapid in its nutritive action. Milk is, of course, a most valuable alternative food; but, in these cases of absence of sufficient meat, we do not find it has ever been taken in any such quantity as to make up for the deficiency; and the quantity of bread consumed, even supposing it to be pure and of good quality, is in such cases entirely inadequate to supply the required quantity of nitrogen. I need not allude to the effect of deficiency in the other constituents of the diet. It is sufficient for my present purpose to show that the nitrogenous elements, while they are of all the most important, are those which are markedly absent in the cases now under consideration.

There are few observations bearing on the subject now under discussion which can be quoted from published works. Professor Voit (*Untersuchung der Kost*, Munich, 1877) mentions an interesting fact. In a public institution, a home for girls, on which, he reports, the diet included an average quantity of 170 grains of nitrogen only, the girls appeared healthy, but their muscles were found to be weak, and menstruation was found to be delayed in many cases until the sixteenth or seventeenth year.

Parkes states that when the nitrogen is reduced to from 70 to 100 grains daily, the body gradually lessens in activity, and passes into a more or less adynamic condition, which predisposes to the attacks of all the specific diseases, especially malarious affections, typhus, and pneumonia, etc.

It must be assumed from what is known that if the minimum

quantity of nitrogen—which, for the sake of argument, we may put as low as 250 grains in the case of a young woman—be not given, a condition of weakness will soon be induced, and with greater or less rapidity, according as the quantity of nitrogen falls much or little below this 250 grains *per diem*. Thus it is easy to understand why, in a year or two, with an average daily consumption of only 100 grains of nitrogen, for instance, important modifications of the nutritive processes are effected, whereby there is produced a direct predisposition to disease.

It is now some sixteen years ago since I first publicly discoursed on the importance of nutrition in the treatment of disease. I have made it the basis of my practice for some years past, and four years ago I described the condition to which I have now again referred as “chronic starvation.” In the United States, Dr. Weir Mitchell has employed a system of rapid feeding, assisted by massage and electricity, for the cure of weakly, nervous sufferers, a system which has been found most successful, and Dr. Playfair has done good service in introducing Dr. Weir Mitchell's method into this country. The method in question is essentially a rapidly acting means of introducing nutritive material into the system, and it is dependent for its success on the fact that the principle of the curative influence and action of food is adequately recognised.

It is probable that it will be found, on further investigation, that particular deficiencies in the dietary result in deterioration of particular tissues of the body. The field for inquiry in this direction appears to be a large one. Deficiency in the quantity of meat taken, certainly, according to my experience, appears to have an enormous influence in predisposing to uterine ailments; but, no doubt, other organs of the body also suffer from deficiency in this article of diet. And I have abundant evidence that it plays a very prominent part in predisposing to the serious wasting diseases so often fatal in this country. I refrain from dilating further on this particular point, because such discussion would in this place be inappropriate.

The conclusion which is obviously suggested by the foregoing considerations, is the necessity for a greater attention to the question of diet, in the bringing up of families, than appears at present to be given to it. We all know that health and a good appetite usually go together. But it seems to be too frequently the case that, when the appetite is absent, such absence is taken as a matter of course, and receives no notice. It results from what has been stated, that absence of appetite may lead to most serious results. It is not immediately dangerous, but it is the first step possibly in the downward course. A continuously bad appetite constitutes a grave condition, and should be seriously regarded.

If it be a rigorously proved fact, that the human body is dependent for its existence, in a state of vigour, on an adequate and regular supply of food, it behoves us to take all possible opportunities of enforcing this great principle; and in making it known as a great and universally applicable measure and precaution in the prevention of disease.

CASE OF ENTERIC FEVER WHICH HAD TWO RELAPSES.—Dr Robt. W. Forrest, in a paper reprinted from *The Glasgow Medical Journal* for May, 1883, describes a case of enteric fever in which the interesting feature was “the unusual duration (*sic*) of the interval between the first and second attacks.” The primary attack ended on the 30th day, when the temperature became normal. After an interval of thirty-two days the temperature again rose. The first relapse lasted twenty-three days, and, after a further interval of seven days, a second relapse occurred. So long an interval between the primary attack and the relapse, as occurred in this case, is very unusual. Dr. Forrest formulates the question whether the first relapse was due to the primary infection or to a new infection. Very wisely he does not attempt to solve it.

BEQUESTS AND DONATIONS.—The Rotherham Hospital and Dispensary has received £175 further, making a total of £675, under the will of Mr. John Brown.—The Queen's Hospital, Birmingham, has become entitled to £100 under the will of Mr. Thomas Aston, £100 under that of Mr. Edwin Cotterill, and £100 under that of Mr. Thomas Lowe.—Mr. William Foster has given £105, Mr. John Foster £105, Mr. James Walton £105, and Mr. John Dawson £100, to the Royal Albert Asylum for Idiots and Imbeciles of the Northern Counties, Lancaster.—The Corporation of the City of London have given one hundred guineas to St. Mary's Hospital.—The Sussex County Hospital, Brighton, has received £104 9s. 7d. under the will of Mrs. M. Sayer.—Sir Philip Rose Bart., bequeathed £100 to the Hospital for Consumption and Diseases of the Chest, Brompton.

AN ADDRESS

DELIVERED AT THE OPENING OF

THE SECTION OF PHYSIOLOGY,

At the Annual Meeting of the British Medical Association in Liverpool, August, 1883.

By E. H. SCHÄFER, F.R.S.,

Jodrell Professor of Physiology in University College, London.

THE STRUCTURE OF THE ANIMAL CELL.

THE subject which I have selected to bring before you in this section is the structure of the animal cell. To physiologists and pathologists this is a subject of so general an interest as to need no apology for its introduction. Especially does this seem an appropriate time and occasion for its review, by reason of the fact that there is no single subject in anatomical or physiological science which has absorbed of late more attention than this one, and moreover because it is one in which I myself who have been honoured by being selected to preside on this occasion have for many years taken a particular interest.

In the first place, let us consider the views which were formerly held regarding the structure of the animal cell. The earliest notion was that expressed by its name, a hollow space enclosed by firm walls. The space was occupied by fluid, which might contain granules. Schwann (*Microscopical Researches*, 1839, Sydenham Society's Translation, 1847) succeeded in crushing a young cartilage-cell; it “suddenly shrank together, whilst a clear fluid streamed out, thus proving the contents of the cell to be fluid and transparent.” The wall is essential to the idea of the cell; even the lymph and pus-corpuscles have an envelope or cell-membrane enclosing the cell-contents. Mucus-corpuscles are exceptional in so far as that “an especial cell-membrane cannot be distinguished” in them; it may, nevertheless, still exist. The nucleus is a simple vesicle, filled with fluid; with, at the most, one or more solid particles—nucleoli—suspended in the fluid, it is not essential to the idea of a cell, or at least may disappear even from living succulent cells, it is often adherent to the cell-wall. This notion of the animal cell is an extension of that which had been already formed from observations on the vegetable cell, no independent line is taken, the facts regarding the plant-cell seem clear enough, the elements are large and conspicuous; what more likely than that the animal cell should possess the same structure?

The most conspicuous fact in this idea of the cell is the importance of the cell-membrane. This did not serve merely to inclose the fluid contents of the cell; it had a direct relation to nutrition and growth. Kölliker (*Manual of Human Histology*, 1852: Sydenham Society's Translation, 1853) expressly declares that “the power of growth is not simply innate in every organic membrane, manifesting itself when sufficient formative material is offered, but requires special conditions, which are only realised by the cell-membrane.” Further: “While the membranes grow by the attraction of material from the surrounding fluid, by virtue of their priority, they allow substances to penetrate into their interior. This filling, however, does not take place by the cells admitting every kind of matter indiscriminately, but they...take up one constituent (of the cytoplasm) and reject another.” And this selection is the property of the cell-membranes, for a little further on we read, “The cell-membranes do not act as mere filters, but allow one substance or another to permeate them, according to their chemical constitution, the constitution of the fluid which imbues them, their condition of aggregation, and their thickness.” This is not a mere process of endosmosis, although this also “must be taken into account as a condition in the absorptive action of cells, though hitherto it has been too freely appealed to, and cells have been too often considered as vesicles provided with merely indifferent porous membranes.”

Given such structural conditions—a vesicle filled with fluid, and enclosed by a structureless membrane—it was not easy to understand how changes were brought about in the cell-contents, although the occurrence of such changes was not to be ignored. Schwann assumed the existence of a special force—metabolic force—capable of producing such changes; but this assumption could not, of course, advance the explanation. Others were disposed to endure