AN ADDRESS

ON THE

TREATMENT OF FRACTURE OF THE PATELLA.

Delivered at the First Meeting of the Session of the Medical Society of London.

BY JOSEPH LISTER, F.R.S.,

Professor of Clinical Surgery in King's College.

SIR JOSEPH FAYRER, and gentlemen,—Some time ago, Mr. Holmes remarked to me that it would be well for me to place before the profession statistics of the operations which I had performed for fracture of the patella. And when you, sir, did me the honour to request that I should open this session of the Medical Society with a paper, it occurred to me that I could hardly do better than act on Mr. Holmes's suggestion. But, before entering on the strict subject of the communication which I have the honour to bring before you, it will be advisable to make some prefatory remarks regarding the circumstances that led me to it. In March 1873, my friend, Dr. Hector Cameron of Glasgow, recommended to my care, in the Edinburgh Infirmary, a case of ununited fracture of the olecranon. Dr. Cameron had formerly been my house-surgeon in the Glasgow Royal Infirmary, and I had afterwards for several years the great advantage of his assistance in private practice; and he reminds me that I had often expressed to him the opinion that the use of a metallic suture, antiseptically applied, which we had employed in ununited fracture of the shafts of the long bones, ought, in suitable cases, to be extended to the olecranon and patella. The patient to whom I refer presented himself to Dr. Cameron in the out-patient department of the Infirmary; and, as he had not at that time beds in the institution, and therefore could not operate himself, he sent him to me. He was a man, thirty-four years of age, who, five months previously, had received a blow from a policeman's *baton* on the left elbow. This occasioned great swelling, which seems to have concealed the true nature of the case from the medical man whom he first consulted. On admission, there was a considerable interval between the olecranon and the shaft of the bone; and, although the limb was muscular, it was comparatively helpless, as he could not extend the forearm at all without the aid of the other hand. On the 28th of the month, I made a longitudinal incision, exposing the site of the fracture, and, at the same time, bringing into view the articular surface of the humerus; and, having pared away the fibrous material from the fractured surfaces, I proceeded to drill the fragments, with a view to the application of the suture. The fracture was oblique from before backwards, as indicated by this diagram. I found no dif-

FIG. 1.

ficulty, with the proximal fragment, in making the drill appear upon the fractured surface at a little distance from the cartilage, (see

b, Fig. 1), but with the other fragment the obliquity of the position in which the drill had to be placed was so great that, instead of the end of the drill emerging at the fractured surface, as I had intended, I found it had entered into the substance of the humerus (d, Fig. 1). I therefore withdrew the drill, and substituted for it a needle (c d), passing the eyed end in first. Then, with a gouge, I excavated an opening (e) upon the fractured surface, opposite to the drill-hole (b) on the other surface, until the needle was exposed. Withdrawing the needle, I introduced a silver wire in its place, and I had no difficulty, by means of forceps passed into the excavation made by the gouge, in drawing out the wire. I was then able to pass it on through the other drilled opening, and thus the two fragments were brought into apposition. The ends of the wire were twisted together and left projecting at the wound. Healing took place without suppuration or fever, and the wire, was removed on May 19th, seven weeks after the operation. The wound made for its extraction soon healed, and the patient returned to Glasgow; and I afterwards had the satisfaction of learning that he was wielding the hammer in an iron shipbuilding yard with his former energy.

I have had two other cases of ununited fracture of the olecranon; and, as these are closely allied to the subject of my paper, I may refer briefly to them. One was a man forty-five years of age, incapacitated for his occupation as a plasterer by inability to extend the right elbow completely. The ununited fracture of nine weeks' standing was oblique laterally. It was treated as in the last case; the operation presenting no difficulty. It was performed on March 20th, 1878. The wound healed without suppuration, but the wire was not completely removed; for, the loop having broken near the twist, the twisted part was alone taken away, and the loop left behind. It never caused any inconvenience, and he afterwards wrote to us from his home at Bristol that he was able to follow his old employment.

The third case was that of a gentleman, 33 years old, who had consulted no fewer than eighteen surgeons on account of the weakness of his left arm, caused by ununited fracture of the olecranon. I operated on July 28th, 1881, paring the broken surfaces as in the other cases, using a chisel and hammer for the purpose; and, having drilled the fragments with a common bradawl, I brought them together with moderately stout silver wire. In this case, however, I did not leave the ends of the wire projecting from the wound; but, having given them one complete twist (or two half twists) cut the ends off short, and hammered the twisted part down flat upon the bone with this small hammer.

The advantage of this practice was strikingly exemplified by the difference in the course of this case from its predecessors. Instead of keeping him under treatment for several weeks until the wire could be removed, I was enabled to allow him to return, fifteen days after the operation, to his home in Wales, with a sound cicatrix; and, trusting to the connecting loop of wire, permitted him to use the elbow freely. I afterwards learned that he was able to drive a four-in-hand as well as ever.

The practice of cutting the ends of the wire short, and hammering down the twist upon the bone, is one to which I shall have to refer again in connection with my later cases of fracture of the patella. It is in every respect an advantage. The hammering down of the twist renders it more secure than if it is left projecting, to be moved by every shifting of the dressing, and perhaps broken, as in the second of the cases just referred to. We also get rid of a source of disturbance, and sometimes of considerable uneasiness, in the wound. The time of healing is greatly shortened, and the knowledge that the loop of wire securely holds the fragments in position allows the use of the joint to be commenced much earlier than when we have only the organic bond of union to trust to. The practice is also of the highest value in ununited fracture of the shafts of the long bones. The thickness of the wire must be proportioned to the force to which it is to be subjected. For the olecranon, that which I have here is amply sufficient, only about one twenty-fifth of an inch in diameter. For the shaft of a femur of an adult male, a piece like this, about one tenth of an inch in thickness, which I have had specially prepared with a view to a case of ununited fracture of the femur which we expect to operate upon in the hospital this week, is requisite, in order to resist with certainty the enormous force of the great muscles of the thigh. The pieces of bone which I hold in my hand were removed, in August 1881, from a case of badly ununited fracture of the femur. The patient was a gentleman from Rio de Janeiro. The fracture had occurred about the junction between the middle and upper thirds of the bone; and it had been so badly ununited, that the fragments overlapped very much, and also were

at a considerable angle with each other, so that the limb was extremely distorted, as well as much shortened. Bloodlessness of the operation having been provided for by elevation of the limb and the application of an elastic tourniquet, I cut down from the outer aspect of the thigh upon the seat of fracture. Then, with a periosteum-detacher, I separated the soft parts from the place of junction of the fragments; and, in the next place, I went through the extremely laborious process of cutting through the osseous union (which was of almost ivory hardness) parallel to the axes of the two overlapping fragments. This having been at length accomplished, and the soft parts still further detached, I found that I was able not only to get the limb into a straight position, but also, by very moderate extension, so to reduce the riding of the fragments that, by sawing off comparatively small portions, I was able to bring their extremities into apposition and apply my suture. This was done with wire of about this same thickness. The limb was put up at first with a long splint. I need not enter into details regarding the after-treatment; but I may say this, that he was a weakly man, and it was some months before absolutely firm union was obtained. It would have been extremely embarrassing to have had the ends of the wire sticking out from the wound all that time. On the contrary, it was a very great comfort to have no occasion to think about the wire; and ultimately he left for South America, with a perfectly straight limb, almost of the same length as the other, and, at the same time, with thoroughly firm union.

But to return to the immediate subject of my communication. Ever after my first case of ununited fracture of the olecranon, I was on the look out for a fracture of the patella to treat on the same principle. Dr. Cameron, however, anticipated me. In October 1876, being now full surgeon in the Glasgow Infirmary, he admitted a man with transverse fracture of the patella. He treated him, in the first instance, in the ordinary way, and dismissed him at the end of eight weeks with a pretty short and strong ligamentous union. Eleven days later, however, he reappeared, having ruptured the fibrous band by a violent movement during a state of intoxication. The fragments were then found widely separated. Dr. Cameron again treated him on ordinary principles for eight weeks, at the end of which time the fragments were still so widely apart, and the limb so feeble, that Dr. Cameron determined to cut down and apply the wire suture antiseptically. This he did on March 6th, 1877. On making a longitudinal incision, he exposed a condition of the parts, of which, through his kindness, I am able to show you a sketch, viz., a ligamentous union one inch in length, connecting pretty equal-sized fragments, with nipple-like projections extending from their attenuated margins much thinned by absorption. He cut away the fibrous material, and, having pared the edges of the fragments, and drilled them in two situations with a bradawl, he connected them with two sutures of stout silver wire (as shown here in another drawing), the ends of the wires being left projecting at the wound. At the same time, he introduced an independent drain into the joint. The wound healed without suppuration or fever; and, though osseous union was not obtained—which, as Dr. Cameron remarks in his report of the case, was not to be wondered at, considering the thinned state of the surfaces—yet he had the satisfaction of discharging the patient with close approximation of the fragments, and a thoroughly useful limb.

In October of the same year, 1877, a patient with transverse fracture of the patella was admitted under my care in King's College Hospital. He was a man forty years of age, who, while riding on horseback, had his horse stumble and fall. He was thrown over the horse's head, falling on the right knee. He could not rise, and was brought to the hospital. In the first instance, I attempted with this patient to bring the upper fragment down, so that it should be in contact with the lower. For this purpose I applied an apparatus, into the details of which I need not enter further than to say that it was so arranged, that the upper fragment, by means of weights and pulleys, was drawn down. Four days later, however, I found that there was still a quarter of an inch interval between the fragments, and I suggested to the patient the operation of cutting down and applying the wire suture. This, however, he would not then consent to, and preferred returning home to be under the care of his ordinary medical attendant. Eight days later, or fourteen days after the accident, he was readmitted, expressing a wish to be operated upon. On October 26th, I accordingly proceeded to operate, making a vertical incision, about two inches in length over the patella, exposing the fragments, which were then one inch apart. My inability to bring down the upper fragment into contact with the lower became explained when the parts were exposed; for there

were found between the fragments extremely firm coagula, with fibrous tissue, fascial and periosteal, mingled with them, constituting so firm a mass as to make it quite impossible for the two fragments to be brought into contact. The clots having been completely cleared away from between the fragments and from the interior of the joint, I applied a common bradawl in the middle line of the patella, drilling each fragment obliquely so as to bring out the drill upon the broken surface a little distance from the cartilage. Pretty stout silver wire was then passed through the drilled openings, and the fragments thus strung upon it were pushed firmly home, and so brought accurately into apposition. Before they were brought together, however, an arrangement was made for the drainage of the joint. This was done on the same principle in all the cases that I have to record, and I may therefore describe the matter once for all. A pair of dressing forceps, with the blades closed, were introduced from the wound made into the anterior part of the joint to the most dependent part of the outer aspect of the articulation. The instrument was then forcibly thrust through the synovial membrane, the fibrous capsule, and the fascia, until the point of the forceps was felt under the skin. An incision was then made with a knife through the skin upon the end of the dressing forceps, so as to allow it to protrude. The blades of the forceps were then expanded so as to enlarge the opening which they had made in the deeper structures without risk of causing hæmorrhage. The drain was then seized in the forceps that protruded through the wound, and drawn into the joint. The ends of the wire were now twisted together, and the twisted ends brought out at the wound, which was closed with sutures and a small drain inserted. I need hardly say that in this case, as in Dr. Cameron's, antiseptic treatment was employed throughout. It is unnecessary for me to enter into details as to the progress of this case. We have here the temperature-chart for as long as it was thought worth while to have it recorded, and you will see that it indicates, after a little temporary disturbance immediately after the operation, an entirely afebrile condition. The wounds healed without any suppuration. At the end of eight weeks, the wire was removed by an incision through the cicatrix. Eight days later, the wound made for the removal of the wire had healed. At the end of ten weeks from the operation, the patient was allowed to get up, and, though no passive motion had been employed, he could move the limb freely through an angle of about thirty degrees. Two days later, he was discharged, and, unfortunately, nothing has been heard of him since. I saw him once in a cart a few days after he was dismissed, but I have not been able to learn any further tidings of him. This, I believe, is the first instance of a recent case of fracture of the patella being treated by wire-suture antiseptically applied.

My next case occurred two years later. William T., a coal-porter, thirty-seven years of age, was admitted December 13th, 1879. He was a muscular man. The patient slipped on December 9th, while carrying a sack of coals, and felt something gave way in one knee. On endeavouring to rise, he found himself unable to do so. On admission into the hospital, the right patella was found fractured transversely, the interval between the fragments being about an inch. There was a considerable amount of effusion into the joint. On December 15th, that is to say, six days after the accident, I proceeded to operate, making a longitudinal incision, as in the last case, about two inches long, over the patella. The lips of the wound being held apart with blunt hooks, a hole was drilled in each fragment in the median line. Stout silver wire was passed, and secured by four half-turns. A drainage-tube was introduced, as in the last case. And here, again, there is no need for my entering in detail into the reports. We have before us the temperature-chart. For the first fortnight after the operation, there was what we may call an absolutely afebrile state. During the rest of the time of his residence in the hospital, there were two accidental elevations to 100.5°, but nothing to indicate anything serious. The wound healed, as in the last case, without any suppuration. Six weeks after the operation, he was allowed to get up. Eight weeks after the operation, an incision was made through the cicatrix for the removal of the wire, the loop of which was cut, and the wire withdrawn. On February 23rd, that is, a fortnight later, the bandages which had been previously applied to the leg were removed, and it was found that he could bend the limb to a right angle; he could walk well, and was able to kick. On February 22nd, 1883, this patient showed himself. We then took the following notes.

"The patella seems perfectly natural, except a trifling irregularity of outline on one side. It is evidently osseously united. The movements of the joint are perfect, from complete flexion to extension. He can kick vigorously. He says the joint is just as trustworthy as ever.

He frequently carries a weight of 220 lbs. for upwards of 100 yards, and he walks without the slightest limp."

[The patient was now introduced and exhibited to the Society. He said his limb was as good as ever, and that it never failed him.] I have not seen the man since the time I have referred to. Except for the linear cicatrix, no one would be able to tell that this patient had had anything wrong with his knee at all.

The next case was one of ununited fracture. The patient, Joseph R., aged 22, was admitted into the hospital on September 27th, 1880. He is a soldier, and stated that, on June 3rd of that year, while running across a green, he slipped upon a piece of turf, and, in trying to recover his balance, he snapped his knee-pan and fell. He was taken to a Military Hospital; and the limb was put in splints, which were kept on for seven weeks, a starched bandage being applied after their removal. On admission, the fragments were found separated by an interval of three-quarters of an inch. There was a firm fibrous band of union between them. The knee, however, was quite stiff in consequence, apparently, of the rigidity of the extremely atrophied quadriceps extensor. He was quite unable to walk. If he attempted to do so, he held the leg in his hand. He complained of frequent uneasy sensations in the knee. On the 22nd of October, I cut down over the patella, as in the former case, making an incision about two inches in length. Having cut away the fibrous tissue between the fragments, I refreshed the osseous surfaces, and then, having provided for drainage as before, drilled the fragments, and drew them together with stout silver wire. Nothing occurred worthy of notice until four weeks after the operation, when I proceeded to attempt passive motion. Without chloroform, this proved impossible; but, under chloroform, I used considerable force, when, the rigid quadriceps extensor refusing to yield, the twist of the wire gave way; the cicatrix, which had quite healed except where the wire projected, opened; and there was heard a sound of air being sucked into the joint, the fragments becoming, at the same time, widely separated. I injected the joint with an antiseptic solution, a procedure which I candidly confess I should now regard as probably superfluous. However, such was done; and, six days later, when all disturbance caused by this second injury and the irritation of the antiseptic injection had passed off, I proceeded to operate upon him a second time. Chloroform was again administered, and an incision similar to the former one was made, the cicatrix being laid open. I then found that the ends of the wire were lying in place, ready to be twisted together. I found, at the same time, a considerable mass of coagula present. These I cleared away from between the fragments and from the joint, twisted the wire again, and thus we had a second operation in one patient. The after-progress was as in the other cases. Here, we have a doubly long temperature-chart, because we have two cases in one, so to speak; but it will be seen that there was no deviation whatever from the normal state; neither as the result of one operation nor the other has any febrile condition as regards temperature been produced. In due time the wire was removed, and the wound made through the cicatrix having healed, he was discharged eight weeks after the second operation. He was then able to walk, but with a stiff knee, with scarcely any mobility. I did not attempt any more passive motion after previous experience, and while I was well pleased to see that he could walk with a stiff knee, whereas he could not walk at all on admission, I hardly ventured to hope for anything better. However, on February 22nd, 1883, he presented himself, when the following report was made. "The patella is perfectly natural, except a little irregularity of the two borders opposite the seat of the fracture. The surface is quite smooth. There is no interval between the fragments. There is evident osseous union. The degree of flexion is increasing. He can bend to an angle of 60°, and extend again completely. He laid aside his stick last summer, and returned to his work as a gardener seven months ago. He can do anything except kneel. In getting over a paling, he has to throw the leg over in a partially extended position. He walks with a barely perceptible limp." We may now say that he walks with no limp at all.

[The patient was introduced and exhibited. He could now bend his knee nearly to a right angle, and said that he was on his legs all day, and was quite equal to his work as a gardener.] The great interest in this case seems to me to lie in the improvement that has occurred without any passive motion on the part of the surgeon, and as a mere result of the natural actions of the limb; from such a very rigid state of things as there was when he left the hospital to the condition of mobility which we now witness.

The next case was also one of ununited fracture. The patient, Martha F., aged 43, was admitted November 5th, 1880, on

account of an accident which had happened eight weeks before, when she felt something snap in her knee, while trying to save herself from falling. She was unable to walk or move the joint, which quickly swelled up to a large size. No splints had been used, and no treatment of any kind employed. On admission, there was fullness over the knee, and a transverse fracture of the patella to be discovered in the middle of the bone. The fragments were separated one inch. There was fluid in the joint, as indicated by fluctuation. On November 12th, I operated as in the last case, paring away the fibrous material between the fragments, and refreshing the osseous surfaces with cutting pliers. The fragments were then drilled obliquely in the middle line, and brought into apposition by means of a suture of stout silver wire. This, however, could only be done when the limb was raised high into the air so as to relax the quadriceps, she being a stout woman, with powerful muscles. Of course, antiseptic dressing was employed; and, as was done in all these cases, the limb was placed in a trough of Gooch's splint, with the upper end oblique, corresponding to the line from the tuberosity of the ischium to the great trochanter, and the lower end excavated in the form of a horseshoe, while the horns of the horseshoe were well padded, to support the sides of the foot. With regard to this patient, again, I have to show a temperature-chart free from any febrile indication. In eight weeks after the operation, the patient was allowed to get up, and she walked about on crutches. She was discharged three weeks later, able to bend the knee slightly, lowering and raising the foot three or four inches.

On February 14th, the patient was readmitted, to have the wire extracted; and, five days later, she was finally discharged. It will be remembered that this was a case of ununited fracture; and that, when the patient left the hospital, she could only move the foot through a distance of about four inches. On February 22nd of this year, we had an opportunity of seeing her, and then the following note was taken. "There is perfect union of the fragments. She can walk from Drury Lane to Billingsgate and back, and walks without a limp. She can bend the knee to a right angle, and says the mobility increases every day. From the position at a right angle, she can raise the leg in a perfectly natural manner to the extended position. She is a stout heavy woman. She cannot kneel."

[The patient was now introduced and exhibited. There seems to be a perfect patella; there is nothing abnormal to be felt about it. She says she can do everything except kneeling.]

The next case was one of recent fracture. William G., 62 years of age, admitted June 21st, 1881, a healthy man, but a pretty hard drinker. On the morning of his admission he slipped, put out his right leg to save himself, fell, and could not rise. His leg doubled under him. On admission the patella was found fractured transversely, with one inch of interval between the fragments, which could be brought together with difficulty. The knee was considerably swollen. On the 24th, that is three days after the accident, I operated upon him. The operation was conducted precisely as in the cases that have been before described, except that, unlike the two last, there was no necessity for refreshing the fragments. It was merely necessary to take away the clots of blood and any effused fluid from the interior of the joint. A few hours after the operation this patient became delirious, and we were apprehensive of delirium tremens. However, he was quieted with a dose of opium, and gave no further trouble. The temperature-chart here, as in the last case, exhibits entire absence of febrile indication. In this patient, for the first time, the ends of the wire were cut short, and the twist hammered down. The wounds healed without suppuration. Exactly six weeks after the operation he was allowed to get up, and I had the satisfaction of exhibiting him before some of the members of the International Medical Congress, and showed them that he could raise his limb, from a position of flexion almost at a right angle, to the completely extended posture. I have not seen him from that time until to-day. Having learned that he was employed in a Birmingham establishment, I wrote to my friend Mr. Chavasse of that city, who replied as follows. "I saw G. this morning, and have made arrangements for him to leave by the 2 o'clock train, to-morrow afternoon. He will be at your house just before 8 o'clock. In case he should not keep his appointment, there is a faint linear cicatrix present over the joint. The knee can be flexed as well as the other. There is bony union of the patella, a faint ridge being felt on the point of union. No suture is to be detected by the touch. At his work, as a stamper, all day long, he works a pulley with the affected limb, which raises a weight of 60 lbs. In damp weather he feels very slight inconvenience in the site of the

old fracture." Happily our patient has kept his appointment, so that we are able to see him to-day.

[The patient was introduced and shown. He said he could do anything with the limb the same as with the other. He worked a stamp-hammer, and he could work that hammer, weighing one hundredweight, all day long. There was no difference whatever between the movements of the two knees.]

There is a barely perceptible cicatrix, and he has a perfect patella.

My sixth patient was a woman, Elizabeth C., fifty-seven years of age. She also was a recent case, admitted on October 21st, 1881. Going downstairs, she fell with her left leg bent under her, striking the knee against the ground. She was brought at once to the hospital. The left knee was very much swollen from effusion, hot, and tender: the fragments of the patella were felt, separated from each other about one inch. I believe it is generally wise to let any distinct inflammatory appearances that exist, as the immediate result of the accident, pass off before the operation. In the former case, there were no such, or scarcely any, therefore we operated three days after the accident. In this case, they were manifest, and we allowed a week to pass. The operation was performed just as in the last case. The ends of the wire were twisted with two half-turns, that is to say, one complete turn. The ends were cut short, and then, with the small hammer, the twist was hammered flat down upon the patella.

Here we have the temperature-chart. As we very commonly find, in cases antiseptically operated upon, the first effect of the operation is a depression of temperature. On the evening of the first day, there was a depression; on the following day, a little rise, but that rise only reached 101°; after that, there was nothing to indicate anything febrile. Two weeks after the operation, passive motion was commenced. The wounds were then so very nearly healed, that the milder antiseptic boracic lint was used for the superficial sore which remained. Three weeks after the operation, the limb was bent to a right angle; it was painful to her, but caused no disturbance of the fragments. Four weeks after operation, passive motion was again employed; the wounds were completely healed. Six weeks after the operation, the patient was allowed to get up, and could already walk fairly. Three days later, she was discharged; she could then walk very well. On the 23rd of February of this year, we took this note regarding her. "The patella appears perfectly natural, with the exception that the wire is felt, causing the slightest projection under the skin, which, however, moves freely over it. The movements of the joint are perfect: from complete flexion at a very acute angle to perfect extension. She kneels as she used to do, and only occasionally the wire comes into contact with anything, and then she feels the skin over it tender. She can walk any distance, as before the accident, and does so without a limp." [The patient was now introduced to the Society, and confirmed in every respect the previous report.]

My last case is a man, sixty-seven years of age, John P., admitted into King's College Hospital, on January 6th of the present year. He had fallen from an omnibus on the previous day, striking his knee upon the ground. There was transverse fracture of the patella, and great swelling of the joint. Six days after the accident, I proceeded to operate. On making a vertical incision and exposing the seat of fracture, I found a condition of things which, possibly, had I known it, might have induced me to abstain from operating. I found that the lower fragment was very small, and was comminuted. There was one entirely loose piece, as big as a filbert, and another about half that size, which, of course, had to be extracted. Such a condition, taken in connection with the advanced age of the patient, did not promise well for satisfactory union. Having sponged the clots out from between the fragments and from the joint, and established a drain, I drilled; but as the lower fragment was so exceedingly small, I was obliged to drill, not through it, but through the ligamentum patellæ; and, by this means, I was able to bring the large upper fragment into contact with the raw surface below. Now, however, I do not regret having operated upon this poor man. You will see that his temperature-chart is free from any febrile indication; and the wounds healed without any supuration. Here, as in the last two cases, the ends of the wire were cut short, and the twist hammered down upon the patella. Three weeks and four days after the operation, the wound was nearly healed. The knee was bent through an angle of thirty degrees, and from that position the patient could himself raise it to complete extension; a thing which, without the wire, and without our feeling that the wire was in a very secure form, we should not have thought of permitting. Four weeks after the operation, the wound was

healed. The knee was then bent to an angle of forty degrees. Eight weeks after the operation, the patient was discharged, having been for a considerable time before (I have not got it exactly recorded when) allowed to walk about the ward, and walking well. You will observe, gentlemen, that this is my most recent case. He has not had the same time for improvement that the others have had.

[The patient was now introduced, walking up the room without the slightest limp. He said he was getting on "first-rate." He found the leg continue to get stronger, and he was able to move it better. He noticed that it was getting stronger every week; he might say every day.]

I confess I am surprised to see how complete a patella this man has, considering how extremely small the lower fragment was. There must have been considerable new osseous formation from the periosteum, because here we feel the central part of the wire, and yet between that and the lower border there is what appears to be a substantial lower fragment connected with the upper one by osseous union. Now we may say he has a perfect patella.

These, gentlemen, are all the cases of fracture of the patella on which I have operated, and I consider it fortunate that I am able to bring before you six out of the seven.

I should like now to say a few words as to the method of operating. The wire employed should be, as I have said, pretty stout, about one-sixteenth of an inch in diameter. I have not found it needful to use more than a single suture of such wire. It is applied in the vertical plane, in the course of the longitudinal incision over the middle of the bone; and in recent cases no dissection of the soft parts from the patella is necessary. It seems important that the cartilaginous surface of the bone should be left quite smooth, or, in other words, that the fragments should be exactly at the same level at their lower part. We cannot be perfectly sure, when we drill, that the bradawl will come out exactly at a corresponding point on the two surfaces. Supposing that on one side the instru-

FIG. 3.

ment should have come out too far down, it may be into the cartilage (as at *d*, Fig. 2) instead of a little above it. We do not regard that at first, but pass the wire through each drill-hole the moment the drill is withdrawn, and then on that side on which the hole has come too far down, by means of the bradawl, we simply chip away a little of the material that is above the wire until the wire comes to be in a position exactly opposite to the hole on the other side, leaving a gap below, as indicated by the dark shade at *d*. This is a perfectly simple matter; at the same time, it might possibly not occur to anyone during the operation. Here (Fig. 3) we have the wire represented twisted, and the twist hammered down. The twist always goes to one side, and, being on the other side in this instance, is not shown in the section represented by the diagram.

I should like, gentlemen, with your permission, to make some few general remarks. I think it must be admitted that these cases show that the mode of treatment which I have recommended, when

applied to recent transverse fractures of the patella, affords a means of restoring the joint to, practically, a perfectly natural condition, provided only that no disaster occurs. That, however, is a tremendous proviso, and no one is more conscious of it than myself. Before I made the incision in the first case that I have recorded to-night, I remarked to those who were assembled in the theatre that I considered no man justified in performing such an operation, unless he could say, with a clear conscience, that he considered himself morally certain of avoiding the entrance of any septic mischief into the wound. Supposing, on the other hand, that a man can say that with a good conscience, then I conceive that he is not only justified, but bound to give his patient the advantages that we see are to be derived from this method of procedure. We know, of course, very well that, by ordinary means of treatment, patients often recover with exceedingly useful limbs. Every now and then, osseous union is obtained; it is a thing which I used to pride myself formerly on striving to get, and I have got such a thing, but it was rather rare, and it was obtained by a very tedious process; and, if ligamentous union occurred, we never had any security that what was a very short ligamentous union when the patient was discharged, might not be a long ligamentous union at a later period. A gentleman consulted me with transverse fracture of the patella a couple of years ago. He happened to come to me just as I was about to start for my autumn holiday. I did not care to operate upon him, and throw the responsibility of the after-treatment upon another, and it happened that the case seemed especially favourably circumstanced for osseous union. There was nothing between the fragments; the two could be brought into apposition with the utmost facility. I applied an apparatus which appeared to have the effect of maintaining perfect immobility of the fragments; and, when I came back from my holiday, I found them as I had left them, with no interval between them, and I hoped there might be osseous union. Eight weeks after the accident, the patient was allowed to get up and walk, and I was told that he was walking exceedingly well. Six weeks after this, I asked him to come and report himself, and I was sorry to find that there was already half an inch of separation of the fragments. He was then about to start for India. What the separation may be now, I do not know. It is true, that there may be a very useful knee-joint, with a very considerable length of fibrous union. Still, when there is a great length of fibrous union, the knee is not equal to the original.

Some of our experience with these cases, where we had the opportunity of inspecting the actual state of things, will serve to explain the uncertainty of the results of ordinary treatment. We have found, for instance, such dense masses of clot so mixed with fibrous tissue as to make it quite impossible that the fragments should be brought into apposition. I found in one case, also, such a tilting up of one of the fragments that it would have been perfectly impossible to get the osseous surfaces in contact by any other means than direct operative procedure. Considering, therefore, the great inconvenience which results in many cases when the treatment is conducted on ordinary principles, I believe that if we can really say that we are morally certain that we do not subject the patient to risk, we are in duty bound to give him the benefit of this method. It has been said that it is not justifiable in recent cases, though it is justifiable in ununited cases where there is an useless limb. I must confess that if I believed I was subjecting a patient to serious risk to life, I should not feel justified in operating on the ununited case, and the ununited case is in every respect worse as a subject of operation than the recent. The ununited case has the fragments probably dwindled by absorption, and these fragments, already dwindled, have to be pared. There is a separation, it may be a very considerable separation, and this has to be overcome, it may be by dividing the quadriceps extensor, an operation of difficulty; and in proportion to the length and difficulty of the operation is the chance that the surgeon may forget some point of importance with regard to the antiseptic element. Then, again, when you come to divide the quadriceps extensor, you divide a very vascular structure, and you may have hæmorrhage; and, further, when the fragments are brought into apposition they very likely are at considerable tension, and the tension may be apt to cause, through the nervous system, an inflammatory disturbance, and this tends to weaken the parts, and to diminish the power of resistance by which the natural tissues are able to combat the entrance of septic agencies, even though they be in contact with the part. With the recent case, on the other hand, everything is favourable. We have a wound involving no bleeding; and there is no need to pare the fragments. All you have to do is to sponge away the clots, and the surfaces are ready

for coaptation. The drilling is a matter of the utmost simplicity. It is an operation of no difficulty, it does not take long; it does not cause anxiety to the surgeon; there is no shock to the system, and no tension. In every respect, the circumstances are favourable as regards the operative procedure. And then, when we come to consider the chances of successful antiseptic management, if there is in the whole body a situation which is well adapted for antiseptic treatment it is this; for of all the conditions requisite according to our present methods of procedure, that which is most important is that the skin on all sides round about the wound should be able to be amply overlapped by the antiseptic dressings. Here we have the wound in the middle of a long limb; from the groin to the foot we may have our antiseptic envelope. Then again we have this envelope surrounded with a secure bandage, and the bandaged dressing encased in a splint, and even if you come to have the patient delirious, as one of my patients was, or supposing a patient to be very curious, as some patients will be, it would puzzle him to get the wound exposed under these circumstances. Now there are wounds so circumstanced that you cannot well guard against this risk. I had a gentleman lately under my care with a psoas abscess; he was very intelligent, and seemed duly impressed with the importance of the antiseptic management; and yet his brother, who was a medical man, coming in one day, saw him drawing the dressing aside, and peeping at the wound. Now a man cannot peep at a wound in connection with a fracture of the patella; it is so circumstanced mechanically that he cannot do it; and I believe that if we use the means that we have now at our disposal, we may say, with a safe conscience, if we use them aright, that we do not subject the patient to risk, not to anything like so great a risk as patients used to be subjected to not many years ago when they had fatty tumours removed in general hospitals in London. We must all of us remember cases in which, after such operations, erysipelas or diffuse suppuration came on, or some other "unhealthy action," which, of course, was nobody's fault, but the patient died.

I have referred to a case of ununited fracture of the olecranon where eighteen surgeons had been previously consulted. I trust no one here will suppose that I mentioned this circumstance for the purpose of glorifying myself. I mentioned it in order to emphasise what I believe to be the truth, that by antiseptic means we can do, and are bound to do, operations of the greatest importance for our patients' advantage, which, without strict antiseptic means, the best surgeon would not be justified in recommending. How wise those eighteen gentlemen were in counselling against operative interference, provided they were not prepared to operate strictly antiseptically, I think we must be all agreed. As regards the operative procedure in that case, it was of the most simple character; any first year's student could have done the operation exactly as well as myself; and, therefore, I trust I shall not be misunderstood by its being supposed that I came here to extol my own skill. That which justified me in operating in that case was simply the knowledge that strict antiseptic treatment would convert serious risk into complete safety.

I should have liked, if time had permitted, to have said a few words as to what seem to be the essential points as regards antiseptic treatment. If I say any words at all now, they will be exceedingly few. I should just like to make this remark, however, that now-a-days antiseptic treatment is not a very complicated business, either in theory or in practice. First, as to theory, we do not require any scientific theory to enable us to believe in antiseptic treatment. You need not believe in the germ-theory at all; if you are not convinced of the truth of the germ-theory of putrefaction and of septic agencies generally, no matter whatsoever with reference to antiseptic practice. All you have to believe is that there are such things as putrefaction and other septic agencies, and that our wounds are liable to these, and that they are very pernicious, and that these things come from without, and that we have the means of preventing them by various chemical agencies. That is all that we require; and I think anybody who knows the present state of surgical practice must admit these to be truisms. It has sometimes been a great grief to me to think that, because gentlemen are not convinced of the truth of the germ-theory out and out, therefore they lay aside antiseptic treatment altogether. And then as to practice; it is not a very difficult thing to wash your hands in a carbolic solution, and have your instruments in this carbolic solution for a quarter of an hour before you operate. It is not a very difficult thing to wrap round the limb a suitable envelope of antiseptic material. What I believe to be one of the most important things of all is, strictly to maintain this rule inviolate, which I insist upon with my dressers, and which I confess I have insisted upon more of

late years than I used, and that is, always when we change a dressing, invariably first to cover the wound with something pure; not to wash the surrounding parts with antiseptic solution, and then, after this has been done, put a dressing on the wound; but before we begin to defile the lotion at all, put on the wound what is pure, and, last thing of all, wash the surrounding parts, which, though they look the same to our eyes, are different *toto caelo*. The edges of the dressing are septic; the wound, if it is as it ought to be, is aseptic. I have known such a thing, for instance, as for a gentleman, in dressing a stump after amputation of the thigh, to wash the perineum with a rag dipped in the carbolic lotion one in forty; and then, having so washed the perineum, immediately to squeeze the rag over the wound. Gentlemen, that makes you laugh; but I assure you these are the kind of things that are constantly going on, and disasters happen in consequence; and gentlemen with whom things go wrong invariably say that with them everything has been perfectly done—a thing which, for my part, I am always loth to say. I am not likely to have many more years of active surgical work; and I have felt that when you, sir, gave me this opportunity, it was my duty to speak what I believe to be the truth; for I feel it to be a grievous thing that patients should be hurried out of their lives or deprived of usefulness of limbs, simply for want of sufficient earnestness with regard to the endeavour to obtain complete exclusion of septic agencies from wounds, according to our present lights and our present knowledge. Gentlemen, I thank you most heartily for your cheers; for there was a time when such remarks might have met with a different reception.

A NEW METHOD OF PERFORMING RESECTION OF THE KNEE-JOINT.

By W. MACFIE CAMPBELL, M.D.,
Surgeon to the Liverpool Northern Hospital.

WITH regard to Mr. R. Davy's clinical lecture on this subject, in the JOURNAL of October 20th, it seems to me that the claims to priority in special modes of operation are often made without due consideration.

On November 20th, 1882, I excised the knee-joint of a woman, aged 35, for acute ulceration of the cartilages. The femur was sawn obliquely before and behind, removing the cartilage completely, leaving a wedge-shaped extremity. A corresponding form was cut in the head of the tibia to the depth of about an inch. The advantage obtained was complete antero-posterior immobility. She was discharged cured on February 5th, 1883, sixty-seven days after operation.

This case was shown at a spring meeting of the Liverpool Medical Institution, and the method was familiar to the surgeons present, as appeared from the subsequent debate. To whom is due the original idea, I know not; certainly not to myself, as I had read it shortly before the operation, but where I cannot remember, though I think in some American journal.

I may add that this excision is only possible in cartilaginous disease, or in such a case as Mr. Davy has described, as sufficient healthy bone must be obtainable, and I, for one, would object to remove as much of the femur as he figures.

A CENTRAL site has been purchased in Cambridge by Mr. Reynolds Rowe, upon which he intends to build and endow a large and free open church dedicated to St. John of Jerusalem. The scheme includes an infirmary for the use of members of the University; a training-school and home for nurses ready for engagement in private houses, and an ambulance centre for instruction.

PRESENTATION.—Mr. Edward Walter Witten, M.R.C.S. Eng., upon resigning as Superintendent of the Medical Mission Dispensary, Brighton, has been presented with a gold pencil-case, and a text-book with the names of his co-workers and patients written in, and inscribed, "Presented to Mr. E. W. Witten, as a small token of gratitude and esteem from his co-workers and the people for whom he has laboured. October 18th, 1883."

DONATION.—Since the re-opening of Eaton Hall, the Duke of Westminster has charged all visitors 1s. for the privilege of inspecting the house and gardens. His Grace has recently sent the whole of the proceeds from 11,200 visitors, amounting to the handsome sum of £560, as a donation to the funds of the Chester Infirmary.

AN ADDRESS ON COLLECTIVE INVESTIGATION OF DISEASE.

Delivered at a Meeting of the Lancashire and Cheshire Branch.

By WILLIAM ROBERTS, M.D., F.R.C.P., F.R.S.,
Professor of Clinical Medicine in Owens College, Manchester; Physician to the Manchester Royal Infirmary and Lunatic Hospital.

GENTLEMEN,—Your presence here in such considerable numbers is a sign of the interest which you take in the movement. It is also a sign, I am bound to say, of admiration for our distinguished visitor, Mr. Jonathan Hutchinson.

We are in some sort under a special obligation in this district to take exceptional interest in this movement, because the initial idea sprang up in our midst. To our friend and neighbour Dr. Arthur Ransome is due the first inception of the plan of collective investigation. In 1864 he read a paper at the Cambridge meeting of the British Medical Association "On the Need of Combined Medical Observation," and he there suggested that the machinery of the Association should be utilised for that purpose. Two years ago, the matter was taken up afresh and independently by Dr. Mahomed. Both gentlemen are to be congratulated on the present flourishing prospects of the movement. It is not necessary for me to go over again the arguments in favour of this mode of investigation—this work has been exhaustively done by Professor Humphry, Sir William Gull, Sir James Paget, and others, whose addresses have been published in the BRITISH MEDICAL JOURNAL. Indeed the time is rather gone by for argument; the plan is actually on trial and in operation, and the first fruits are already garnered.

I have before me the first volume of a series of reports intended to be issued by the General Committee, and I may be permitted to compliment the editors, Professor Humphry and Dr. Mahomed, on this new birth of a wholly new kind of medical literature. I can only regret that this little volume is not in the hands of every member of the Association. The object in view is only half accomplished unless these reports as they come out pass into the hands of every member, or at least every contributor, by some automatic process without the trouble of separate ordering and paying. I hope the Committee will be able to see their way to do this. I may say that I was one of those whose confidence in this mode of investigation was not strong, but after a careful examination of the first number of the *Collective Investigation Record* my opinion has been greatly modified. No one can read it attentively without acquiring a feeling almost amounting to a conviction that the movement is one of real importance, and that it will result in a large addition to our knowledge that could not have been got in any other way. I should like to refer for a moment to the admirable report on the communicability of phthisis drawn up by Dr. Burney Yeo. I think no candid person can read this report, and the detailed evidence on which it is based, without coming to this practically important conclusion, that no healthy person should be permitted to occupy the same bed with a sufferer from pulmonary consumption, and that no person with a hereditary predisposition to tuberculous disease should be allowed to have continued and intimate personal contact with a phthisical patient. And I cannot help adding that a grave responsibility would in my opinion be incurred by a medical man knowing this evidence, no matter what his theoretical opinions may be, who would permit such cohabitation and close personal contact.

One of the most promising fields of labour for combined observation is the self-study of disease by medical men. In regard to serious organic fatally tending disorders, this would, indeed, be a task too tragic for contemplation, but in regard to a number of minor ailments this method is admirably adapted. Take, for example, the dyspepsia of healthy persons, or primary dyspepsia. Judging by the number of books written on the subject one would imagine that we knew a great deal about dyspepsia.

I am satisfied that, of exact knowledge, we have very little. The dyspepsia of healthy persons is a very common and peculiarly human disorder; it causes an immense amount of misery, but it kills no one, and the pathologist ignores it. The several types of it have never been scientifically defined, and its case-history remains un-