nitric acid, which I now never carry about with me, it is perfectly harmless, even if it escape from the bottle.

In testing with the powder, about one third of a grain, equal in bulk to a peppercorn, may be shaken up in a test-tube, with a column of urine about one inch in height. As the powder dissolves, the urine becomes turbid with coagulated albumen. The object is to add as much of the powder as the urine will dissolve, and no more. In the absence of a test-tube, the powder may be stirred up with the urine in a wine-glass. The solution of the picric acid in the urine, and the coagulation of the albumen, are quickened by heating the tube over a spirit-lamp or candle, or by immersing the tube or glass in hot water.

The dry picric acid should be used in the form of fine powder, not in crystals. One of my correspondents was perplexed by finding that crystals of the acid dropped into a highly albuminous urine became surrounded by a film of coagulated albumen, and, becoming caked together, formed a dense precipitate. This can never occur if the fine powder be shaken up with the urine.

Another convenient mode of using the powder at the bedside is to add twenty or thirty minims of water to the peppercorn bulk of the acid in a tube, and quicken the solution by the application of heat; an equal bulk of urine is then gradually added to the hot saturated solution, when albumen, if present, is at once detected. It has been objected to the picric acid solution that it stains the fingers; but this stain, unlike the yellow stain caused by nitric acid, which penetrates into the skin, and remains for several days, is merely a surface-stain, and is immediately and completely removed by washing with soap and water.

Dr. George Oliver has devoted much time and labour to a comparison of the various tests for albumen, and also to the preparation of test-papers for bedside use. His observations agree with my own, that the three most delicate liquid tests for albumen are solutions of potassium-mercuric iodide, picric acid, and tungstate of soda. He finds "these three albumen-precipitants of equal keenness;" but I have found the tungstate of soda decidedly inferior to the other two. In the form of paper-test, Dr. Oliver states that picric acid is "the weakest of the series" (*Lancet*, Jan. 27th and Feb. 8th, 1883). The explanation of this result is that, in consequence of the comparative insolubility of picric acid, the small slips of paper dried after immersion in a saturated solution do not retain sufficient of the acid to render them a satisfactory means of testing. But there is no need for picric acid papers. The main advantage of the test-papers consists in their ready portability; but this advantage is shared equally by the picric acid powder, which, moreover, possesses these advantages over any forms of test-papers, that it requires no elaborate preparation or the addition of any other agent, that it is therefore cheaper and more simple, that it affords a more ready and trustworthy indication of the amount of albumen, and, lastly, as I shall show you in a future lecture, that, in combination with caustic potash, it forms an admirable qualitative and quantitative test for sugar.

INSANITY IN FRANCE.—One of the most striking features of modern French life is the rapid increase of insanity; the number of cases of which, and especially those induced by alcoholism, is becoming larger each year. During 1882 there were 13,434 admissions into the asylums, of which 10,184 were new cases, the total number under treatment in the year being 58,760, of which about 27,000 were men and 31,000 women, showing that females are the most liable to the disease. That they are less influenced by treatment is shown by the fact that the average duration of treatment for men is 276 days, while for women it is 295 days. The following statistics represent the character of the brain-diseases, and the average time that each class requires for supervision in the asylums:—Simple insanity, 33,629 patients, average time 287 days; softening of brain and paralysis, 8,665, average time 250 days; senile dementia, 4,495, average time 289 days; idiots and *crétins*, 5,377, average time 298 days. The latter, therefore, appears to be the most intractable form. In all France there are only 61 public and 42 private asylums, of which nine are for men only, and 14 for women only, the remainder being for both sexes. No fewer than 34 departments have no establishment whatever for the care and treatment of the insane.

PUBLIC BATHS AT SMETHWICK.—At the meeting of the Local Board on Friday, November 9th, a letter from Mr. Tangye was read, stating that, if the Board would establish public baths, the firm of which he is a member would contribute £1,000. The offer was accepted, and a committee appointed to have plans prepared without delay.

A LECTURE ON NETTLE-RASH.

By MCCALL ANDERSON, M.D.,

Professor of Clinical Medicine in the University of Glasgow; Physician to the Western Infirmary, etc.

URTICARIA (URTICA—A NETTLE) NETTLE-RASH—URTICAIRE
—NESSELAUSSCHLAG.

THE symptoms of this disorder are very familiar to most persons, seeing that the rash is identical with that resulting from the sting of the common nettle (*urtica urens*); hence the term nettle-rash. It makes its appearance in the shape of circumscribed elevations, rarely larger than the finger-nail, which are rounded or oval, or which assume the form of segments of circles (pomphi or wheals); and, when the patches are present in their most typical form, the centre of each is pale, while the periphery is red. This eruption is accompanied by itching, burning, or stinging sensations, which are increased by scratching, and are often very distressing; but its most remarkable character—that by means of which it can be distinguished from most other eruptions, and which often enables us to say that a rash is a member of the nettle-rash group, although it does not assume the typical characters of that disease—is the wonderful rapidity with which it appears, and its transient character; for, in a few minutes, it may be fully developed over the greater portion of the body, and within an hour it may all have vanished, although sometimes two or three days elapse before it disappears; it is never followed by desquamation. Its tendency to resolution is indicated by the wheals feeling softer, by the fading of the peripheral redness, and by the subsidence of the irritation. Occasionally, vesicles or bullæ make their appearance upon the patches if the inflammatory action run high, so that the careless observer might mistake the eruption for herpes or pemphigus. And not unfrequently the rash is accompanied by œdema, especially where there is much loose cellular tissue; or œdema may take the place of the eruption—oftenest on the hands and face (*urticaria edematosa*). The extent of the rash is very various; sometimes it is partial, being limited to the hands or face, while at other times the whole surface is more or less involved. Sometimes the patches are distinct from one another, sometimes they are confluent (*U. conferta*). It presents, too, certain peculiarities, according to the region affected. "When it occurs on the face, it generally produces an œdematous swelling, especially of the eyelids and lips; the wheals, however, are less distinct than usual, and the rash for the most part assumes the character of an urticaria rubra, and consists of red lines and striae. The neck is comparatively rarely the seat of this eruption, which is, however, more commonly seen on the chest and back, where, as on the face, it often takes the form of striae, and sometimes of peculiar wavy lines. On the limbs, it is observed less frequently than on the trunk. When nettle-rash affects the neighbourhood of a joint, the skin over the articulation becomes swollen and œdematous. If the hands and feet are attacked by it, the patient often complains merely that they feel as if covered by some woollen substance..... and no particular change in the appearance of the skin of these parts is to be detected. In some cases, however, the fingers and toes become so swollen, that their movements are interfered with." (*On Diseases of the Skin*, by Ferdinand Hebra, M.D., Sydenham Society's translation, vol. i, p. 304.)

The rapidity with which nettle-rash comes out, and its evanescence, have led to much speculation as to its cause. There are some who hold that spasm of the muscular fibres of the skin leads to the development of pomphi; while Liveing is of opinion that they are the result of a spasmodic contraction of the muscular coat of the vessels. The most generally received opinion, however, is, that they are due to an acute inflammatory œdema, having its seat in the papillary layer of the corium; and the reason why the rash is so fleeting is, apparently, that the exudation is thinner and more serous than in the case of most other inflammatory affections, and that the vessels soon recover their tonicity, and absorb the exudation.

Urticaria is usually an acute affection which disappears in a few hours (*U. ephemera*), or at most within a very few days (*U. evanida*), and sometimes it is preceded and accompanied by fever (*U. febrilis*). The presence of febrile disturbance is rather a favourable feature

than otherwise, for then there is a reasonable hope that the attack is an isolated one, and will be of short duration, disappearing with the transient cause which has produced it.

It may seem strange to speak of chronic urticaria, seeing that the grand characteristic of the rash is its evanescence; but what is meant is that, although each individual rash is of short duration, the disease is kept up by constant relapses, and thus may continue even for years (*U. perstans*; *Nesselsucht*; *Urticatio*). In this form, errors of diagnosis are apt to arise, for the eruption is, in the majority of cases, nearly or completely absent when the patient presents himself for examination, and then all that we can see upon the skin is the eruption produced by the nails of the patient in scratching (elsewhere designated a pruriginoid eruption); but an inquiry into the history of the case will prevent errors, and generally we are told that the eruption comes out when the patient is undressing at night or after he gets warm in bed, or under the influence of nervous excitement. In these cases, too, we may be helped in our diagnosis by writing letters upon the skin with a pencil, which is generally followed almost immediately by a nettle-rash tracing.

There is a variety of nettle-rash to which the term *U. nodosa seu tuberosa* has been given. It is a rare affection; it appears in the shape of pretty hard nodosities about the size of a split marble, and of a reddish colour, which involve the skin and subcutaneous cellular tissue, which occur oftenest at night, and disappear in a few hours, frequently to recur. They may involve any part of the body, but the extremities and loins are specially liable to attack. In very rare cases, owing to excessive congestion of the nodosities, rupture of the capillary blood-vessels ensues, so that, after their subsidence, round ecchymoses are left, which gradually disappear. Generally, the nodosities are multiple; but, occasionally, only a single one makes its appearance. A case of this kind came under my observation some years ago. A gentleman, about fifty years of age, and otherwise apparently in good health, came to show me a swelling upon the left side of the neck, just below the ear. When I saw it, it was beginning to subside, but it was still three inches in diameter, and raised about an inch and a half above the level of the surface. He told me that he had been subject to this for ten years, the swelling coming on about once in two months, or oftener in damp weather. From its commencement to the time of its attaining its full size, no more than five minutes ever elapsed, and sometimes its growth was so rapid that its increase in size could actually be seen. As it grew larger, it became hard, and was the seat of a slight tickling sensation, and, when very large, it interfered with mastication. It always disappeared within a few hours, generally within two or three. It never developed upon any other part of the body, and no neighbouring irritation—caries of teeth, etc.—or other cause was apparent.

Only one other variety of urticaria is worthy of mention, and is, less fleeting than those previously alluded to. In it, the inflammatory exudation occurs around the cutaneous follicles, and is accompanied by the deposit of lymph, leading to the formation of large red papules. These are exceedingly itchy; and as the patient does not spare himself, the summits of many of the papules are torn off by the nails, and the blood which exudes dries up into little blackish crusts, thus somewhat resembling prurigo. Mingled with the papules are generally found some of the typical nettle-rash wheals. This is the variety of urticaria most frequently met with in young children; it is most commonly seen upon the hips and extremities, and often lasts for months. It corresponds with the *Urtica urticatus* of Willan.

Etiology.—This is one of the few diseases of the skin to which the term neurotic may, not inappropriately, be applied, seeing that the vaso-motor nerves are principally at fault. This vaso-motor disturbance may result from direct irritation of the skin, or may be reflex, arising from the irritation of distant organs and tissues; but whatever the cause may be, the first consequence is contraction, which is succeeded by dilatation and paralysis of the capillary vessels of the affected parts.

The most familiar illustration of local irritation, resulting in nettle-rash, is to be found in the sting of the common nettle (*Urtica urens* or *dioica*), the sting resulting from the irritation of the fluid in the glands on the under surface of the leaves connected with the prickly hairs, which contains sulpho-cyanogen. It is also often called forth, in those who are predisposed, by scratching the skin, or by the bite or sting of insects, such as the flea, the bug, the mosquito, and the wasp; in which case, in the centre of each wheal, the seat of puncture, in the shape of a dark point, is to be seen (*Urticaria traumatica*). Some years ago, I received a telegram asking me to visit immediately a well-known gentleman in the West

of Scotland, under the following circumstances. While picking strawberries in his garden, he put one into his mouth containing a live wasp, which stung him on the right side of the tongue near its root. In about five minutes, his tongue was so much swollen that he could scarcely move it; and within ten minutes the whole surface was covered with nettle-rash, which, commencing on the head and neck, rapidly spread over the whole body. The affected parts were of a deep red colour, were very much swollen, and intensely itchy. The first medicine at hand was citrate of magnesia, of which he had half an ounce; and about an hour afterwards he vomited, putting up with the vomited matter some strawberries and gooseberries of which he had partaken. When he was sick, the itching almost disappeared, but returned again afterwards, though not with the same severity. Three mild antibilious pills were then administered, which acted in about two hours; and within four hours after the commencement of the nettle-rash, it had entirely disappeared. The other day I saw, in consultation with Dr. Miller of Dundee, a single lady who had been troubled with nettle-rash for eighteen months, in whose case the washing of the face with warm water brings it out at once, so much so, that for half an hour thereafter she is unable to go down to breakfast on account of the disfigurement.

Amongst other internal causes which may produce nettle-rash by reflex action may be mentioned the following.

a. Irritation of the uterine nerves in connection with uterine disorders of various kinds. Hebra mentions the case of a patient who had flexion of the uterus, and in whom nettle-rash was induced fifteen times in succession as the result of the introduction of the uterine sound; and cases have been recorded in which the rash appeared in connection with each pregnancy.

b. In some persons, mental emotion is sufficient to call it forth, such as an excess of joy or grief; and, once it has appeared, it is very apt to return from the slightest causes, and even from speaking of it. Some remarkable cases of this kind have been reported by Alibert. He once saw a young woman who could not enter a drawing-room without having the whole skin covered with nettle-rash, so much so that she could not dance or enjoy any other recreation; an ecclesiastic who could not celebrate divine service because the eruption immediately came out, and caused him to scratch himself with the greatest violence; and a poor woman who for sixteen years was the victim of this complaint, and who could not speak without the whole body being covered with the rash.

c. Derangements of the digestive organs are very apt to produce it, or partaking of certain articles of food, or even food to which patients are unaccustomed (Dr. Thomson). The kind of food which produces it varies in different persons, but they soon get to know what they cannot take with impunity. Shell-fish, such as muscles, oysters, crabs, and lobsters; fruits, such as nuts and almonds; vegetables, such as onions and garlic—especially if underdone; meat, such as pork and sausages; and medicines, such as valerian, copaiba, cubebs, turpentine, and quinine, are, perhaps, most apt to induce it. In a letter which I received some time ago from a medical friend on the subject of nettle-rash, he mentions the following substances as being apt to disagree with him; "*In primis*, nuts of all kinds; haws from the hawthorn, especially if very ripe; raisins, figs, prunes, and dried fruit of all kinds, especially if containing sugar. Dates do so very rapidly; sometimes grapes, if I eat the skins. Almonds, wheat, new oats, peas (green and dried); beans of all kinds, unless cooked; most pastries if they contain a good deal of oil, and are what are called heavy or rich; infusion of senna; and common tea, if strong, and without cream and sugar. Neither coffee nor chocolate injures me, but cocoa does; and common scones and rolls, if the loose flour be left on them, especially if they be taken hot, and spread with butter. Opium and Dover's powder sometimes produce a like effect. The attack begins in this wise. One day, at a dinner-party, I thoughtlessly began to eat a few nuts, when, almost instantly, even when they were on the tongue, I began to feel a tingling sensation, with heat, and a sense of fullness in the throat, and swelling of the fauces. In a few minutes, itching and tingling began in the palms of my hands and soles of my feet, and within twenty minutes the whole body was covered with rash, as if I had been thrashed all over with nettles..... My usual remedy is brandy or whiskey; indeed, I can eat most of the above-mentioned articles if I am drinking whiskey-toddy at the time."

The following case is also worthy of being put upon record. I quote from a letter of a relative of my own. "My experience of nettle-rash is anything but recent, as it is now nearly thirty years since I discovered that I could not eat butcher's meat in any form without causing it, upon which I finally gave up the indulgence of

that taste. Since then, I have once or twice had slight attacks of nettle-rash from partaking of very strong soup, but none of those violent symptoms which the solid meat used to occasion. I first made the discovery after a long fever I had in 1830-31. I had previously suffered occasionally from nettle-rash, but not violently, nor uniformly, on eating meat. After my fever, however, it was a clear case. It was long before I found that everything in the shape of butcher's meat was inadmissible. Many trials were made with meats, and portions apparently as tender as, or more so, than fowl, as for instance, rabbits, ox or sheep's tongue, sweetbreads, etc.; but all with the same inflexible result, and that whether I knew what I was eating, or expected to suffer from it, which satisfied me and the most incredulous around me, that imagination had nothing to do with it. The symptoms did not begin for an hour or two. The first was the feeling of a lump in my stomach, perceptible even to the touch; then appeared nettle-rash on my wrists, my arms, my groins, and other tender parts of the skin; at first, in separate white blisters (as if an army of fleas and bugs had attacked me), which shortly agglomerated into large masses of white blisters. Along with this the inside of my throat and nose became swelled, my voice hoarse, and a feeling as if I had a violent stuffy cold in the head. If the attack were less severe, I used to go to bed, and was well by morning. If more violent, I used to take magnesia, which acted strongly on my bowels, causing first faintishness, and then severe purging, after which I became well. Various members of my wife's (she is a blood-relation of his own) family have been subject to nettle-rash, but not from the same cause. My mother-in-law could not eat barley-meal, nor my brother-in-law oatmeal, without suffering from it, though not, I believe, so severely as myself. My wife cannot let figs or wall-flowers touch her face without producing a rash.....If you will make it worth my while, I will come down at the Whitsuntide holidays and be exhibited. I will also eat the *Ornithorhynchus paradoxus*, if you can catch one unstuffed, and finally determine whether it be bird or beast."

This case illustrates the occasional hereditary nature of the disease—a point which has also been brought out by other writers; amongst others, by Trousseau, in his work on *Clinical Medicine*, Sydenham Society's translation, vol. ii, p. 285.

In many cases, specially in chronic urticaria, no cause can be made out. In the latter, the reason may be that the cause which originally produced the attacks has passed off, and the disease has been kept up owing, so to speak, to the skin having contracted a bad habit; or it may be the result of some peculiar idiosyncrasy, which is a convenient term to hide our ignorance.

Diagnosis.—When the eruption appears in its typical form—in the shape of wheals which are pale in the centre and red at the edges—it cannot be mistaken for any other; and when it is due to the sting of an insect, the dark point in the centre of each wheal is characteristic; but when it assumes one of the less usual forms, the lesion being erythematous or papular (*Lichen urticatus*), or tubercular (*U. nodosa*), mistakes may sometimes arise, if due care be not taken. Such errors may, however, be generally avoided by noting the presence of the four following points, which almost invariably characterise the members of the urticaria group: 1, the rapidity with which the eruption makes its appearance; 2, the itching, burning, or stinging sensation to which it gives rise; 3, its short duration, although the disease may be kept up indefinitely, owing to the occurrence of successive crops; 4, its not being followed by desquamation.

Treatment.—The first point in the management of any case of urticaria is to endeavour to find out, and, if possible, to remove, the cause or causes, the nature of which has already been sufficiently considered in a former section.

In acute cases, the eruption generally subsides within two or three days, when no treatment is adopted; but generally a sharp purge is of use, especially when, as in the majority of instances, the eating of some indigestible food, or digestive derangement, is at the root of the mischief; and, if we have reason to believe that such food is still in the stomach, as indicated by nausea, etc., an emetic of mustard, ipecacuan, or sulphate of zinc, may be administered at the outset. In all such cases, stimulating food and drink should be avoided, and the diet should be of the simplest kind.

In chronic cases, a similar line of treatment should be pursued under similar circumstances, and, where we have reason to suspect that it results from the digestion being disturbed by some particular kind of food, the nature of which varies in different persons, owing to their peculiar idiosyncrasy, we may with advantage follow the advice of Willan, who wrote, "I have desired several persons, affected with chronic urticaria, to omit first one, and then

another article of food or drink, and have thus been frequently able to trace the cause of the symptoms. This appeared to be different in different persons. In some it was malt liquor, in others, spirit, or spirit and water; in some, white wine; in others, vinegar; in some, fruit; in others, sugar; in some, fish; in others, unprepared vegetables." Like most other observers, however, he found that, in some cases, a complete alteration of the diet was not of the least avail. It would be quite out of place to refer to the means to be taken for the removal of the many other causes of this affection, as these must be treated upon general principles, and in the same way as we should do if they were independent of urticaria.

When no cause can be made out, or where the supposed cause has been removed, and the eruption continues to crop up, we must treat it empirically. We may, for example, try the effect, as Trousseau suggested, of the administration of sulphuric ether in doses of twenty to forty drops in water, or of quinine in full doses, or of arsenic, which is only exceptionally useful. But the medicines from which, perhaps, most is to be expected are atropia and bromide of potassium; the former may be administered subcutaneously at night, or night and morning, the initial dose for an adult being $\frac{1}{10}$ of a grain (e.g. 5 minims of a solution of one grain of sulphate of atropia in 500 of water); the latter in doses of ten grains dissolved in water three times a day. In either case the dose should be gradually increased, either until the disease begins to yield, or until the superintention of the usual physiological effects renders it unsafe to push the experiment further. Occasionally, good results are obtained from the continuous current of electricity for ten minutes night and morning, one sponge (the positive pole) being applied to the top, and the other to the bottom of the spine.

In obstinate cases, a complete change of air and scene and occupation is desirable, and sometimes advantage is obtained from visiting one of the alkaline spas, as Vichy, or from a course for three or four weeks of the baths of Leuk, in Switzerland.

Local treatment is generally resorted to, more with the view of alleviating the distress of the patient than in the hope that it will out short the disease. The parts, for example, may be sponged with vinegar and water, or with eau de Cologne, or with a lotion of carbolic acid.¹ An ointment containing chloroform or a mixture of chloral and camphor² may sometimes be of service, and in exceptional cases not only temporary relief, but permanent benefit may result from the use of the tarry preparations, such as a lotion composed of equal parts of tar, soft soap, and rectified spirit.

¹ *A Practical Synopsis of Cutaneous Diseases, According to the Arrangement of Dr. Willan*, by Thomas Bateman, M.D., F.R.S. Seventh edition, edited by A. T. Thomson, M.D., F.R.S. London: Longmans. 1820.

² $\frac{1}{2}$ Acidii carbolicii cryst. 3ii; glycerini (Price) 5vi; eau de Cologne 5i; aquæ distillatæ 3iv.

³ $\frac{1}{2}$ Chloralis hydratis, camphoræ aa 5ss; misce et adde glycerini (Price) 5i, unguenti simplicis ad 5i.

HOW FEVER IS SPREAD.—At the Manchester Police-court last week, Elizabeth Leigh was charged with stealing wearing apparel. Prisoner, who was a nurse, was called in to nurse a lady suffering from scarlet fever. She was so engaged for a fortnight, and during that time she stole a quantity of the underclothing which the patient had worn during her illness, and this she forthwith pawned. The magistrate described the prisoner's conduct as outrageous, and sentenced her to nine months' imprisonment.

CANTOR LECTURES.—The first course of Cantor Lectures, at the Society of Arts, was commenced last Monday, and the subject was "The Scientific Basis of Cookery," by W. Mattien Williams, F.C.S. The introductory lecture treated of modes of applying heat; radiation, conduction, and convection; roasting, grilling, baking, boiling, and stewing. The second lecture will deal with the constituents of flesh; the action of heat on albumen, gelatine, fibrin, &c.; exosmosis and endosmosis as operating in the kitchen; maceration; caseine; the cookery of cheese and its nutritive value; milk, butter, and "Bosch." In the third and concluding lecture will be considered the nutritive constituents of vegetables; the change effected by cookery on vegetable substances, &c. The lectures will be illustrated by a selected exhibition of cooking apparatus and appliances.

The Thorne Rural Sanitary Authority, Yorkshire, have received a letter from the Local Government Board, intimating that they are not prepared to assent to the continued employment of the poor-law district medical officers as medical officers of health; and trusting to receive a proposition to appoint one independent medical officer of health for the whole district. It was decided to consider the matter at the next meeting, and to give special notice to all the members, so as to ensure a good attendance and full discussion.