We select the letters for these pages from the rapid responses posted on bmj.com favouring those received within five days of publication of the article to which they refer. Letters are thus an early selection of rapid responses on a particular topic. Readers should consult the website for the full list of responses and any authors' replies, which usually arrive after our selection.

LETTERS

MARTIN KURZER

HERNIA REPAIR

Outcomes other than recurrence should be analysed

Nordin and van der Linden's attempt to link outcome after groin hernia surgery with surgeon volume, using the Swedish hernia register, is commendable.¹ But even they must admit that using recurrence as the only outcome is crude, outdated, and of little relevance nowadays. In addition, not all hernia operations are equivalent—size and difficulty vary greatly—and their study took no account of this.

Twenty years ago, the success of hernia repair was measured mainly by recurrence. Since the advent and widespread use of mesh patches, recurrence rates have fallen greatly,² and other outcomes are now of greater clinical importance.³ A "good result" is not simply a hernia that does not recur.

Long term postoperative pain and discomfort are currently the major problems after groin hernia repair, with an incidence of 10-25% one year after operation. They may be related to intraoperative nerve damage and hence associated with surgical technique and surgeon's experience.⁴ Other important outcomes include complications, time to return to normal activities, local anaesthetic and day case rates, quality of life, and patient satisfaction.⁵ Unfortunately, none of these was assessed in this study.

As pointed out in the linked editorial, recording national data has an important role in planning the delivery of services and comparing peers, and if surgery is going to progress and become more evidence based, outcome analysis must remain a priority, with relevant clinical input. The modern outcomes for hernia surgery are postoperative pain, complications, and

quality of life. It is a pity that this large review did not look at these outcomes.

Martin Kurzer surgeon, m.kurzer@mac.com

Alan E Kark surgeon, Tahir Hussain surgeon, British Hernia Centre, London NW4 4RS

Competing interests: All authors are surgeons at a private specialist hernia centre.

- 1 Nordin P, van der Linden W. Volume of procedures and risk of recurrence after repair of groin hernia: national register study. *BMJ* 2008;336:934-7. (26 April.)
- 2 EU Hernia Trialist's Collaboration. Repair of groin hernia with synthetic mesh: meta-analysis of randomized controlled trials. *Ann Surg* 2002;235:322-32.
- 3 Jenkins JT, O'Dwyer PJ. Inguinal hernias. *BMJ* 2008;336:269-72.
- 4 Kehlet H. Chronic pain after groin hernia repair. *Br J Surg* 2008;95:135-6.
- 5 Burney RE, Jones KR, Coon JW, Blewitt DK, Herm A, Peterson M. Core outcomes measures for inguinal hernia repair. *J Am Coll Surg* 1997;185:509-15.

SENTINEL LYMPH NODE BIOPSY

Let's get back to basics in managing melanoma

I find the debate about sentinel node biopsy puzzling.¹ Clinicians are getting bogged down with this question while the fundamentals of managing cutaneous melanoma are being neglected. A recent Australian study showed that doctors perform poorly in key aspects of managing this aggressive tumour.² Only a third of doctors excised cutaneous melanoma with the margins recommended by the Australian guidelines—a third used larger margins and, more worrying, a third used narrower margins. Most doctors failed to check the skin at follow-up, and they often diagnosed suspicious lesions by biopsy not local excision. Australian surgeons are slow to acquire dermoscopic skills that improve early diagnosis of melanoma. A patient with a thin melanoma is more likely to develop another cutaneous primary than metastatic disease. Yet dermoscopy and skin checks are often neglected.

If surgeons used time spent doing sentinel lymph node biopsy in routinely examining the skin at follow-up, there would be a tangible gain for our patients. In contrast, a procedure with a 10% incidence of complications,³ which does not improve five year survival,⁴ is hardly a tangible gain. Let us get back to basics. Let's offer our patients skin checks for life, ensure dermoscopy is a routine part of this examination, and excise suspicious lesions rather than biopsy them to gain histology. Most importantly, let's give our patients with invasive melanoma a minimum 10 mm margin of normal skin rather than skimp.

Anthony Dixon censor, Australasian College of Skin Cancer Medicine, Geelong, VIC 3216, Australia
anthony@skincanceronly.com

Competing interests: None declared.

- 1 Thomas JM. Sentinel lymph node biopsy in malignant melanoma. *BMJ* 2008;336:902-3. (26 April.)
- 2 Kelly JW, Henderson MA, Thursfield VJ, Slavin J, Ainslie J, Giles GG. The management of primary cutaneous melanoma in Victoria in 1996 and 2000. *Med J Aust* 2007;187:511-4.
- 3 Morton DL, Cochran AJ, Thompson JF, Elashoff R, Essner R, Glass EC, et al. Sentinel node biopsy for early-stage melanoma: accuracy and morbidity in MSLT-I, an international multicenter trial. *Ann Surg* 2005;242:302-11; discussion 311-3.
- 4 Morton DL, Thompson JF, Cochran AJ, Mozzillo N, Elashoff R, Essner R, et al. Sentinel-node biopsy or nodal observation in melanoma. *N Engl J Med* 2006;355:1307-17.

PREMATURE MENOPAUSE

Article's recommendation of HRT is highly questionable

I thought we had laid to rest the myth that hormone replacement therapy (HRT) has a beneficial effect on cardiovascular disease. The American women's health initiative study, published in 2002,¹ which was stopped short because combined HRT increased the risk of breast cancer 2.5-fold, also showed that HRT actually increased stroke and venous thrombosis by 29%. The same study showed an increase in bone mineral density and reduction in fractures. Short course HRT has been shown to confer long term benefit,² so the short course is the only sensible recommendation. It should be given only for severe menopausal side effects, because the risk of breast cancer increases with the duration of treatment.³

If we add in the higher risk of recall for further investigation after breast cancer screening and cervical screening shown by the million women study, with all the psychological distress that it causes, then it seems irresponsible for Jones et al to recommend combined hormone replacement therapy for 15 years or more.⁴

I sympathise with women who have menopausal symptoms, but find it hard to pass by advice that is so contrary to good practice.

Mark D Lewars consultant radiologist, Southend Breast Unit, Southend University Hospital NHS Trust, Essex SSO ORY
mark.lewars@southend.nhs.uk

Competing interests: None declared.

- 1 Roussouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, et al. Risks and benefits of oestrogen plus progestin in healthy post-menopausal women: principal results. *JAMA* 2002;288:321-33.

- Middleton E. The effect of short-term and low-dose hormone replacement therapy on long-term bone mineral density. *RAD Magazine* January 2008:19.
- Short Cuts. HRT associated with lasting risk of cancer. *BMJ* 2008;336:582-3.
- Jones GL, Ledger W, Mitchell C. Suspected premature menopause. *BMJ* 2008;336:833. (12 April.)

LUNG AGE

Study's conclusion about screening is unwarranted

Although Parkes et al's study is interesting,¹ the only conclusion is that for asymptomatic smokers who undergo spirometry, communication about lung age is a more effective motivator for smoking cessation than providing patients with uninterpreted spirometry measurements. The study does not establish the independent motivational effectiveness of doing spirometry screening versus not doing it, which would require a randomised trial in which the control arm did not receive spirometry. Only two randomised controlled trials of this type have been conducted and both had negative results. For this and other reasons, the US Preventive Services Task Force recommends that adults (regardless of smoking status) should not be screened for chronic obstructive pulmonary disease using spirometry.²

Kenneth W Lin medical officer, Agency for Healthcare Research and Quality, Rockville, MD 20850, USA
Kenneth.Lin@ahrq.hhs.gov

Competing interests: None declared.

- Parkes G, Greenhalgh T, Griffin M, Dent R. Effect on smoking quit rate of telling patients their lung age: the Step2quit randomised controlled trial. *BMJ* 2008;336:598-600. (15 March.)
- US Preventive Services Task Force. Screening for chronic obstructive pulmonary disease using spirometry: US preventive services task force recommendation statement. *Ann Intern Med* 2008;148:529-34.

SELECTING MEDICAL STUDENTS

Beware of self selection and self fulfilling prophecy

Because the *BMJ* is mostly read by clinicians, we should be aware of the risk of self selection.¹ Just as companies can develop an internal culture, whereby only similar personality types are recruited, so the same happens with doctors. We select only those with high academic results, we train them to be problem focused, and, not surprisingly, they tend to select similar people. There is a danger in simply selecting what you already have, as it may not be what you need.

Historically, doctors needed good memories—they will probably continue to do so, but because medical knowledge has outstripped our ability to remember it all, we may do better with someone who is adept at using computer based tools.

We select people who relish solving problems and train them in high stress situations, so we

select those who get a buzz out of "winging it." This may explain why medical practice is slow to adopt care pathways and protocols as "it takes the fun out of medicine."

Teams often fail because they miss the "finisher" person—we select clinicians with the "shaper" profile, who want to see their solution put into action.² Some of the most dysfunctional teams consist of many shapers, all vying to get their idea through.² Because we need to move to a team based approach to medicine and more flexible roles, perhaps we should review how we select healthcare staff, including whether they need to be "professionals" and whether they should be managers rather than divas. I admit to somewhat caricaturing clinicians here, but, having run focus groups across a range of stakeholder groups, I have found that clinicians are different in the way they approach matters.

Peter D Singleton principal research fellow, University College, London N19 5LW
peter.singleton@chi-group.com

Competing interests: None declared.

- Brown CA, Lilford RJ. Selecting medical students. *BMJ* 2008;336:786. (12 April.)
- Belbin RM. *Management teams: why they succeed or fail*. 2nd ed. Oxford: Butterworth Heinemann, 2004.

WHY I LOVE A FREE LUNCH

Free lunches are not the answer

I share Dr Patel's concerns about doctors' continuing education and the quality of hospital food.¹ However, I disagree that "free" lunches are the best solution to either of these problems.

No clinician can be expected to follow the progress of experimental drugs in hundreds of journals. That is why we have independent sources of information about drugs—such as the *Drug and Therapeutics Bulletin*, *NPS Radar*, and *Prescrire International*—which aim to distil a complex evidence base into accessible guidance for busy clinicians. *Prescrire International* published a review of ivabradine last year, which deemed it "best avoided."² According to the abstract, concerns include a lack of evidence of superiority to standard antianginal drugs and evidence of increased adverse cardiac events in patients taking ivabradine compared with those taking atenolol or amlodipine.² Did Dr Patel hear these messages from his smiling drug representative? I found this information via a PubMed search in about the time it would have taken Dr Patel to eat his lunch.

Contact with drug representatives is positively associated with prescribing new drugs, prescribing costs, and non-rational prescribing.³ Industry paid meals are associated with formulary addition requests.³ New drugs are not necessarily safe; how many patients may have been harmed by free lunches promoting rofecoxib? And "free" lunches are not really free—

they are paid for by drug sales. It would be wise to improve investment in doctors' prescribing education, but let's not deceive ourselves that free lunches are a substitute for this.

Brett D Montgomery clinical senior lecturer, Discipline of General Practice, School of Primary, Aboriginal and Rural Health Care (SPARHC) University of Western Australia, Crawley, WA 6009, Australia
brett@multiline.com.au

Competing interests: BDM is a member of Healthy Skepticism, an organisation that aims to reduce harm from inappropriate drug promotion.

- Patel K. Why I love a free lunch. *BMJ* 2008;336:962. (26 April.)
- Anonymous. Ivabradine: new drug. Best avoided in stable angina. *Prescrire Int* 2007;16:53-6.
- Wazana A. Physicians and the pharmaceutical industry: is a gift ever just a gift? *JAMA* 2000;283:373-80.

You must be joking

How are doctors supposed to get new information? Perhaps by reading during your lunch break, rather than listening half heartedly while munching on your Marks and Spencer wrap.¹ Or even, maybe, a journal club? Drug sponsored lunches are socially uncomfortable, and consume time that could be used to chat with your colleagues (team building); to learn about something that might change your practice or might be evidence based, not just speculative; to catch up on the news—a mildly interesting fact or two that probably has no direct bearing on your life, personally or professionally.

"The NHS is unwilling to devote serious capital to fund most educational activities."¹ The free lunch does not contribute to education; indeed, I can afford my own lunch and would rather the money be used for patients. What helps education is having protected time. Not money. Or free food.

Adrian H Sie specialist registrar in paediatrics, Crosshouse Hospital, Kilmarnock KA2 0BE
a_sie@doctors.org.uk

Competing interests: None declared.

- Patel K. Why I love a free lunch. *BMJ* 2008;336:962. (26 April.)

REDUCE, REUSE, RECYCLE

Cool for cats

In striving to reduce the use of plastic to wrap the *BMJ*,¹ why not post it rolled up in a brown paper wrapper as was done a few decades ago? After his retirement my father tended to look at the *BMJ* just "to see who's died," but some copies remained unopened and were kept by the back door. He disliked marauding cats in his garden and found a rolled up *BMJ* was an excellent deterrent up to 20 yards.

Charles Oxley retired GP, New Buckenham, Norfolk NR16 2AG
charlesoxley@yahoo.co.uk

Competing interests: CO gets cats in the garden.

- Groves T. Reduce, reuse, recycle. Editor's choice. *BMJ* 2008;336. (26 April.)