resected portion of the diaphysis. Approximately at right angles to this outline the line A B is seen in Fig. 2 and indistinctly in Fig. 3. The inner portion of the line A' B'

Fig. 2.—Diagram showing the position of the parts when the operation is complete.

stands out in both the model and the photographs. In the skiagrams the continuation of this line is seen as an irregular shadow through the shaft of the femur. In the original of Fig. 3 the new internal lower end of the diaphysis appears exactly as if it were received into a depression of the resected portion of the diaphysis.

Fig. 3.—Skiagram of the right knee nineteen days after operation. (Taken with the plate behind the limb.)

The study of the evidence presented by the skiagrams and illustrated diagrammatically in Fig. 2, shows us that the alterations brought about by the operation are somewhat as follows:

1. The inferior epiphysis of the femur retains its relation to the resected portion of the diaphysis, and the two together may be described as the lower fragment in a complete fracture.

2. The lower fragment is dislocated inwards, causing the diaphysis to project outwards beyond the external condyle. (Vide Fig. 3.)

3. The lower fragment is also rotated upwards and inwards round an antero-posterior axis passing through the epiphysal line in the neighbourhood of the intercondylic notch. This displacement is more clearly understood if we consider the direction of the forces applied in straightening the limb. The above rotation or tilting up of the lower fragment has the following consequences: (a) The separation of the epiphysis from the diaphysis externally; (b) the impaction of the compact tissue of the lower end of the upper fragment into the cancellous tissue of the resected portion of the diaphysis in the lower fragment, and the approximation of the line A B to the epiphysal line; (c) the assumption of a more vertical position of the internal portion of the epiphysal line.

4. The relations of the tibia to the condyles of the femur are not materially altered.

5. The relation of the external to the internal condyle is unaltered. The latter is not fractured through, although the osteotome penetrates the inferior epiphysis of the femur near the intercondylic notch.

In conclusion, these skiagrams appear to show the importance, in rectifying genu valgum, of being able to produce a displacement of the ununited epiphysis, and possibly also in treating other similar deformities.

THE INFLUENCE OF ACID ON GUINEA WORM LARVAE ENCYSTED IN CYCLOPS.

By ROBERT T. LEIPER, M.B., F.Z.S.,

HELMINTHOLOGIST AT THE LONDON SCHOOL OF TROPICAL MEDICINE.

THE practical study of dracontiasis, one of the oldest known and, in some regions, the most prevalent of tropical diseases, has been singularly neglected.

Although not a fatal disease or one which attracts much sympathy, its incidence nevertheless is commonly the precursor of much suffering and physical incapacity. Modern methods of treatment are at once so unsatisfactory and necessarily prolonged and so inaccessible to the vast majority of sufferers that it can only be from the application of prophylactic measures that the diminution or eradication of the disease is to be expected.

In dracontiasis a knowledge of the life-history of the *Filaria medinensis*, and particularly of the stage and manner of its entrance into the human body, must form the essential basis on which such measures may be framed, yet Manson appears to be the only observer who has made any important contribution to Fedschenko's original discovery of cyclops as the intermediate host thirty-six years ago.

Based on many theoretical considerations, but especially influenced by the recent discovery that the young stages of certain worms, such as *Ankylostomum duodenale* and *Strongyloides intestinalis*, make their way into the body of their host through the follicles of the skin, modern scientific opinion has become more and more prepared to postulate a like mode of entry in dracontiasis, while at least one authority has suggested that a second intermediate host might even be necessary. In view of the fact that a discussion in which these theories were in turn maintained took place on this subject at the last annual meeting of the British Medical Association, and was recently fully reported in the BRITISH MEDICAL JOURNAL, thus attracting considerable attention, I am induced to publish somewhat in advance an observation made by me while studying this disease on the Gold Coast, which seems to throw suggestive light on the question.

The embryos of *Filaria medinensis* during the first fortnight of their stay in the body-cavity of cyclops—that is, during their entire metamorphosis—continue to exhibit active vermicular movements. They also not infrequently change in position. During the third and fourth week, however, these changes gradually cease, and ultimately the larva lies almost wholly quiescent. If a cyclops containing such a larva be placed on a slide in a very small drop of water, and if, while it is being watched under the low power of the microscope, there be added some 0.2 per cent. hydrochloric acid, representing the

acidity of the gastric juice, a remarkable change will be seen to take place. The cyclops is gradually killed, showing little motion beyond a few tremors of the antennae and rami. On the other hand, as the acid penetrates, the apparently quiescent filaria regains its former activity. At first slowly, but with every moment with increasing strength and rapidity, it coils, stretches, and recoils, until at last, having divested itself of what probably is the remains of a very delicate cuticle, it bursts into the general body cavity of the now moribund cyclops, and in a very short time completely destroys the organized contents by its increasingly violent motions. The larva is soon seen to be testing with obvious purpose every joint of the cuticular integument. At one moment it is attempting to advance along the entire length of a narrow antenna, at the next it may be found attacking one or other of the prolongations of the bifurcated tail. Retreating again, it will endeavour to ascertain the patency of the anus, the genital openings, or the mouth. By one or other of these apertures, or, in some cases, it may be by a rupture of the cuticle, the young worm eventually reaches the exterior, and there swims about with great rapidity like a minute snake. One specimen, watched for three hours, continued with brief rests, in which it coiled and uncoiled, to manifest the same marvellous vitality.

In contrast, other cyclops containing embryos of the same date of infection, and in all respects similar to those used in the above experiments, were kept alive for a further period of two weeks—that is, until 6 weeks old—but their contained larvae did not exhibit change either in complexity or in gross increase in size. Some of these cyclops died in the meantime, but the guinea-worm embryos did not appear to escape; they were also found dead.

This experiment seems to indicate—in so far as an artificial experiment may—that the *Filaria medinensis* is introduced while still in a quiescent state in the body of cyclops into the stomach of the human host, whence, on being set free by the influence of the gastric juice, it proceeds to its further development within the body.

MEMORANDA:

MEDICAL, SURGICAL, OBSTETRICAL.

A SERIES OF CASES OF ICTERUS NEONATORUM IN A FAMILY.

AFTER an extensive perusal of the literature on the subject of icterus neonatorum I have come to the conclusion that the following cases are worthy of publication. The series appears to be unique as regards the number of deaths, the rapidity with which death ensued, and from the fact that there were intervening cases of recovery. Amongst the hitherto reported instances the nearest in correspondence with these is the example recorded in Ashby and Wright's *Diseases of Infancy and Childhood*. In that case the first two infants remained alive and well and were not jaundiced, but the third, fourth, fifth and sixth infants died from the ninth to the eleventh day.

Mrs. L., a woman of the working class, suffered from small-pox in early life, but has otherwise always been a well-nourished and healthy woman. She was married at 17, and her husband, apart from occasional rheumatism, has manifested no evidence of constitutional disease. The children were all born at full term between the mother's eighteenth and thirty-fourth year of age.

The first, a boy, did not suffer from jaundice, but died at 5 months from bronchitis.

The second, a girl, suffered from jaundice, but recovered.

The third, fourth, and fifth suffered from jaundice, and died. These cases occurred before I knew the family.

The sixth labour presented no unusual feature, but by the second day the infant—a boy—began to show signs of jaundice, which gradually deepened into almost a copper colour, and death followed a comatose condition about the eighth day.

The seventh, a girl, and the eighth, a boy, also suffered from intense jaundice, but recovered. These cases were treated from birth with a small dose of calomel, followed by large and repeated doses of castor oil, together with a soda and rhubarb mixture. The success which apparently

followed my treatment in these two cases led me to anticipate a favourable termination to the ninth pregnancy, but in this I was disappointed, as the infant, a very fine well-nourished boy, died from jaundice on the fourth day. In this case I was permitted to make a *post-mortem* examination, but, apart from the bile-stained condition of all the organs, I could discover no morbid condition. The gall bladder contained bile, but was not distended, and the bile duct was not occluded.

The tenth, a girl, I did not see until about six hours after birth, but it was then markedly jaundiced, and in spite of the treatment, which had apparently been successful in the seventh and eighth cases, it died early on the fourth day. The second, seventh, and eighth children, although all suffered from jaundice for a few days after birth, are now alive and very fine children. They show no evidence of specific taint. Any suspicion of syphilis being the cause of the trouble was negatived by the *post-mortem* appearances, and also by the fact that the disease, whatever its pathology, does not show any diminution of intensity after the lapse of fifteen years, but rather tends to a more rapidly fatal issue. As the mother is still far from the end of the child-bearing period, I shall be glad to receive any hints or suggestions as to future treatment of mother or infant should pregnancy again occur.

JAS. BUSFIELD, M.R.C.S., L.R.C.P., F.I.C.
Clapham Park, S.W.

IMPACTION OF A HATPIN IN THE MALE URETHRA.

I WAS called into the country recently to see in consultation a young man, aged 16. A few hours previously, while pushing a blackheaded hatpin head first down his urethra, he had lost his hold on the point of the pin, and it slipped out of sight. At the time of my examination, the head could be felt in the perineum 5 in. from the external meatus; the point had perforated the urethra and lay in the peri-urethral tissues, $1\frac{1}{2}$ in. from the meatus. I removed the pin in the following way: The patient having been anaesthetized, pressure was made in the perineum, posterior to the head of the pin, so as to prevent it passing further up the urethra, at the same time causing the point to perforate the skin of the penis on its under surface. The shaft of the pin was then withdrawn through the skin puncture until the head prevented further progress. The point was then depressed towards the perineum, and the pin, measuring $3\frac{1}{2}$ in., removed by pushing it head first through the external meatus. Recovery was uneventful.

This little manoeuvre, although a recognized procedure, is hardly so well known as it deserves to be, since it is the only method which can be relied on to save a patient who has the point of a pin embedded in the periurethral tissues from the risk of damage to his urethra by the use of forceps or from the performance of an external urethrotomy.

C. HAMILTON WHITEFORD, M.R.C.S., L.R.C.P.
Plymouth.

THE TREATMENT OF A COMMON COLD.

IN the case of an ordinary head cold the quickest result, in my experience, is obtained by giving 30 minims of sp. ether. nit. and 30 drops of sp. ammon. aromat. in 1 oz. of water, repeating the dose in two and then every four hours. Three or four doses are generally sufficient to put a stop to the discharge. Should the discharge happen to be thick when first seen, then a snuff composed of 1 gr. of cocaine, 2 gr. of menthol, and 100 gr. of boric acid quickly effects a rapid cure. When the cold has run down into the trachea, as shown by a tickling of the throat whenever a long breath is taken, then a mixture of liq. ammon. acet. $\mathfrak{z}$ ij, sp. ether. nit. $\mathfrak{m}$ x in 1 oz. of water, every four hours, rapidly gives the required relief.

Bexhill-on-Sea.

F. P. ATKINSON.

MEDICAL GRADUATES' COLLEGE AND POLYCLINIC.—Sir Felix Semon, K.C.V.O., will deliver the opening lecture of the winter session of the Medical Graduates' College and Polyclinic, at 22, Chenies Street, W.C., on Tuesday, January 9th, at 5.15 p.m. The subject will be Some Unusual Manifestations of Syphilis in the Upper Air Passages. All members of the medical profession interested in this subject are cordially invited to be present.