explanation of the arterial spasm which up to the present time has been offered has been, to use Sir R. Douglas Powell's words, that of "a pure neurosis of the cardiovascular system."

Owing to the varying degree of intensity of the symptoms I would suggest that in classifying the cases the simplest distinction might be found in the terms "angina pectoris major" and "angina pectoris minor," the former being confined to those cases in which there was believed to be permanent anatomical change in the heart or its vessels.

REFERENCES.

- ¹ G. A. Gibson, *Nervous Affections of the Heart*, 1904. ² Thomas Oliver, *Lancet*, September 16th, 1905, p. 809. ³ James Mackenzie, *BRITISH MEDICAL JOURNAL*, October 7th, 1905. ⁴ Pawlow, *The Work of the Digestive Glands*. ⁵ Horter, *Lectures on Chemical Pathology*. ⁶ Chittenden, *Physiological Economy in Nutrition*. ⁷ W. Heberden, *Med. Trans. Coll. of Phys., Lond.*, vol. ii, p. 69, 1772; do., vol. iii, p. 1, 1785. ⁸ C. W. Parry, *An Inquiry into the Symptoms and Causes of the Syncope Anginosa, commonly called Angina Pectoris*, 1799. ⁹ Allan Burns, *Diseases of the Heart*, 1809, p. 138. ¹⁰ Jurine, *Mémoire sur l'Angine de Poitrine*, 1815. ¹¹ Laënnec, *Diseases of the Chest* (first edition, 1819), Forbes's translation, fourth edition. ¹² Forbes, *Cycl. of Pract. Med.*, vol. i, 1832. ¹³ Bertin, *Diseases of the Heart*, etc., 1833. ¹⁴ Hope, *A Treatise on Diseases of the Heart and Great Vessels*, 1839. ¹⁵ Latham, *Diseases of the Heart*, 1846. ¹⁶ Walshe, *Diseases of the Heart* (first edition 1851), third edition, 1862. ¹⁷ Stokes, *Diseases of Heart and Aorta*, 1854. ¹⁸ Trousseau's *Lectures on Clin. Med.*, New Sydenham Soc., vol. i, p. 592, et seq. ¹⁹ Nothnagel, *Deut. Arch. f. klin. Med.*, Bd. 2, s. 173; Bd. iii, s. 309. Leipzig, 1867. ²⁰ Sir B. C. Brodie, *Lectures on Pathology and Surgery*, 1846, p. 360. ²¹ Boullay, *Arch. Gén. de Méd.*, vol. xxvii, p. 425, 1831. ²² Charcot, *Gaz. Méd. de Paris*, 1859, *Progrès Méd.*, 1887, p. 99. ²³ Lauder Brunton, *Lancet*, vol. ii, p. 97, 1867, *Trans. Clin. Soc. Lond.*, vol. iii, p. 191, 1870. ²⁴ W. T. Gairdner, *Reynolds's Sys. of Med.*, Lond., 1877, vol. iv, p. 535. ²⁵ Sansom, *Diagnosis of Dis. of Heart and Aorta*, 1892. ²⁶ Osler, *Lectures on Angina Pectoris and Allied Conditions*, 1897. ²⁷ Sir R. Douglas Powell, *Albutt's System of Med.*, vol. vi. ²⁸ Byrom Bramwell, *Diseases of the Heart*, 1884. ²⁹ G. A. Gibson, *Diseases of the Heart and Aorta*, 1898. ³⁰ George Oliver, *Blood Pressure and Tissue Lymph Circulation*, 1903. ³¹ ³² W. Russell, *Lancet*, June 1st, 1901; *Ency. Méd.*, vol. xiii, Vessels; *BRITISH MEDICAL JOURNAL*, June 4th, 1904. ³³ Prof. Pal, *Gefässkranken*, Leipzig, 1905. ³⁴ Rosenbach, *Die Krankheiten des Herzens*, 1896. ³⁵ H. Huchard, *Maladies du Cœur*, Tome 2, 1899.

FOOD FEVER IN CHILDREN.

By EUSTACE SMITH, M.D., F.R.C.P.,

SENIOR PHYSICIAN TO THE EAST LONDON HOSPITAL FOR CHILDREN.

I PROPOSE to describe, under the name of "food fever," a class of cases with which I have been familiar for a number of years. The disorder consists of an attack of fever which comes on suddenly, is accompanied by signs, more or less pronounced, of digestive disturbance, lasts in its acute form for several days, and may linger on in a modified degree for some weeks. If these attacks occur frequently—once a month or so, as they sometimes do—their effect upon the nutrition of the patient is highly injurious.

Dr. H. Davy, in the *Lancet* of September 24th, 1904, published an interesting paper on this subject, which he called "Fever in children caused by the indigestion of certain kinds of carbo-hydrate foods." Dr. Davy instanced the case of his own son, who suffered repeatedly from this form of pyrexia. The attacks came on quite suddenly and without discoverable cause, and by their frequency interfered greatly with the boy's nutrition, brought his education to a standstill, and kept his parents in a state of continual perplexity and alarm. I have already described these cases in my large work on *Disease in Children*, under the name of "acute recurring gastric catarrh." It is not, however, every case of recurring gastric catarrh which presents these symptoms, but those which do exhibit them are so characteristic, and to many practitioners so puzzling, that I have thought it might be convenient to group them together under a distinctive name.

SYMPTOMS.

The subjects of the complaint are usually neurotic children of either sex between the ages of 3 or 4 and 10 or 12 years, although I have seen cases in children still younger. The attack begins with headache, accompanied sometimes by vomiting, sometimes by looseness of the bowels, but always by signs of some kind of digestive upset. The child lies about with a dull, pasty face, a dirty tongue; he does not care to play, and has a temperature of from 101° to 105°. If the abdomen is examined, the stomach resonance is found to reach higher than normal—often nearly as high as the left nipple—and the edge of the liver can be felt one or two fingerbreadths below the costal margin. The water is very high-coloured, or thick, and very acid.

If the case is left alone, or not treated properly, the temperature remains for some days continuously higher than the normal level, and at some period of the twenty-four hours shoots up still higher. It generally rises a little after food, and is almost always 102° or more in the evening. At the end of a few days it subsides, either suddenly after a profuse sweat or gradually, but is often moderately raised (99° to 101°) for a week, or even longer.

If, when the fever begins, the child is given a good dose of calomel or blue-pill, the temperature generally falls directly the medicine has acted; but instead of remaining low, as it would in an ordinary case of upset, it rises again in the course of a few hours, and in the evening generally reaches 100°, or a little higher. For several days afterwards the temperature may remain uncertain—not continuously higher than normal, but rising above a healthy level at some period of the 24 hours, in the evening or after a meal. The children, as a rule, make no complaint of bodily discomfort, but are fretful and irritable in the day, and at night lie awake or sleep badly, grunting and groaning and tossing from side to side. Often they do not care about their food, and sometimes it is most difficult to make them eat anything at all. The younger children, when forced to take food into their mouths, roll it about from one cheek to another, as if they could not bring themselves to swallow it.

Children who have these attacks only at rare intervals soon recover their health after the temperature has fallen to its normal level, and seem in no way the worse for their indisposition. If, however, as sometimes happens, the attacks are repeated every five or six weeks, nutrition soon begins to suffer. Such children are pale and thin, have little or no appetite, and even if lively and active, have habitually a languid circulation and cold feet. These signs of malnutrition are generally accepted by the parents as the necessary outcome of a supposed delicacy of constitution. It is the frequent recurrence of fever which causes so much anxiety, especially as the attacks come on quite suddenly and without obvious cause. A child seems in his usual health, is perhaps going to school regularly and joining briskly in all the school games. One morning he leaves his home apparently as well as usual, but returns in the evening flushed in the face and complaining of headache. His temperature is taken and found to be 102° or higher. The complaint then follows its usual course. Some of the attacks, however, have special symptoms. A highly neurotic child may have a fit of convulsions, which may be repeated again and again. In a certain proportion of the cases a prominent symptom is vomiting, which may persist for several days, and I am strongly of opinion that many of the cases described as "periodic" or "cyclical" vomiting are of this nature. The vomiting is sometimes accompanied by pain in the epigastrium or umbilical region.

A little girl of 6 was subject to sudden attacks of fever, with vomiting and pain in the pit of the stomach. The febrile condition always lasted five days, during which time she looked yellow without being actually jaundiced, and her bowels were very confined. The child, when I saw her, was supposed to be well, but she looked very pale and thin and her feet were bitterly cold. She was brought to me that I might suggest some means of preventing the attacks.

The pain is sometimes the most distressing symptom.

A child of 8 was brought to me because she had suffered for more than a year with attacks of acute abdominal pain and high fever which came on every three or four weeks. The child, after a restless night, would complain in the course of the morning of severe abdominal pain. Her temperature would be 103° or 104°. The pain always went on continuously for three days, the child sitting or lying with her knees drawn up and crying with the pain. Sometimes she was sick. At the end of the three days the pain and pyrexia would subside, leaving the patient much weakened, with a poor appetite and dislike to any exertion. I put the child upon a strict diet and enjoined special precautions against chills. I heard two years afterwards that the attacks had ceased from the time the treatment was begun.

In other cases there are signs of intestinal catarrh, generally of the descending colon, and the child passes loose stools containing mucus and blood.

Not long ago I saw a boy 11 years old, who was brought with the history of many attacks of this kind during the last four years. In these attacks the temperature would rise as high as 105° or 106° for the first few hours, then on the second day it would fall, but remain higher than normal in the

evening (99.5° to 101°) for a week or more, or as long as the intestinal disturbance persisted. But it was not only during the attacks of looseness that the temperature was raised. Often the boy would have an afternoon or evening temperature of 100° to 101.5° for a week at a time; but whenever this happened it was noticed that the stools were unhealthy and offensive, although not necessarily increased in number. Just before I saw the boy he had been at the seaside for five weeks. All through his stay not a day passed without the temperature rising above the normal level at some period of the twenty-four hours; generally it was 99.5° or 100°, but once it reached 105°. During this time the boy, although gaining flesh slowly, was pale and, as they said, "weary looking"; he complained at times of abdominal pains and occasionally of nausea. His bowels only acted after an enema and the stool passed was offensive and fermenting. His feet were always cold. The lad's diet all this time was strictly confined to Benger's food and milk (by medical advice), but during the last week of his stay he had had a little chicken in addition.

The special feature of these feverish attacks is that the rise of temperature is always accompanied by some signs of digestive upset, although not always by signs which make much impression upon the minds of the parents. Often they tell you nothing but that the child is feverish in the attacks and lies about, and it is only in answer to questions that you learn that the patient has no appetite, looks livery, and has unhealthy stools. The length of the attack is usually three days to a week; but it may last longer, and sometimes, as in the case just referred to, the temperature may remain uncertain for a month or more, often rising in the evening to 102° or 103°, although the patient's condition is fairly good and he does not seem to be actually ailing.

I always find these cases to be exceedingly puzzling to the practitioner. He usually suspects tuberculosis, and may even commit himself to that diagnosis. Certainly if the attacks occur frequently—I mean every few weeks—and the digestion in the interval remains poor, so that the patient if he do not lose flesh keeps pale and thin, with a bad appetite and little care to exert himself, the general state, coupled with the frequent, often moderate pyrexia, is suggestive of some such explanation of the symptoms. The diagnosis depends, of course, upon the history of the case—its irregular course, and intervals of comparative health, and the presence of undoubted signs, however slight, of gastro-intestinal disturbance.

PATHOLOGY.

As to the cause of the fever, I believe the cases are primarily attacks of acute gastric catarrh. They can often be traced to a definite chill; and the patients have always cold feet, and consequently their resisting power is so feeble that they are prepared to suffer from comparatively slight exposure.

But when the liver has been unloaded by a good mercurial purge and the temperature has fallen, I think the moderate rise which follows is due not to the catarrh, but to reabsorption from the bowel of injurious products of decomposition. "Auto-intoxication" is the word at present in fashionable use, and it probably expresses what is going on in these cases. The appearance of the stools in the majority of the children so affected shows fermentation, and it is a natural inference that some of the products of this fermentation must be absorbed into the blood. Moreover, the results of treatment suggest that this explanation is correct, for when the child is put upon a strict diet the temperature very quickly becomes normal.

Dr. Francis Hare, in his interesting work on *The Food Factor in Disease*, has proposed another explanation of the derangement. According to this view the phenomena which constitute a "bilious attack" are the direct consequence of over-formation of glycogen in the liver. The hepatic cells are swollen by the excess of glycogen they contain, and the whole organ becomes enlarged. The capillaries of the portal vein which ramify between the lobules are pressed upon, the flow of blood in them is impeded, and mechanical congestion is induced in the mucous membrane of the alimentary canal. The congested mucous membrane swells and blocks up the openings of the common bile duct and pancreatic duct, shutting off the secretion of those organs. Probably, also, as a consequence of the catarrh, the gastric and other digestive juices are checked or arrested, digestion and absorption cease, and we find anorexia, nausea and vomiting, and perhaps diarrhoea. Under these circumstances glycogen-

forming material can no longer enter the portal vein, glycogen ceases to be produced, and that already deposited in the cells is gradually consumed by the uninterrupted combustion in the tissues. The functions of the organs involved are then gradually restored, and the attack passes off. Dr. Hare, as I understand him, not only excludes chill from any share in the production of the disturbance of function, but goes so far as to declare that external cold actually tends to retard glycogenic distension of the cells, and should therefore be a protection against the attacks. In reply to this, I can only repeat that the actual conditions present in these cases seem hardly to favour the theory, for the feet and legs of the subjects are always bitterly cold, the resisting power of the system against chills is reduced to its lowest point, and it is only when the circulation has been improved and the body guarded by a sufficiency of warm woollen under-clothing, that the occurrence of the periodical attacks can be put a stop to.

TREATMENT.

To prevent and cut short the attacks, a careful diet is of the utmost importance, but it is not only excess of the carbo-hydrates which has to be considered. We have also to forbid all articles of food which are capable of undergoing an unwholesome fermentation in the bowels, and thus providing septic matter for absorption into the blood. Starches and sweets must be given with caution, and a combination of starch with milk, such as we find in the ordinary milk pudding of the nursery, rice, sago, tapioca, etc., is to be strictly forbidden. Milk itself, except in very limited quantity, is harmful, and a milk diet, which consists of milk in conjunction with various kinds of farinaceous foods, is actively injurious. I believe that one reason why the temperature is in many cases so slow to return to a healthy level is that the children are generally put upon a milk diet under a mistaken notion that it is the food the least likely to disagree. But surely it is an error to suppose that a fermenting process can be brought to an end by giving a fermentable food; and in all cases where the temperature continues uncertain day after day, rising above normal in the evening or after a meal, it is wise to exclude milk altogether from the diet. Acids, again, such as baked apples, grapes, oranges, and lemonade, must be condemned, for acid tends to make acid, and promote fermentation in all the fermentable things it mixes with.

The proper diet for these children consists of mutton, poultry, white fish, well-boiled green vegetables, and eggs. They may have plenty of butter with stale bread, toast, and rusks. The salted foods are good, such as bacon and, if they are old enough, ham and tongue; bloaters also and sardines, and the pastes made of anchovy, bloater, and shrimp may be allowed. These things, which are usually unknown in the nursery, are really perfectly harmless, and make good substitutes for marmalade and jam.

My experience is that if the case be treated in this way the temperature very quickly becomes normal, but it is wise to caution the parents that the dieting must be thorough. The object of it is to put a stop to the fermenting process; but a very small quantity of fermentable material will keep this process up. And that, I think, is the reason why in many cases the temperature is slow to fall, because milk is still allowed to be taken.

But while it is of the utmost importance to supply the child with a diet which is unfermentable and wholesome, it is of almost equal moment to take steps to prevent any fresh attack of catarrh which would keep up the derangement of the digestive organs. The child should go out as much as possible, but it must be our care to see that he does so without risk to his health. The chief points to attend to are the sufficient clothing of his body and the warmth of his feet and legs. Children with bare legs cannot be cured until their legs are properly covered with long woollen stockings, indoors as well as outdoors; and this applies to summer as well as winter, for it is in summer that the changes of temperature are so rapid and unexpected. But there are many ways in which a child may catch cold. In every individual case it is wise to go into the ordinary nursery routine (which may differ, of course, in different families) and find out if possible what is being done that is wrong. I may add that in doing this it is necessary to remember that children who have suffered from many attacks of this nature have become

very susceptible to slight chills, so that precautions which would be successful enough with an ordinary child are not effectual with them.

ON DRUG IDIOSYNCRASIES IN RELATION TO OFFICIAL DOSAGE.*

By C. O. HAWTHORNE, M.D., M.R.C.P.,

EXAMINER IN MEDICINE AND CLINICAL MEDICINE IN THE UNIVERSITY OF ABERDEEN; PHYSICIAN TO THE CENTRAL LONDON OPHTHALMIC HOSPITAL; ASSISTANT PHYSICIAN TO THE NORTH-WEST LONDON AND ROYAL WATERLOO HOSPITALS.

THE personal factor provided by the individual patient forms an element in medical practice the range and command of which can hardly be overstated. It weighs largely in the choice of a physician and sometimes with equal unreason determines his dismissal; it modifies the expression of the processes of disease and thus complicates the art of diagnosis; and last, but by no means least, it influences the actions of medicines and therefore demands an attempted estimate in the construction of a scheme of treatment. Hence it must ever remain true that the prescription of remedies in the treatment of disease can never become a mere mechanical art. For even when remedies have been accurately accommodated to definite and exact standards patients will continue to differ, and this difference will express itself in a more or less conspicuous fashion in the shape of a varying response to the influence of drugs.

The existence of idiosyncrasies in relation to drugs is a doctrine which, it is true, receives some measure of recognition in the discipline of medical education. But it may be questioned whether it might not advisably be presented with greater emphasis. For the most part it is illustrated by the announcement that occasional individuals are found who are excessively susceptible to, or extraordinarily tolerant of, some few of the more potent agents among the official materia medica. Thus drug idiosyncrasies are presented as extraordinary or exceptional occurrences which appear as "sports" amidst the generally stable and constant phenomena of the therapeutic world. So far from being merely extreme expressions of a general law, these illustrations seem to be the negation of all law. They are apt to appear, like so many big gooseberries, to enliven the dull season of a course of lectures on materia medica, rather than to enforce a practical therapeutic lesson. So considered, it must indeed be admitted that instances of unusual sensitiveness to such remedies as opium and arsenic have this value, namely, that they teach the student and junior practitioner to be careful in the use of these remedies. But it is submitted that this is only a limited application of the significance of such facts.

The suggestion is here offered that these facts are merely extreme and conspicuous examples of the general truth that every individual patient has a degree of reaction, special and personal to himself, in relation to each member of the materia medica. The majority of individuals may, it is true, closely approximate in this respect to a common standard, but between the extremes of excessive susceptibility and excessive tolerance all gradations are possible, and successful therapeutics largely consists in the determination of the exact position of the individual patient in relation to the individual remedy. The probable truth of such a statement may well be expected *a priori*. If one individual is found to react powerfully to a minute dose of atropine, for example, whilst another receives a relatively large dose without manifest change, it is at least probable that individuals exist who form a graded series between these two extremes. And equally it is probable that what is true of one drug—for example, atropine—is true in a greater or less measure of all others. Such a conclusion, further, has the support of experience, and hence it seems desirable to emphasize conspicuous drug idiosyncrasies not as extraordinary and exceptional occurrences, but as impressive illustrations of a general law, the existence of which applies over the whole field of drug administration. Such a presentation involves an exhortation to the prescriber to free himself from the yoke

of a rigid and authoritative scale of doses, and to make himself ready, both in one direction and the other, to adjust his quantitative orders to the idiosyncrasies which exist, not in exceptional cases only, but in each individual patient.

The question now to be asked is, What influence is exerted upon the view of drug idiosyncrasy as just defined, and upon the practice which is the inevitable outcome of such a view, by the erection of an official scale of doses in the national *Pharmacopoeia*? The first remark which arises on such a question is that the two schemes appear to be in absolute opposition to one another. In one flexibility, and in the other rigidity, is the predominant note. It is submitted that as a matter of fact the pharmacopoeial doses do exercise a prejudicial influence upon the habit of the practitioner in the quantitative adjustment of his remedies in dealing with the individual patient. These doses are for the most part committed to memory by the student in anticipation of the demands of the examiner. A similar consideration determines their appearance in the pages of the current manuals of materia medica and therapeutics. And the examiner, it may be hinted with all due deference, not infrequently adopts an attitude which justifies these procedures. Thus in the early and flexible phases of the practitioner's career a more or less arbitrary scheme of doses is stamped upon his memory, and these are the more impressive as they appear to have an authoritative sanction. Is it surprising to find that in many cases these powerful influences exercised on plastic and receptive minds become stereotyped as the habits of a lifetime, and the rigid rules which may be necessary for the inexperienced remain to limit the freedom of the man of full stature?

It is therefore argued that the existence of an official scheme of doses, so far from favouring the cultivation of a free and responsible personal judgement in the selection of the quantities of remedial agents suitable for administration, tends to oppose such a habit, and to encourage mechanical and arbitrarily-restricted methods of prescription. It suggests to the practitioner an appeal to authority rather than a study of the personal peculiarities of the patient, and it indicates a limitation of dose by precept instead of by physiological and therapeutical effects. The dangers of over-dosage attract many warnings, but who shall tell the disasters which have attended the timid and restricted prescriber? If, therefore, as will be admitted, the careful adaptation of remedies, both qualitative and quantitative, to the wants and idiosyncrasies of the individual patient is an essential part of good practice, and if, as will also be admitted, each individual patient has his own claims in this respect, it is obvious that conformity to a stereotyped scale of doses is out of harmony with the liberty and power of scientific prescribing. In other words, the existence of a scale of official doses encourages neglect of some of the conditions which favour the successful use of drugs in the treatment of disease.

Now the logic of all this would appear to be an appeal for the removal of the official doses from the pages of the *Pharmacopoeia*. Were the desired end to be attained only by such a step, the present writer would not hesitate to adopt it. At the same time, it would have to be recognized that this policy would remove one of the arrangements by which an accidental error in dosage is prevented. The existing plan affords an added protection to the public by forming a platform on which the dispenser can approach the physician should the latter by a slip of the pen have ordered a dose manifestly excessive and dangerous; and, in view of this, the suppression of the official doses cannot be lightly advocated.

Further, it may be urged that the disadvantage which is attached to their existence, and which has already been discussed, may be avoided by a method short of wholesale abolition. These stated doses may remain as a convenient opportunity for communication between the dispenser and prescriber, but they should be plainly recognized as having little other significance. In particular, they should be deprived of any suggestion of official force. In the preface to the *Pharmacopoeia*, it is true, the real position and significance of the so-called official doses is recognized and defined. But the limitations and qualifications there introduced are not vividly before the minds of medical students, nor are they always emphasized in the manuals of materia medica or in the discipline

*A paper read before the Therapeutical Society on November 28th, 1905.