In case of hæmoptysis with a high-tension pulse, could we rationally, and with a due sense of our responsibilities, give so powerful a pressure-raiser as the suprarenal? And in lung bleeding with low pressure—probably the more usual case—should we not rather assist Nature by lowering circulatory tension by any means in our power than empirically use a drug which, though powerful for much good, when rightly and rationally used, is also capable of much harm.

MYCETOMA (MADURA FOOT) IN THE YEMEN.

By FRANK G. CLEMOW, M.D., D.P.H.,

CONSTANTINOPLE.

THE geographical distribution of mycetoma, or Madura foot, is very imperfectly known. At one time believed to be confined to the Indian peninsula, it has more recently been shown to exist in many other tropical and semi-tropical countries. So far as I am aware, no authentic evidence of its presence in Arabia has hitherto been published. Suspecting its possible existence there, I last year requested some of the medical officers who are yearly sent from Constantinople to the Camaran lazaret for the period of the Moslem pilgrimage, to inform me of any case or cases which might come under their observation, particularly in native inhabitants of the neighbouring provinces of the Hedjaz, Assyr, and the Yemen. I am much indebted to Dr. Pappathanassiou, a member of last year's and of this year's missions to Camaran, for a careful record of ten cases, together with a number of photographs, which show conclusively that mycetoma exists, and is perhaps common, in the Yemen. The following is a brief summary of these cases.

CASE I.—Seit Abdou, aged 16, Imam, from Zodiah. Site of the disease, the right buttock. Has been ill for five years. He attributes the disease to the bite of an insect. He was operated upon in 1903, but the disease returned. When seen on October 4th, 1904, there were all the signs of mycetoma of the black variety. A second operation was performed on October 25th, and he left Camaran fifteen days later completely cured.

CASE II.—Hassan Hussein, aged 26, field labourer, from Zodiah. Site of the disease, the left foot; has suffered for seven years. He attributes the disease to the bite of a snake on the sole of the foot. When seen on February 3rd, 1905, the whole foot was swollen; the big toe alone was healthy. From a number of sinuses an oily liquid was constantly flowing, containing some pus, a little blood, and coarse black grains in suspension, resembling coarse gunpowder. On February 5th the foot was amputated, and seventeen days later the patient left the lazaret in good health.

CASE III.—Hassan Abdou, aged 28, field labourer, from Wady Aghem, near Hadjour. Had been ill for eight years. He also attributed the disease to the bite of a snake on the sole of the foot. When seen on February 3rd, 1905, the right foot presented much the same appearance as in Case II, save that all the toes were healthy. On February 6th amputation was performed, and fourteen days later the patient left the lazaret cured.

CASE IV.—Moeddin, aged 22, field labourer, from Zodiah. The disease began near the right elbow two years before. No cause could be assigned for it; he first noticed, while bathing, some small swellings on the external surface of the lower third of the upper arm, which later extended to the middle third. When seen on February 8th, 1905, there were three or four such tumours, with one or two sinuses from which an oily liquid escaped; this liquid was then clear or slightly purulent, but the patient stated that some time before it had contained in suspension black grains. The patient left without consenting to any operative treatment.

CASE V.—Otabi, aged 55, field labourer, from Loheia. Has been ill for ten years; he ascribes the disease to the bite of a snake in the left foot. When seen on February 20th, 1905, the foot presented all the signs of mycetoma of the yellow variety. On February 22nd the foot was amputated, and twenty-one days later the patient left the lazaret cured.

CASE VI.—Seit Ahmet, aged 38, field labourer, from Badjel. He also ascribes the disease to the bite of a snake ten years ago. As in the other cases attributed to a similar cause the swelling began at the site of the bite, was of very slow growth, and only after several months began to soften and break down, with the formation of sinuses and the discharge of an oily liquid, containing a little pus, some blood, and black grains in suspension. When first seen on April 8th, 1905, the left foot was in an advanced stage of mycetoma (see Fig. 1); amputation was performed on April 10th, and seventeen days later he left the lazaret cured.

CASE VII.—Mollah, female, aged 57, bedouin from Zodiah. The disease began in the right foot seven years ago. When seen on May 4th, 1905, only a cursory examination could be

made, but it was seen that the foot was much swollen, and sinuses were present from which an oily liquid was flowing,

Fig. 1.—Case vi.

the woman's companions stated that the fluid had contained black grains, and some of these were produced wrapped in a rag. The woman refused operation and disappeared.

CASE VIII.—Selim, aged 32, field labourer, from Zodiah. The

Fig. 2.—Case ix.

disease began in the right ankle ten years ago. When seen on June 2nd, 1905, the foot presented all the signs of mycetoma of the black variety. On June 4th amputation was performed, and twelve days later he left the lazaret cured.

Fig. 3.—Case x.

CASE IX.—Omah, aged 30, field labourer, from Zodiah. The disease began seven years ago in the left foot. When seen on

June 8th, 1905, it was very far advanced (see Fig. 2); the foot was enormously swollen, and from numerous openings an oily liquid was escaping in abundance, in which yellow particles, resembling fishes' roe, were suspended. Amputation was performed on June 10th.

CASE X.—Dr. Pappathanassiou has since sent me photographs of a tenth case—that of a woman from the neighbourhood of Zodiah. From one of these photographs (Fig. 3) it can be seen that the disease is present in the right hand, and has begun in the right foot.

In all the above cases, where amputation was performed, the pathological appearances characteristic of mycetoma were found to be present on cutting into the tumour. It was unfortunately not possible to make microscopic observations.

The above cases may be briefly summarized as follows: of the 10 cases, 8 were in males and 2 in females; 7 of the males were field labourers and 1 an Imam; 7 came from Zodiah or its neighbourhood, 1 from Wady Aghem, 1 from Loheia and 1 from Badjel; in 4 the left foot was affected, in 3 the right foot, in 1 the buttock, in 1 the right elbow, and in 1 both right hand and right foot; the duration of the disease had varied from two to ten years; in 7 the disease was of the black variety, in 2 of the yellow variety, and in 1 the colour is not stated; in 4 the disease was ascribed to a snake bite, in 1 to the bite of an insect, and in the rest no cause for it was suggested by the patient.

All the above-named places are situated in the Yemen, within a comparatively short distance of the Red Sea coast. Loheia is one of the principal ports of the Yemen; Wady Aghem is some thirty hours inland from Loheia; Zodiah (where the majority of the cases came from) is three and a half hours inland from Ibn Abbas, which lies on the coast, nearly opposite to Camaran, and between the ports of Loheia and Hodeidah; Badjel is ten hours inland from Hodeidah. All the cases were therefore drawn from within a comparatively short radius of the Camaran lazaret, where the known presence of European medical men would naturally tend to attract such cases from the neighbouring mainland. It still remains an open question whether the disease also exists further inland in the Yemen, and in the Hedjaz, Assyria, and the other Arabian provinces.

Since writing the foregoing I have myself seen the following two cases during a fortnight's stay in Camaran:

The first case was that of an Arab boy, aged about 16, from Deir-Mohammad, near to Zodiah, the district from which so many of the cases already reported came. The disease was said to have begun three years ago in the sole of the left foot. The foot was in an advanced stage of mycetoma, of the black variety, and there was marked marasmus.

The second case was that of an older man, aged about 25, also from Zodiah. Although the disease was said to have appeared four years earlier, it had only affected the middle toe of the right foot. It was said to have begun on the upper surface.

These cases are a further proof of the existence—and indeed of the frequency—of Madura foot in the Yemen province. I cannot hear that it is met with in the Hedjaz.

THE TEETH OF SCHOOL CHILDREN.—In an official report of Mr. Joseph J. Brittain, United States Consul at Strassburg, quoted in the *Medical Record*, he refers to the good results of the establishment of a dental clinic for school children in that city. In 1902 a clinic was opened for the care and treatment of the teeth of the children in the public schools. Attendance at the clinic is compulsory, and the children are treated free of charge. In the first year 5,343 children were examined and 2,666 received treatment. The third annual report, just published, states there were 12,691 visits to the clinic in 1904 and 6,282 children were treated, for whom 7,065 teeth were filled and 7,985 teeth were extracted, and 4,382 other children had their teeth examined. The children are taught to clean their teeth three times daily, and especially before going to bed. The dentist also instructs the children in the use of the toothbrush, each child receiving a brush for home use. Since the introduction of the treatment there has been a marked improvement in the general health of the public school children. Since the foundation of the Strassburg clinic similar establishments have been opened in Darmstadt, Mulhausen, and other cities in Germany.

MEMORANDA: MEDICAL, SURGICAL, OBSTETRICAL.

ANGINA PECTORIS WITHOUT APPARENT DISEASE OF THE HEART OR BLOOD VESSELS.

DR. CURTIN is so good as to invite me to comment upon a case—a remarkable case—of angina pectoris.¹ He considers that his case stands against my hypothesis that the seat of origin of angina pectoris is the first portion of the aorta, and lies not in the heart itself. As Dr. Curtin's skill was successful in saving the life of his patient, the nature of the attack, decisive enough clinically, must remain pathologically ambiguous. But Dr. Curtin thinks the case clear enough for some argument. He surmises, and probably with truth, that in his case the intense irritation of a sensory nerve or nerves had its common consequence of a widespread arterial constriction, "with a consequent damming back of the blood on the left ventricle, which, owing to the attenuated condition of its walls, became suddenly stretched, and therefore unable to bear the stretch on its increased superficial area."

I will not follow Dr. Curtin into "pseudo-anginas," which are no more angina pectoris than colic is appendicitis.

It may be my fault, but on repeated perusal of Dr. Curtin's note I fail to find any argument which, while supporting his postulate of excessive intraventricular pressure, excludes that of excessive intra-aortic pressure; or which pits one against the other. Yet surely such a difference is of the essence of his discussion? I would dwell on excess of pressure in one area, Dr. Curtin on this in the other. If I admit his plus intracardiac pressure he must surely admit my plus intra-aortic pressure?

Again, it is the misfortune of both of us in this instance that Dr. Curtin did not see his patient during the attack. A statement of the behaviour of the heart during the attack is of the first importance, and we are without it. We rarely see our patients in these attacks, yet in this good hap I have been more fortunate than most physicians, and am able to testify that in the large majority of seizures the heart goes on its way with but little perturbation. Fifteen minutes after the attack Dr. Curtin finds the heart "intermittent, losing about one beat in four"—that is, the reflex (vagus) inhibition, which might well have stopped the heart in the seizure, still showed traces of activity. (This reflex, I admit, may have been either splanchnic or aortic, or, again, as I rather suppose, binary, with an unknown distribution of the responsibility between the two areas.)

If Dr. Curtin feels as insecure as I do in estimating the dynamic value of the heart by its physical signs, he will not pronounce upon the value of this patient's heart; it seems, at any rate, to have been good enough to ride through a terrific storm. So, if Dr. Curtin is disposed to attribute some defect to the fabric of this heart, I am disposed to suspect also some defect in the spring of the arch of the aorta of a man aged 59. I consider one surmise as good as the other, and my surmise is supported by my review of the whole subject in my Cavendish Lecture, a review I cannot of course repeat here. I must content myself with one quotation, and this is from Dr. Curtin's own note: "The most typical condition for angina pectoris is aortic regurgitation and stenosis, without mitral lesion"—that is, as I put it, a diseased aorta at a moment of excessive arterial pressure, the excess being either positive or relative to the morbid state of this part. That before the attack the pulse was of "low tension," even if Dr. Curtin attaches more value than I do to such an impression, does not affect the argument of either of us. Nor can we attach much importance to "softness" of the radial artery as an indication of the histological factor in other arterial areas.

In conclusion, I will venture to comment upon the way of nearly all writers in confusing the pain of angina pectoris with "feeling of dyspnoea, tension, or oppression about the heart . . . pain in the region of the apex," etc. (the italics are mine). As in Dr. Curtin's case, the pain in unmixed angina pectoris is not epicardial but substernal; and "dyspnoea" is not a feature of it.

Cambridge.

T. CLIFFORD ALLBUTT.

¹ BRITISH MEDICAL JOURNAL, April 14th, 1906.