it is clear that this portion of the posterior root is more liable to show signs of impaired nutrition at an early stage than either the extramedullary portion or the cells of the spinal ganglia. Some conditions probably exist favouring the incidence of noxious influences on this portion of the sensory protoneurone. In this connexion Marie and Guillain have laid great stress on the peculiarities of the lymph circulation in the posterior columns of the spinal cord. They maintain that in the pia mater of the posterior columns, between the outer and inner layer, there exists a lymphatic system, the spaces of which communicate with those of the posterior columns, but are independent of those of the antero-lateral columns. In this system the current of lymph is upwards and towards the central canal. In tabes they have found this system dilated, and, by the Marchi method, black granules in the cells of the ependyma in cases where the central canal is permeable. They attribute the condition to a syphilitic affection of the posterior lymphatics, and derive the lymphocytosis of the cerebro-spinal fluid from the elements of this system.

That noxious substances or toxins originating in the periphery ascend to the spinal cord more readily by the posterior than by the anterior roots has been shown by Homén, and also by Orr and Rows; and that in this way, or by peculiarities in the lymphatic circulation in the cord, the posterior roots and posterior columns are specially exposed to injury is extremely probable. But that there is in tabes a posterior meningitis of a syphilitic character, as maintained by Marie and Guillain, is not generally admitted. It is not supported by the results of anti-specific treatment, nor can the lymphocytosis of the cerebro-spinal fluid be adduced as a conclusive argument in its favour.

[Bibliography will be given at end of third lecture.]

Demonstrations

ON

THE CYSTOSCOPE

AND A METHOD OF ILLUSTRATING DISEASES OF THE BLADDER AND THE APPEARANCES IN RENAL DISEASE BY THE OPAQUE PROJECTOR.*

By **DAVID NEWMAN, M.D., F.F.P.S.G.,**
SURGEON, GLASGOW ROYAL INFIRMARY.

[WITH SPECIAL PLATE.]

PART II.—CYSTOSCOPIC APPEARANCES IN SOME DISEASES OF THE BLADDER AND KIDNEYS.†

In the cases to be presented to you to-day I have selected those suited to illustrate the conclusions to be submitted.

Fig. I.—Normal ureter opening.

the mouth is easily recognized on the ridge of the trigone.

I have avoided taking what might be regarded as "typical cases," choosing by preference (with one or two exceptions) those most commonly met with in practice.

Fig. I represents the appearance of the normal ureter opening and the mucous membrane of the bladder when the viscus is distended to 6 fluid ounces. The lips of the opening are distinctly seen and slightly parted so that

A. CALCULUS.

1. Stone Lying Free in Bladder.

Probably the most easily-observed morbid object within

* The second of a series of Post-graduate Demonstrations given at the Royal Infirmary, Glasgow.

† The writer has much pleasure in owing his indebtedness to the Editor of the *Practitioner* for the use of the blocks to illustrate Figs. III, X, XII, XVIII, XXIV, and XXVI.

the human bladder is a stone lying on its floor (Fig. II). The presence of such a foreign body can easily be discovered by means of a sound, but its probable composition, its form and size can only be accurately determined by the cystoscope. The spherical, smooth phosphatic calculus caused comparatively little irritation, and its presence did not induce much change in the appearance of the mucous membrane of the bladder.

Fig. II.—Stone loose in bladder.

The patient had complained of symptoms of stone for over two years, the most pronounced being pain at the end of micturition, but the suffering was not sufficient to preclude him from following his occupation (a salesman in a grocer's shop) until within two weeks of the time I saw him. He sought advice on account of a hæmorrhage; when examined the stone was at once seen, and the following day it was removed.

In many cases, however, without the aid of the cystoscope, stone may elude detection, either from being covered by a thick layer of mucus or by being placed in such a position that it cannot be reached with the sound.

2. Stone Obscured by a Thick Layer of Mucus.

When the surface of the stone is regular but finely honeycombed without having prominent projections, thick mucus may become

adherent to it, so that it presents the appearance of being covered with wet cotton-wool (Fig. III), and this mucous layer may form a very efficient protection to the mucous membrane of the bladder. I have seen several instances of this kind, some where a stone has been present in the bladder for a considerable period without inducing signs of irritation, others where the symptoms have abated in a most remarkable way, as a consequence of the protective coating.

Fig. III.—Stone obscured by mucus.

Fig. III is an illustration from a case in point. The history was that of chronic cystitis, frequent micturition, pyuria, but no hæmaturia. The pain was so slight that the presence of a stone was not suspected. The case was rather regarded as one of septic cystitis. I found, however, a white woolly mass occupying the floor of the bladder, the shape and size of a pigeon's egg. From its appearance and form, the presence of a stone was suspected, but on passing a sound into the bladder no hard object could be found. The bladder was washed out regularly twice daily for a week, and a considerable quantity of mucus was in this way removed. Then a second examination was made, when a phosphatic stone was seen lying on the floor of the bladder, only with a light covering of loose flaky mucus.

3. Calculus in Sacculæ.

A calculus may be hidden in a sacculæ or diverticulum in which to find a stone may be extremely difficult (Coloured Plate, Fig. IV). The mucous membrane of the bladder was practically normal. The colour, except immediately around the opening of the sacculæ, was pale, but the lips of the opening were deeply congested, rounded, thickened, and, in marked contrast to the bright red lips of the orifice of the diverticulum, a pale white object was seen.

In such cases the difficulty when first seen is to determine whether such a white object is a stone or a tuberculous ulcer covered with a slough, but by carefully washing out the bladder the mucus covering the stone is removed and by passing the finger into the rectum or the vagina, while at the same time observing through the cystoscope, if the object be a stone, the bladder wall

is pushed forward in a degree represented by the size of the stone and not by that of the point of the finger.

4. Stone behind Prostate.

Not uncommonly a stone is hidden behind the prostate (Coloured Plate, Fig. V), and can only be detected with a sound when the patient is placed in the elbow-knee, or in the Trendelenburg position. The case from which the drawing was taken was one of enlarged and greatly congested prostate, a stone was suspected but could not be detected with the sound, but was readily seen with the cystoscope.

5. Calculus Impacted in the Orifice of the Left Ureter.

The following case is an instance of this condition:

There was evidence of obstruction to the escape of urine over a period of fully four years, and the family medical attendant correctly diagnosed a recurring hydronephrosis. For the first two and a half years the attacks came on about once in four months, and were neither prolonged nor very severe, but during the last eighteen months the patient had never been free from pain for a longer period than six weeks, and sometimes the pain recurred as often as twice in the fortnight. The attacks of pain in the lumbar region were associated with all the symptoms of hydronephrosis, but on no occasion within the first three years since the onset of the disease was blood discovered in the urine. The first haematuria was on January 3rd, 1900. The bleeding at that time was very slight, but, on account of a rather profuse haematuria, I was asked to examine the patient. On cystoscopic examination the mucous membrane of the bladder was found to be slightly congested, and the arterioles were more numerous than normal. The mucous membrane was slightly thickened, and, considering the degree of distension, the folds were numerous and distinctly marked. The line of the ureter was unduly prominent and rounded, and at the orifice somewhat club-shaped. The mouth of the ureter was widely dilated, and between its lips an irregular-shaped, pointed calculus was seen (Coloured Plate, Fig. VI). The surface was of a dark brownish-red colour, roughly nodulated, and frosted over with a slight covering of phosphates. A suprapubic operation was performed and the calculus removed.

B. BALLOONING OF URETER.

1. Slight Ballooning of the Mucous Membrane due to Impaction of a Small Calculus.

The history was that of repeated attacks of renal colic without suppression of urine, but followed by haematuria, marked irritation of the bladder, frequent but not painful micturition, and latterly almost constant presence of blood. There were no renal symptoms. The swelling in the bladder was at first diagnosed a tumour, but at the operation it was discovered to be a small calculus included in a sac formed of the mucous membrane of the bladder. The following is a short history of the case:

J. H., aged 35, a marine engineer, first suffered from renal colic for about twenty-four hours only, on a homeward voyage from India in December, 1893. Complete relief from pain followed, and lasted for seven weeks. Travelling from London to Leeds brought on a second seizure, which lasted for five hours, but although shorter in duration, the suffering was more severe in this than in the first attack, and when the pain passed off the patient noticed that the urine contained blood. During 1894 the patient suffered from seven attacks, three of which were followed by haematuria. At no time were there any symptoms suggestive of suppression of urine nor of hydronephrosis. Towards the end of 1894 the patient complained greatly of frequent micturition; during the act he suffered little or no pain, but on many occasions blood appeared in the urine in moderate quantities. Only twice could the haematuria be described as profuse. There were no symptoms pointing to any renal lesion, nor could swelling or tenderness be made out in either lumbar region. No information was gained by rectal examination. No tube casts and no albumen were in the urine except from the presence of blood. A cystoscopic examination showed the presence of a rounded swelling immediately over the orifice of the right ureter, about the size of a small cherry, very vascular on the

Fig. VII.—Ballooning (slight).

surface, but smooth and free from ulceration (Fig. VII). The swelling was diagnosed as a fibroma of the bladder pressing upon the orifice of the ureter, and so causing an

obstruction to the escape of urine. Suprapubic cystotomy was performed, and the swelling was seized with ring vulsellum forceps. Immediately pressure was made the swelling cracked, and a little traction brought away a calculus, and a thin but complete covering of mucous membrane.

2. Extensive Ballooning of the Mucous Membrane of the Bladder (Fig. VIII).

The patient was sent to me with a very complete history of her illness, and the conclusion her medical attendant came to was that there was some obstruction to the escape of urine from the right ureter, and he suspected an impacted calculus.

The symptoms were those of intermittent obstruction to the ureter. Beyond the elevation of temperature during the acute attacks, and the presence of pus in the urine, pain in the right renal region and in the bladder was the only symptom which attracted attention. Suspecting a calculous obstruction to the right ureter, I made a cystoscopic examination on November 22nd, 1905, when a swelling was found occupying the right half of the bladder. The surface of the swelling was paler than that of the vesical mucous membrane, perfectly smooth, translucent, and meagrely supplied with blood vessels. The swelling was distinctly pedunculated, floated in the fluid contents of the bladder, and moved freely with respiration (Fig. VIII). The bladder, distended with 10 oz. of solution, allowed the greater part of the rounded swelling to be seen, but the site of its attachment to the bladder wall could not be ascertained. On opening the bladder in front, the cyst was found to be about the size of a duck's egg, and to have a pedicle about $\frac{1}{4}$ in. in diameter. On pressure between the fingers it was readily emptied, and its fluid contents escaped into the bladder, after which a round opening was discovered in the bladder wall, in the situation of the orifice of the right ureter. It was simply a sac protruding into the cavity of the bladder. The precise cause of obstruction was not ascertained, but a careful examination failed to reveal the presence of a calculus. The sac was incised, a tube was inserted into the bladder, and the urine was drained off until November 28th, when a second examination was made with the finger, and by means of a speculum through the wound. The cyst was found to have completely collapsed. The wound in the parietes was sutured, allowed to heal; the patient made a good recovery. Prior to returning home a cystoscopic examination was made, when no trace of the cyst could be found beyond some wrinkling and irregularity of the mucous membrane on the right side of the bladder. The orifice of the right ureter was represented by an irregular lacerated opening.

Fig. VIII.—Ballooning (large).

Cases of ballooning of the ureter have been published from time to time as pathological curiosities, but I think this is one of the first in which such extensive ballooning has been observed during life, and for which an operation has been performed.

C. TUMOURS.

1. Papillomata.

Papilloma is one of the most common tumours of the bladder, and arises from the mucosa and submucosa. At present there are three cases in the ward and these I will show you at the next demonstration. The tumour may simply be a large group of villi, with an extensive vascular network, and attached to the bladder by a long pedicle, or the fibrous tissue elements may predominate and the papilloma consist of a sessile wart. The illustration (Fig. IX) represents one of the former class.

The patient was admitted to the Royal Infirmary on October 20th, 1905, complaining of blood in his urine and pain of two years' duration. About two years ago he caught a chill, and soon afterwards noticed that blood was present in the urine, which lasted for

Fig. IX.—Papilloma.

two or three days at a time. Recurrence took place only at long intervals until within the last six months. Since then blood had been noticed to be present two or three days every week, sometimes the haemorrhage being so severe that the evacuation appeared almost like pure blood, and frequently it contained large clots. The pain generally came on immediately after the act of micturition, was shooting in character, but no pain was present before or during the act. Immediately prior to admission the pain became so severe as to be almost unbearable, extending on some occasions to both sides of the back. During the last two years the patient has lost considerably in weight, has become anaemic, but otherwise there is nothing of special note.

On October 24th a very clear view was obtained of the bladder, the mucous membrane of which was pale in colour and was ramified by unusually large veins, which, however, were not greatly distended with blood at the time of the examination. On the left side of the bladder there was a large pale papillomatous growth, with long fimbriated processes, which floated freely in the solution.

Fig. IX represents the appearance of a small portion of the tumour with a shaded part of the bladder in the background which throws the soft fimbriated processes prominently into relief. During the examination there was no bleeding, and beyond the presence of the tumour the bladder was strictly normal.

2. Epitheliomata.

Epitheliomata usually attack the base in the vicinity of the ureteral orifices, and as a rule remain solitary. In the case from which Fig. X (see Coloured Plate) was taken, two separate tumours were seen—a large one close to the mouth of the left ureter, and a small isolated nodule on the left side of the middle line posteriorly. Both growths were similar in appearance, differing only in size, and were quite free from villous projections on the surface; the margins irregular in outline, elevated, overlapping the mucous membrane, coarsely granular, and deeply pigmented. The surface of the tumour was flat, finely nodulated, and covered by glistening material, like pure mucus. At one or two small points ulceration was seen. The symptoms were frequent micturition, and occasionally slight bleeding. There was almost no pain, and, until discovered by the cystoscope, the presence of the tumour was not suspected. The growth was rather an unusual one to find in the bladder, namely a flat-celled epithelioma, of dense consistence. Further examination showed that the growth involved to a slight degree the walls of the rectum; there was no breach of surface, but with a sound in the bladder and a finger in the rectum, the connexion was easily established. The case was not a suitable one for operation, on account of secondary involvement of lymphatic glands.

3. Fibromata.

Fibroma of the bladder is also a rare disease, but probably some of those growths described as myomata are really fibrous tumours, such as the one about to be described. Little was known of the history of the case, except that the urine was normal at all times, with the exception of three occasions when it contained blood; the last haematuria, being considerable, led to a cystoscopic examination, when the tumour was discovered attached to the anterior wall of the bladder. The growth (Fig. XI) was about the size of a horse-bean, pale in colour and regularly nodulated on the surface, almost like a yellow raspberry, and the colour vividly contrasted with the deeply-injected mucous membrane which formed

Fig. XI.—Fibroma.

the background. The growth was pedunculated and easily removed.

4. Sarcomata.

This variety of tumour in the bladder was considered to be very rare; out of 41 cases of tumour, Thompson found 3 of sarcomata; Albarran, out of 82 cases, 2; Guyon in his practice met with 1 case, and that *post mortem*; and I have seen only 2.

The most characteristic case was that of a man, aged 26, who had been operated upon several times for "recurrent fibroids," one on the right knee, a second on the hip, and a third on the right lumbar region. When I saw him for the first time there were seven such tumours below the level of the waist, and one over the left shoulder in front. He complained of frequent and painful micturition, and had occasional attacks of haematuria. Cystoscopic examination revealed a large rounded tumour on the anterior wall of the bladder (Fig. XII). In colour the surface was mottled bluish-red, and there was an irregular lozenge-shaped pale-blue patch almost in the centre; over the surface there was a large number of small capillaries. The mucous membrane was deeply injected and the vessels dilated. Operation was deemed inadvisable, but a section of the tumour *post mortem* showed it to be a round and spindle celled sarcoma.

Fig. XII.—Sarcoma.

In 24 cases collected by Hintertroisser, it has been shown that these growths arise from the submucosa and muscular layers, that metastasis is comparatively rare, and that the sarcomata are relatively most frequent in childhood.

5. Polypi.

Mucous polypi, which have been erroneously described under the term myxomata, are found usually in young children, and may protrude from the urethra if the patient be a girl.

Fig. XIII (Coloured Plate) shows a polypus hanging from the bladder, about the level of the arch of pubis. The symptom in this case was due to the ball-valve action of the polypus. The bladder could not be completely emptied on account of the polypus hanging into the urethra. After spontaneous micturition, when a catheter was passed, fully 7 oz. of healthy urine could be drawn off. This was a constant feature in the case. The cystoscope showed a translucent pale growth hanging from the anterior wall of the bladder by a narrow neck, and on any agitation of the fluid contents the tumour swung backwards and forwards freely. The polypus was removed through a suprapubic incision.

D. ULCERATION.

1. Solitary Ulcer with Necrosis of a Portion of the Mucous Membrane (Coloured Plate, Fig. XIV).

The symptoms in this case were frequent micturition, pain on and for a short time after micturition, especially marked at the base of the penis, and occasionally followed by distinct but not very severe spasm. During the day the symptoms were not marked when the patient took large quantities of fluid, and the specific gravity of the urine remained low—say 1010 to 1015; but from 8 p.m. till 8 a.m., although fluid was taken freely, the quantity of urine was persistently small and the specific gravity almost constantly high. Whenever the specific gravity was above 1020 the patient complained of pain, and was unable to retain more than 4 or 5 oz., and when the specific gravity rose to 1030—as at the onset of the disease it did almost every night—spasms of pain occurred, lasting usually for ten or fifteen minutes. The mucous membrane of the bladder was only slightly

COLOURED ILLUSTRATIONS.

- IV. Stone in sacculi.
- V. Stone behind prostate.
- VI. Stone in ureter.
- X. Epithelioma.
- XIII. Mucous polypus.
- XIV. Solitary ulcer.
- XV. Tuberculous ulcer.
- XIX. Pneumococcus infection (right side).
- XX. Pneumococcus infection (left side).
- XXI. Acute septic pyelitis.
- XXII. Acute tuberculous pyelitis.
- XXV. Ascending ureteritis.

Fig. IV.

Fig. V.

Fig. VI.

Fig. X.

Fig. XIII.

Fig. XIV.

Fig. XV.

Fig. XIX.

Fig. XX.

Fig. XXI.

Fig. XXII.

Fig. XXV.

injected, and there was little change in its appearance, except around the ulcer, which was situated on the posterior wall. The ulcer (Coloured Plate, Fig. XIV) was solitary, irregular in shape, and extended from 165 to 180 degrees to the left of the middle line. It was superficial and its edges very slightly raised. The margin of the mucous membrane was of a bright red colour, and contrasted very clearly with the pale yellow slough which covered the ulcer. Passing outwards from the margins, the coloration gradually became paler, until it merged into that of the pink mucous membrane. The urine passed during the night contained pus, blood, and gave pure cultures of the *Streptococcus pyogenes*; what was evacuated during the day was clear, and a deposit could only be obtained by using the separator. After treatment for eight weeks the symptoms disappeared. The urine became normal and the ulcer was found to be healed, leaving only a slight pale scar.

2. Tuberculous Ulceration.

In the early stage this may readily be mistaken for solitary ulcer.

In the case from which Fig. XV (Coloured Plate) was taken, the symptoms were very insidious at the beginning, consisting of frequent micturition, evacuation on an average every two hours, and pain at the commencement and at the end of the act only. The urine was generally clear in colour, but what was passed first in the morning contained more or less blood, almost no pus, and no organisms could be discovered by the microscope. At this time (May, 1902) the patient appeared in typically good health, and weighed over 14 st. Gradually the discomfort increased, and by August, 1902, the irritation was so great that he was confined to bed. Occasionally blood was present in the urine, but not in large quantities, and was not increased in amount by exercise or relieved by rest. Ten examinations were made for tubercle bacilli up till August 25th, 1902, with negative results. At that date several groups of bacilli were discovered, and subsequent inoculation experiments proved the presence of tuberculous infected matter in the urine. Cystoscopic examination revealed one large and several small tuberculous ulcers. They were situated within the area of the trigone. The largest was irregular in shape, and was covered by a white slough, while the margins were deeply injected, and of a bright red colour. The smaller ulcers were rounded, with minute caseous centres. The ureter orifices were normal in appearance. Suprapubic cystostomy was performed, the ulcers curetted, and when seen in April, 1905, the patient could retain urine for four and a half hours, suffering no discomfort, and the urine was free from tuberculous infected matter, as proved by inoculation.

3. Tuberculous Infiltration.

In this case there was recent tuberculous deposit close to the neck of the bladder (Fig. XVI).

The symptoms did not point to any serious bladder lesion, the only inconvenience complained of being frequent micturition. The urine contained a trace of albumen, tubercle bacilli were found in considerable numbers, it deposited a minute quantity of pus, but no tube casts were discovered. The kidneys were free from disease, and no evidence of tuberculous disease could be discovered except in the bladder. The mucous membrane of the bladder was moderately injected, and the vessels in its walls were more numerous and larger than normal. Extending along the floor of the bladder from a little below the orifice of the left ureter to the neck of the bladder there was a pale mass, like oedematous granulation tissue, and in it a number of small, white, caseous points were seen; these were tubercles just beginning to break down. There was no distinct ulceration. Two years ago the urethra was dilated, and the whole of the tuberculous deposit was removed with a curette. The bladder was washed out regularly for ten days, after which silver nitrate solutions of varying strength were injected at intervals. The patient made a good recovery, and when seen in August, 1905, there had been no recurrence of the disease.

Fig. XVI.—Tuberculous infiltration.

4. Advanced Tuberculous Ulceration.

The history of the case, stated briefly, was:

Gradual increasing frequency in micturition, at first most marked in the afternoon, if the patient had been active during the day. Rest in bed afforded relief at first only; after a time no benefit was derived from it. Two months after onset of urinary trouble suddenly the irritation in the bladder increased, and was accompanied by persistent pyuria and occasional haematuria. Temperature febrile. Constant pain at the same time was complained of in the right lumbar region, short remissions of the suffering only following evacuations of highly purulent urine. There was evidence of tuberculous disease of the right ureter and left testicle, and of chronic pulmonary tuberculosis limited to the left apex. Cystoscopic examination revealed at some parts considerable thickening and marked hyperaemia of the mucous membrane, while at others it was thin and anaemic, thus presenting a peculiar coarsely-mottled appearance (Fig. XVII). There was considerable ulceration at the base; some of the ulcers were rounded, covered with grey and white sloughs, and their edges were undermined, others were irregular in shape, with bright red margins, and these surfaces were covered with blood and blood clots. The mucous membrane at the neck of the bladder was very highly coloured, swollen, and oedematous, while in the area of the trigone the mucous membrane was of a dark reddish-grey slate colour, with a large number of distended arterioles showing on its surface.

Fig. XVII.—Advanced tuberculous ulceration.

The cystoscopic examination was accomplished only after several unsuccessful attempts on account of haemorrhage from the neck, and difficulty in retaining sufficient fluid in the bladder.

The appearance presented by advanced tuberculous ulceration of the bladder, as viewed during life, contrasts markedly with what is seen *post mortem*; in fact, to see a bladder the site of tuberculous disease after death does not convey the slightest idea of the appearances revealed by the cystoscope. Unfortunately, in advanced tuberculous disease, the bladder is so irritable and contracted that one seldom has the opportunity of seeing what was observed in this case.

5. Malignant Ulcer.

In the case of carcinomatous ulceration at the base of the bladder, shown in Fig. XVIII, the point of interest was that, while the bladder symptoms were insignificant and the ulcer in the bladder small, the primary vesical lesion led to very wide secondary extension.

The cancerous disease was situated in the trigone, to the inner side of the orifice of the left ureter, and occupied an area less than an inch square. The mucous membrane was hyperaemic, but beyond the area above described it was not involved in the cancerous disease.

The only urinary symptoms were frequent micturition and very slight haematuria. The patient died of acute and widely disseminated cancerous disease, probably as a consequence of infection of the blood stream.

Fig. XVIII.—Malignant ulcer.

THE ORIFICE OF THE URETERS IN RENAL DISEASE.

The appearance of the normal ureter opening is shown in Fig. I, with the bladder distended to 6 oz. The mucous membrane was pale in colour, and was ramified by a few small-sized arterioles. The ridge of the trigone is marked, the lip of the orifice distinct, and the opening is easily seen. Observe, not only the form, but also the colour.

Beyond determining the precise nature and extent of vesical lesions, the cystoscope is most useful in enabling us to diagnose renal disease. Many years ago I pointed out the importance of ascertaining the appearance of the orifices of the ureters in diseases of the kidney, and stated that "when there is a distinct evidence of irritation of the orifice of one ureter, while the other is normal in appearance, it may be safely inferred that the renal lesion, if unilateral, is on the side of the morbid ureter."

From that original axiom we have during the last fifteen years made enormous advance; and, while it may be safe in some cases and rash in others to dogmatize as to the conclusions to be drawn from the appearance presented by the ureter orifice, in the majority of cases we may gain information of the highest value. In some instances it may be necessary to make more than one examination before a definite opinion can be expressed regarding the condition of each kidney, but a patient seldom raises serious objection when the importance of knowing the precise condition of both kidneys prior to operation is fully explained. When the second kidney is known to be healthy, the surgeon can recommend an operation, which he would not be justified in advising if he had not ascertained the condition of the opposite kidney. In ignorance of this he might very readily remove only one of two diseased kidneys, and lose his patient.

I cannot here enter into a full consideration of the various points to be attended to in an examination conducted with a view to ascertaining the presence of unilateral renal disease. I may here simply state that, in examining the mouth of the ureter, three main points demand attention: (1) The appearance of the orifice of the ureter; (2) the character of the material which escapes from it; and (3) the nature of the shoot as regards force, quantity, frequency, and rhythm.

A. ACUTE DISEASES.

A condition which has interested me within the last few years is one which has not been sufficiently recognized; a malady of a very grave nature, and sometimes involving the life of the patient. As it is one in which the appearances of the ureter orifices is very striking, I have selected it to illustrate my fundamental assertion that when disease is unilateral, it is on the side of the morbid ureter.

1. *Acute Unilateral Disease of the Kidney due to Pneumococcus Infection.*

The patient came under my observation on November, 1905, complaining of pain in the back in the region of the right kidney, but most marked over the last rib, where there was considerable tenderness on pressure. The onset of the pain was accompanied by slight oedema of the feet and face. It was dull and aching in character, but on admission to the "Home," the patient said that while she lay in bed on the right side the pain became relieved. There was no increased frequency in micturition, and the patient never complained of any pain on urination. On admission the temperature was 102.5°, and remained elevated for some days. On examination of the abdomen the right kidney was found to be much enlarged, slightly movable; on gentle pressure considerable tenderness was elicited, and on firmer pressure the patient complained of severe pain. The left kidney could not be felt, and certainly there was no enlargement, nor was there any pain on pressure. Urine specific gravity 1018, acid in reaction, albumen and blood present in considerable quantities; deposit—abundant granular tube casts, mucus, blood, and pus.

A cystoscopic examination was made on November 11th, when the mucous membrane of the bladder was seen to be normal, with the exception of hyperaemia in the area of the trigone, at the orifice, and along the line of the right ureter (Coloured Plate, Fig. XIX). The lips of the orifice were dilated, thickened and swollen, and extending upwards, and on either side there were a large number of widely-dilated capillaries. This gave the appearance as if the ureter orifice was covered by a pear-shaped veil of dark red lace, the apex of which pointed upwards. Within this deeper coloration there was a ring of paler pink. From the mouth of the right ureter dark blood-stained urine was seen to escape in small quantities at short intervals—on an average event twenty to thirty seconds. The left ureter orifice (Coloured Plate, Fig. XX) appeared as a little round opening, the mucous membrane surrounding it being normal in appearance, and from it clear urine flowed at intervals of about a minute and a half.

On November 15th the segregator was used and the following results obtained: From the right kidney, urine specific gravity 1025, acid, dark red colour, pus, blood, tube casts, finely granular tube casts present; almost pure culture of pneumococcus obtained. From the left kidney, urine specific gravity 1010, acid, dark yellow colour, no blood, pus, or tube casts,

slight trace of albumen, a few *B. coli* present and very few pneumococci. The general condition of the patient was that of one suffering from acute febrile disease. The skin was dry, the tongue furred, red at the point and margins, the pulse rapid, and she complained of loss of appetite, headache, and a constant inclination to be sick. From November 18th the symptoms began to abate, temperature was reduced, the enlargement of the kidney diminished, the tenderness on pressure less marked, the quantity of blood in the urine smaller, and the general condition more satisfactory.

On November 25th, on cystoscopic examination, the trigone was found to be practically normal in appearance, but there was still some thickening about the orifice of the right ureter, although the hyperaemia had subsided. The urine escaping from the right kidney was still darker in colour than that flowing from the left, although the urine passed from the bladder did not contain any blood. (The difference in colour was probably due to greater concentration of the urine from the right kidney.) The urine from the left kidney was pale in colour. Bacteriological examination failed to reveal the presence of pneumococci. No tube casts could be discovered, and only a slight trace of albumen. The patient ultimately made a good recovery, and the kidney became reduced in size.

2. *Acute Septic Pyelitis.*

The onset of the disease was very acute, and followed upon a carbuncle at the back of the neck, so that probably the infection was due to blood contamination. The illness began with a rigor and severe pain in the region of the left kidney. There was no evidence of increase in the size of the kidney, but pressure in the renal region caused pain; there was considerable irritation on micturition, and at the end of the act small quantities of blood escaped, but at this period there was no pus in the urine. The symptoms resembled those of renal calculus impacted in the ureter, but from the circumstances that the pain was suddenly relieved by the escape of pus, coincident with a fall in temperature from 103° to 100.5°, it was inferred that the symptoms were due to an infective lesion rather than to mechanical obstruction simply. On cystoscopic examination, the orifice of the left ureter was seen to be greatly dilated, the margins clear and sharply defined, the wide space between the lips deeply injected, and of bright red colour, in marked contrast to the pale pink mucous membrane of the bladder (Coloured Plate, Fig. XXI). The right ureter was strictly normal. From the left ureter pus mixed with urine was seen to escape in small quantities, at short intervals, the shoots sometimes being as rapid as fifteen seconds apart.

3. *Acute Tuberculous Pyelitis.*

The involvement of the kidney, which was secondary to phthisis pulmonalis, in this case came on very suddenly, with severe pain over the region of the right kidney. The symptoms were frequent micturition, with lacerating pain at the external meatus and in the perineum. On pressure over the kidney the patient complained of severe pain, but no swelling could be discovered. Prior to the present attack, at intervals there had been symptoms of cystitis for four months, and on vaginal examination a thickened right ureter was discovered. The urine was acid, specific gravity 1015, pale straw colour; contained a small quantity of albumen and a slight deposit of pus; no tube-casts nor blood. Microscopic examination revealed the presence of large numbers of tubercle bacilli both in the urine and in the sputum. A cystoscopic examination showed (Coloured Plate, Fig. XXII) the mucous membrane to be deeply injected and pigmented, especially along the line of the ureter, so that there was a distinct elevated ridge extending like a bar underneath the mucous membrane. The orifice was oval-shaped, and the space between the lips was of an unusually dark-red colour. On the surface of the bladder there were a number of little round white pellets, which, on agitation of the fluid, floated about freely, otherwise they presented very much the appearance of miliary tubercles in the mucous membrane.

B. CHRONIC DISEASES.

1. *Septic Pyonephrosis of the Left Kidney.*

This case illustrates very well the appearances presented by the mouth of the ureter in a case of very chronic septic pyonephrosis, due originally to infection following the impaction of a phosphatic calculus in one of the kidneys. There had been repeated attacks of renal colic, sometimes with symptoms of complete obstruction to the left ureter. Each attack was preceded by a rigor and accompanied by sickness, vomiting, and profuse perspiration. The suffering was sometimes so severe as to cause the patient to at once lie down; relief was obtained by hot applications and the administration of morphine. Palpation of

Fig. XXIII.—Septic pyonephrosis.

the abdomen showed marked increase in size of the right kidney. The swelling was rounded, and extended to within 3 in. of the umbilicus, and fluctuation would be readily made out. Cystoscopic examination showed that there was no fluid, either urine or pus, escaping from the right ureter, but the duct could be easily felt on vaginal examination as a thickened cord. The orifice was distinctly retracted, almond-shaped, the lips thickened, widely parted, forming a large, gaping mouth; the mucous membrane seen between the lips, and covering the distal aspect of the ureter, was bright red, a much deeper hue than the lips of the orifice, which formed an oval ring mounted on an elevated ridge (Fig. XXIII). In cases of septic kidney the involvement of the ureter is almost a certain consequence.

2. Ascending Ureteritis (Fig. XXIV).

The case was one of subacute septic cystitis, due to mixed infection of *Bacillus coli* and *Streptococcus pyogenes*, following an old-standing stricture of the urethra, which had been dilated on many occasions, but owing to neglect on the part of the patient the treatment had not been consistently carried out. Prior to the patient coming under my care the urine was alkaline, contained pus, with large quantities of mucus and a considerable deposit of triple phosphates. When the stricture had been well dilated, and the bladder condition improved sufficiently to permit it retaining 6 oz. of fluid, a cystoscopic examination showed marked thickening of the orifice of the left ureter, with deep pigmentation of the mucous membrane, marked dilata-

Fig. XXIV.—Ascending ureteritis.

tion, and a varicose condition of the arterioles (Fig. XXIV). The mouth of the ureter was situated on a rounded eminence, and appeared as a hole punched out of very swollen and congested mucous membrane. There was marked hypertrophy of the muscular coat, shown by prominent trabeculae.

Fig. XXV (Coloured Plate) shows the appearance of a less chronic case, in which changes were not so marked. In both the septic infection extended to the corresponding kidney, and the walls of the ureter were greatly thickened and indurated.

In chronic cystitis the nature of the changes induced in the ureters depends upon the nature of the primary lesion, and the mode in which the secondary infection takes place. In those instances where the infective material gains access along the surface of the mucous membrane, the inflammatory changes induce early impediment to the escape of urine through the orifice. Consequently, dilatation of the ureter in such cases is the first, the most common, and most marked change. When, on the other hand, the infection is by the lymphatic channels, the lowermost limits of the ureter may almost entirely escape, and the most marked morbid changes consist in fibrous induration of the submucous connective tissue, the wall of the ureter being thereby much thickened, while until late in the case the lumen is not materially altered. The circumstance that the interparietal portion of the duct is free from disease accounts

Fig. XXVI.—Ascending ureteritis, pinhead opening.

for the absence of dilatation, and the valvular action of the ureter is maintained. In septic cystitis, when the infection is conveyed along the surface of the mucous membrane, the lips of the orifice are greatly thickened, congested, and may even be ulcerated. Frequent and spasmodic contraction of the bladder, the resistance offered to the escape of urine from the duct consequent upon the

changes I have just described, cause protrusion of the ureteral orifices into the bladder, and gradual separation of the lips, so that the valve action of the mouth of the ureter is ultimately lost. The physiological barrier to septic infection is removed, the door being open for infection of the ureters and kidneys. When the infective process proceeds through the lymphatic channels, the walls of the ureter become sclerosed, the mucous membrane being enveloped in a dense connective tissue sheath of considerable thickness, but, while the duct for a considerable distance above the level of the bladder may be converted into a dense cord, the lumen may remain undiminished, and the mucous membrane may not be seriously involved in the infective processes. Consequently, for a considerable period the kidney may escape, but sooner or later the dense cicatricial tissue contracts, and more or less impediment may be offered to the passage of urine. In these old-standing cases the ureter orifice is contracted to a pin-head opening (Fig. XXVI), and the mucous membrane extending along the line of the ureter is elevated into a sharply-defined ridge of deeply-pigmented mucous membrane.

3. Renal Calculus of Old Standing.

The case was one of renal calculus, with symptoms extending over a period of ten years. They consisted of severe attacks of renal colic with occasional haematuria and almost persistent presence of pus in the urine. The quantity of pus varied from time to time, and was always associated with an acid condition of the urine. Although frequently urged to have an operation performed, the patient refused, until he was compelled to submit by reason of the frequency of the attacks and the increased severity of the pain. Haematuria was by no means a marked symptom. It occurred seldom, and was never severe. On cystoscopic examination the orifice of the left ureter was seen to be greatly dilated (Fig. XXVII), and the margins thickened and of a dark purple colour. The mucous membrane of the bladder in other parts was little altered, but around the orifice of the left ureter it was deeply injected, and one or two of the blood vessels in close proximity to the orifice were distended. The mouth of the right ureter was normal in appearance.

Fig. XXVII.—Renal calculus in ureter.

4. Calculus Impacted in the Left Ureter.

The interest in this case was that while it was under observation the stone was passed.

The history was that of repeated attacks of renal colic with the passage of gravel on several occasions. On September 9th, 1905, an x-ray photograph was taken, when the shadow of a stone was found corresponding in position to the union of the lower and middle third of the left ureter.

On September 10th a cystoscopic examination was made, when the mucous membrane of the bladder was found to be practically normal, but, on account of the enlargement of the middle lobe of the prostate, a cystoscopic examination was difficult. The orifice of the right ureter was normal, that of the left was considerably dilated and the shape of a melon seed (Fig. XXVIII). The margins were very slightly thickened, and on the left lip the line of the mouth was obscured by a trickle of blood escaping from it, so that the whole contour was not clearly seen. The line of the ureter was very distinctly marked, and the mucous membrane covering it was pigmented and congested. On September 11th I received a letter from the patient, saying that he had passed the stone on the evening of September 10th, and that he "felt all right now," but should I wish him to do so he was willing to undergo another cystoscopic examination.

Fig. XXVIII.—Renal calculus in ureter while impacted.

A second x-ray photograph was taken on September 11th, and

failed to show the shadow seen in the first plate, proving that the shadow seen was due to the stone which had been passed. A second cystoscopic examination showed the same appearances as formerly presented, with the addition that close to the middle of the upper lip there was a small submucous haemorrhage, and a similar spot was seen on the margin of the lower lip, but no blood was found escaping from the orifice. The stone passed was the size of a horse bean, but very irregular in shape, and distinctly nodulated. It is worthy of remark that the passage of this concretion caused no

Fig. XXIX.—Renal calculus in ureter, passed.

laceration of the lips of the ureter or other injury recognizable with the cystoscope further than the minute submucous haemorrhages just described (Fig. XXIX).

CONCLUSIONS.

It is impossible with our present knowledge to dogmatize, and the conclusions which I have come to may be either confirmed or contradicted by a larger experience. I think, however, it is well to state as clearly as possible the opinions I have formed after a somewhat lengthy series of investigations. They may form a foundation for others to build upon.

1. When one orifice is altered and the other normal, the renal lesion is on the side of the abnormal ureter, and the character of the deformity of the orifice may indicate the nature of the renal lesion.

2. Active congestion and swelling of the mucous membrane in the neighbourhood of the orifice and along the line of the ureter (Fig. XIX), also swelling and pointing of the lips, denote acute irritation of the pelvis or parenchyma of the corresponding kidney.

3. A dilated orifice, the lips being sharp and clearly defined, the mucous membrane between the lips acutely congested, while the colour of that of the bladder is little changed, denotes recent and acute inflammation or mechanical irritation in the corresponding pelvis (Fig. XX, etc.).

4. A dilated orifice, the lips being thickened and only slightly rounded, the mucous membrane between the lips of a dark-red colour, while the mucous membrane of the bladder is deeply injected and pigmented, especially along the line of the ureter, denotes descending ureteritis with dilatation (Figs. XXI, XXII, and XXIII).

5. An elongated or dilated mouth with hyperaemia of the lips indicate acute inflammation of the parenchyma or recent distension of the renal pelvis.

6. A punched-out orifice, marked thickening of the lips, induration and congestion of the surrounding mucous membrane, indicate a dilated ureter with ascending ureteritis, the infective process extending along the surface by continuity (Figs. XXIV and XXV).

7. A pinhead opening on a well-defined ridge of deeply-pigmented mucous membrane denotes induration of the walls of the ureter from ureteritis, the infective material being conveyed from the primary focus in the kidney or bladder, principally through the lymphatic channels (Fig. XXVI).

8. Pinhead contraction of the mouth without other vesical changes denotes spasm induced by the presence of a rough calculus impacted in an irritable ureter.

9. Inflammation of the mouth and thickening of the lips, with inflammatory changes limited to the mucous membrane immediately surrounding the ureter orifice, denotes mechanical irritation of old standing (Fig. XXVII).

10. Simple dilatation of the mouth, without much thickening of the lips, and with congestion limited to the orifice and neighbouring parts, suggests mechanical irritation of recent origin (Figs. XXVIII and XXIX.)

I may now indicate the inferences which may be drawn from a study of the changes to be observed in the rhythm, size, form, frequency, and composition of the efflux.*

A. When the Urinary Shoots are in Distinct Jets, but are more frequent on one Side than on the Other.

1. Greater functional activity is indicated when the shoots are large, uniform in size, and regular in rhythm.

2. Undue irritation and congestion of the kidney is to be inferred when the shoots are more frequent, small in size, unequal, and irregular in rhythm.

3. Stricture, stone, or ureteritis should be suspected when the shoots are distorted in form, or irregular in amount.

4. The action of drugs, administration of fluid, excitement, pain, or sudden surprise, may cause very rapid secretion of clear urine from both orifices.

B. When the Urine does not escape in Distinct Jets.

1. Inefficient secretion, or dilatation of the ureter without paralysis of the sphincter is indicated when the urine dribbles into the bladder at intervals.

2. Destruction of the sphincter action is denoted when the urine floats into the bladder almost continuously.

C. Absence of Efflux may be noted when

1. Inhibition of the secretion is temporary only, the ureter orifice being distinct, but not gaping.

2. Absence of secreting tissue is indicated when no shoots are seen after several examinations conducted under varying circumstances; the orifice may be difficult to find, or may be wanting.

D. The Urinary Shoots contain some Morbid Material.

1. It consists of blood-stained urine.

(a) In acute inflammation of the parenchyma of the kidney marked hyperaemia (passive), the effluxes are of a dark-brown colour, and may be copious and continuous.

(b) In carcinoma or sarcoma where the tumour has originally invaded the pelvis, or has ultimately broken down and communicated with the pelvis, and in injury to the kidney large quantities of blood may escape.

(c) In ulceration of the pelvis in malignant disease and in syphilis haemorrhage may occur into the pelvis in large quantities, and blood may be expressed from the ureter by pressure with the hand over the affected kidney.

(d) In chronic interstitial nephritis, in stone, and in early tuberculous disease, occasional small quantities of blood may be seen trickling from the ureter. If the trickle of blood succeeds a urinary shoot, the bleeding is probably due to calculus impacted at the lower end of the ureter.

2. It consists of pus-stained urine.

(a) In septic pyelitis and pyonephrosis, where the kidney is still active, pus-stained urine may be ejected at regular intervals. Pus is constantly present, is intimately mixed with the urine, and flakes are not numerous.

(b) In tuberculous pyelitis the pus is not so intimately mixed with the urine, many shoots are not purulent, and pus, when present, is usually flaky.

3. Effluxes of thick fluid or semi-fluid. When the kidney is inactive, as, for example, in old-standing pyonephrosis, in malignant disease where the greater bulk of the organ has become involved in the neoplasm, in advanced tuberculous disease, and in syphilitic disease of the kidney, the parenchyma of the organ may become so destroyed, or its function impaired, so that little or no

escapes with the urinary shoot, as, for example, in a case of renal calculus where the kidney is comparatively healthy; (2) a case of ulceration of the pelvis of the left kidney, where the cavity was distended with fluid blood; (3) a case of pyelitis in which puriform urine

Fig. 9.

escaped in distinct shoots: (4) a case of chronic pyonephrosis where the function of the kidney was in abeyance; and (5) a shoot containing flaky pus, such as is seen in tuberculous pyelitis.

*The shoots were demonstrated in the projector by means of the coils (Fig. 9) described at p. 666. So as to give a clear idea of effluxes, five illustrations were given: (1) Where a small quantity of fluid blood

urine is excreted. Under such circumstances, pure blood or blood clot may be passed through the mouth of the ureter, or pus may escape in putty-like bands which issue slowly from the ureter.

When haemorrhage takes place into the pelvis of the kidney, and the blood is retained there for a considerable time, it becomes so altered that it does not mix freely with the contents of the bladder. When it passes from the ureter it may appear like a ribbon or band of melted red wax, which flows slowly from the mouth, and, after it has attained a certain length, breaks off and floats away.

ERRATUM.—In the BRITISH MEDICAL JOURNAL for March 24th, p. 665, col. 2, line 17 from bottom, the words, "includes 75 per cent. of a circle" should read, "includes 75 degrees of a circle."

THE PROGNOSIS IN POSTERIOR BASIC MENINGITIS.

By O. HILDESHEIM, M.D. OXON.,
LATE HOUSE-PHYSICIAN, HOSPITAL FOR SICK CHILDREN,
GREAT ORMOND STREET.

POSTERIOR basic meningitis is by no means so rare a disease as is generally supposed. It accounts for one-third of all the cases of meningitis which occur among the in-patients at Great Ormond Street Hospital—133 out of 396 consecutive cases. Some account of the general prognosis in this disease, therefore, may be useful, the more so since there is no illness in which it is more difficult to forecast the future. This difficulty arises in chief from the frequency with which the inflammatory exudate interferes with the circulation of the cerebro-spinal fluid, giving rise to some degree of hydrocephalus. The symptoms of concealed hydrocephalus have been described elsewhere.¹ They were divided for purposes of description into four groups.

1. Sudden death; relapse; sudden attacks of vomiting, headache, and pyrexia.
2. Convulsions; insanity.
3. Paralysis; retarded development.
4. Inability to gain flesh, headaches, incontinence, peculiarities of temper, morals, or emotions.

There are, however, certain other sequelae which cannot be ascribed to hydrocephalus, which must be considered in some detail before we proceed to a general survey of the prognosis.

Amaurosis without Optic Neuritis.

This is the most remarkable of these. Carr² does not mention this condition. Lees and Barlow³ say: "Although optic neuritis is rare yet blindness more or less complete is not uncommon," and they cite cases. Stephenson in 1902 reported from the Royal London Ophthalmic Hospital 4 cases of "fleeting amaurosis in infants presenting symptoms of meningitis."

In more than one-third of my cases the child was either blind, or its power of vision was uncertain at some period of the disease. Of these, over 60 per cent. were under 6 months at the time of onset of the disease, showing that the condition is more common in younger patients. The oldest child was 2½ years. In most instances the blindness begins insidiously. In half my cases it was not recorded until more than a month after the disease set in. Generally the condition endures for a few weeks—from three to six weeks; sometimes it lasts much longer. In 3 cases it was present ten months, one year, and two years respectively, but the following history presents my most remarkable example of the condition.

CASE X.—Aged 3 months. He "fell out of bed and injured his brain." In the next five weeks he had nine fits. A week after the fall he was found to be blind. He was an in-patient in three London hospitals. He remained blind for five years, and then sight gradually and completely returned. Two years later, at the age of 7, he began to talk. My own notes say: "Age 10. Can't walk. Legs slightly spastic. Big, healthy-looking boy. Idiot. Very hilarious and good-tempered. Vision normal." There was evidence in the old notes of an early hydrocephalus which became arrested.

Vision returns gradually. In one case two years after the illness the mother told me that, although the sight was perfect usually, yet "when the child is indisposed vision seems blurred." Stephenson gives an almost identical description. Six months after the attack "the sight seems to fade away when the baby is poorly." The fundus oculi is usually pale, but otherwise normal.

The condition is often referred to the excessive intra-cranial pressure so commonly found. The following facts militate against such a view:

1. There was other evidence of hydrocephalus in 29 per cent. of these cases, as against 36 per cent. of all my cases of posterior basic meningitis.

2. Blindness sometimes begins long before there is reason to suspect increased pressure—for example, Case IX became blind immediately after the onset of the disease. Eight months later the head was first noticed to enlarge.

3. The amaurosis disappears, although the hydrocephalus may be progressive.

4. The condition is not included in descriptions of chronic hydrocephalus.⁴

The condition is probably never permanent. In only one case, which I was unable to trace, was the amaurosis persistent when the child was last seen.

Optic Neuritis.

Optic neuritis does undoubtedly occur in posterior basic meningitis, but it is very rare. In many cases there are ill-defined appearances of some change in the fundus insufficient to warrant a diagnosis of neuritis. A necropsy of a case in which optic neuritis and retinal haemorrhages occurred is described in vol. xv of the Great Ormond Street records. It occurred in only one undoubted case in my series⁵; the boy's general condition, seven months after the attack, was steadily improving, but he remained totally blind.

Pseudo-glioma.

Lees and Barlow⁶ describe one case; another has been recently recorded by Dr. Blaber.⁷ I have two instances to add.

CASE LXXXIX.—The child was an out-patient, and the notes are scanty. About the third week, the right eye became affected. "Mr. Lister and Dr. Batten are agreed that the condition is probably a mild degree of pseudo-glioma." A year later vision is normal.

CASE XXXVIII.—Aged 1½ years. An illness which was probably posterior basic meningitis. At the age of 2 years and 3 months he was admitted for fits and blindness. Well nourished; head retracted; legs strongly adducted. Whilst in hospital hydrocephalus rapidly developed. Note by Mr. Lister: "In my opinion there is undoubted chronic iridocyclitis (pseudo-glioma)."

Deaf-mutism.

Apparently this condition is rare. Its frequency is apt to be overrated because it is quite common in certain epidemics of cerebro-spinal fever. I have found only two cases, of which one (Case XXIX, loc. cit.) developed the condition after several operations for subdural drainage.

CASE XXXIV.—Aged 2½ years. Posterior basic meningitis. At the age of 2 years and 10 months, in hospital, "comparatively well, but is not sane; her memory is gone; she does not know her people; she is dirty in her habits; her mother thinks she does not see at all. The child is an idiot." At the age of 6 years and 10 months the father writes, "I am pleased to state she enjoys the best of health, and appears to have all her faculties. I am sorry to say she cannot speak, and her hearing is very dense."

Insanity.

Insanity is not very uncommon, and it often lasts for some months. Consequently the patient is often sent from the hospital *in statu quo*, and this has given rise to the idea that the condition is not rarely permanent. I have records of three such cases where the child ultimately became quite sane. My only permanent cases are the idiot child described above (Case X), and another who had frequently recurring fits during the two years that she survived (Case XXX, loc. cit.).

General Prognosis.

Out of 133 cases in which the diagnosis of posterior basic meningitis was made, 43 were sent out of hospital alive. This does not indicate that one-third of all the cases recover, because:

1. The more severe cases are more commonly sent to hospital.

2. Mild cases are treated in the out-patient department.

3. Many of those who are sent out alive die subsequently.

Probably the survivors of the original illness number considerably more than 1 in 3, whilst in distinctly less than one-third of all cases can ultimate recovery be expected.