differences between the two may be easily enough recognized, and the intermediate forms distinguished. When the spirochaetes are numerous they are likely to be *Spirochaeta refringens*. The variation in the degree and kind of the curves (slightly waved, straight, or spiral existing on one film) proclaims the *Spirochaeta refringens*. When *Spirochaeta pallida* exists side by side or in the same specimen, the divergence of form removes all doubt as to the organism under observation. The fact that *Spirochaeta refringens* has been found in the lymphatic glands at least once should cause one to take note of the presence of this spirochaete more carefully.

Fig. XI.—Diagram of short *Spirochaeta refringens*, showing: (a) longitudinal fission; (b) spore formation; (c) transverse division by attenuation of part; (d) various forms taken by the organism. Drawn from a film stained with acetone gentian violet from a primary commissural chancre. The film also contained the *Spirochaeta pallida* in fewer numbers.

I have to thank Dr. Thomas Kay, the Visiting Surgeon, and the Resident Staff of the Glasgow Female Lock Hospital for their assistance in this investigation; and I am much indebted to Dr. Leslie Buchanan for the vast amount of time and trouble he has taken in photographing the specimens.

Since writing this paper I have read Siegel's third paper,⁴ and while I have named the bodies which I think are the progeny or the progenitors of the *Spirochaeta pallida* as *Cytorrhycles luis*, they bear a resemblance only to some of those figured by Siegel. With the subject in its present stage, it is not advisable to multiply names given to the organisms found, though already this tendency is very apparent. Schutz's paper,⁵ published in March, entirely supports the views set forth in my preliminary communication and in this paper.

REFERENCES.

- ¹ Herxheimer und Opificius, Weitere Mitteilungen über die *Spirochaeta pallida* (*Treponema Schaudinnii*). *Munch. med. Woch.*, 1906, No. 7.
- ² Rille, Ueber Spirochätenbefunde bei Syphilis. *Munch. med. Woch.*, 1905, No. 29.
- ³ Neisser, Versuche zur Uebertragung der Syphilis auf Affen. *Deutsche med. Woch.*, 1906, No. 3.
- ⁴ Siegel, Weitere Untersuchungen ueber die Aetiologie der Syphilis. *Munch. med. Woch.*, 1906, No. 2.
- ⁵ Schutz, Mitteilungen ueber *Spirochaeta pallida* (Schaudinn) und *Cytorrhycles* (Siegel). *Munch. med. Woch.*, 1906, No. 12.

A FATAL CASE OF ACUTE CARDIAC BERI-BERI.

By HAMILTON WRIGHT, M.D.,

LATE BERI-BERI RESEARCH COMMISSIONER, LONDON SCHOOL OF TROPICAL MEDICINE; PATHOLOGIST, WEST LONDON HOSPITAL AND POST-GRADUATE COLLEGE.

THE case of acute beri-beri here recorded is one of several that I have had the opportunity of studying through the courtesy of the Committee of the London School of Tropical Medicine. It is the only case that proved fatal. The result of the *post-mortem* examination of this single case, however, largely confirms my previous observations on this disease. The bacterial investigation of the case has not been finished, but enough has been done to warrant pushing the investigation to a conclusion. For this purpose I have handed over to Dr. Dudgeon of St. Thomas's the cultures, not only of a suspicious bacillus found in this case, but of a similar bacillus recovered from the stools of other acute beri-berics. Dr. Dudgeon will report in due course. In the meantime it seems wise to publish the case itself and my own findings in it.

A. A., age 37, a seaman, was admitted to the Branch Hospital of the Seaman's Hospital Society (Dr. Duncan's Clinic) on November 23rd, 1905.

History and Condition on Admission.

Dr. Stanton, House Physician, noted as follows: The patient, a native of Calcutta, made his last voyage from Calcutta on the ss. *Sumatra*, and arrived in London about November 2nd, 1905. The ship went to Antwerp and returned to the Royal Albert Docks on November 22nd. His present illness began on his first arrival in London about November 2nd. There was cough, with pain in the deep epigastric region which he attributed to the cough. He had no appetite and had marked constipation. In a few days his legs felt weak and began to swell. Both the weakness and swelling gradually increased. No history of other illnesses can be obtained. He had not had any venereal disease. He said that ten others occupied the same quarters as himself, and that as far as he knew none of them had contracted his disease. The physical examination was made by Dr. Hamilton Wright and Dr. Stanton. The temperature was 97.4° F., the pulse at rest 112. Respirations 32. The patient was much depressed; brandy had been administered for threatened heart failure. The pulse was small, weak but fairly regular. There was marked dyspnoea. The lips were cyanosed. The apex beat was beneath the sixth rib in the left nipple line. The precordial beat was somewhat diffuse. The first sound heard at the apex was weak and the diastolic pause decidedly shortened. Occasionally a soft systolic murmur was heard at the apex. The pulmonary and aortic second sounds were frequently reduplicated. The right cardiac dullness extended to the left border of the sternum. There was marked pulsation in the cervical veins. The general character of the cardiac rhythm was embryonic. Large and small moist râles were heard in all parts of the lungs, and there were signs of fluid in the posterior pleurae. The abdomen was distended, and appeared to obtain a considerable amount of fluid. There was marked oedema of the legs, thighs, and sacrum. The patient was slightly dulled mentally from the effects of the brandy administered, and no certain facts could be determined in regard to his sensori-motor condition. The knee-jerks were absent; the superficial reflexes normal. The myotatic irritability of the muscles of the legs and thighs was excessive to direct mechanical stimulation.

The urine for the first twenty-four hours measured 10 oz., specific gravity, 1024; acid, no traces of albumen. The faeces contained some eggs of *A. lumbricoides*, also eggs of *T. dispar*. These disappeared on the exhibition of drugs.

November 24th. A pin can be driven through the skin of the patient's legs without his noticing it if his attention is engaged by conversation. When distraction ceases he correctly locates the pin, and complains of pain. It would seem that dolorific pain is at least delayed. Flexion of the ankles is slow; also extension of the toes. It is difficult to decide whether or not this is due to the massive oedema. The cardiac condition is as first noted. The patient had been given a mixture containing tr. digitalis, spt. ether. nit., and aq. chlorof. every four hours.

On November 27th the patient was as when last noted; 5 gr. of santalin was given.

On November 29th he was still in the same condition as when admitted; sodium sulphate was given.

On December 5th it was noted that many remedies had been exhibited for the dyspnoea and cardiac distress without avail. The oedema had increased, and extended to the trunk. The abdominal wall was markedly oedematous as high as the umbilicus. The scrotum was massive with oedema. The râles in the lungs had cleared up. The sensori-motor condition was not clear; there was certainly no disturbance of either function in the trunk or upper limbs. The legs, however, were apparently dulled to painful and tactile sensations. This might, however, have been due to the oedema. The knee-jerks had been tested nearly every day, and had not been elicited. All other reflexes were normal. Mentally he was clear, but talkative and difficult to keep to the point.

On December 11th, though the lungs were practically clear of all râles, he expectorated a small quantity of blood-stained mucus. The oedema had not subsided. The cardiac condition was as on admission, except that the embryonic character of the rhythm was more pronounced. There was no certain sensori-motor disturbance, except that painful stimulation appeared to be delayed as high as the knees. The reflexes were as on admission. Flexion of the ankles and extension of the legs was certainly slower than the opposite movements; but certain weakness was somewhat doubtful.

On December 18th it was possible to get a certain response to sensory stimulation. Tactile sensation was diminished as high as the knees. There was a slight weakness of the extensors of the legs and of the flexors of the ankles. The cardiac condition was as on admission, except that the irregularity was greater.

On December 24th the condition was unchanged, except that the urine was diminishing in quantity. The patient gradually failed. The oedema did not clear up, and he grew weaker in the legs. It was not possible to determine if the sensory disturbance had extended, as the patient showed considerable irritation and uncertainty on examination. The cardiac sounds and rhythm became more irregular. At 9.30 a.m. on January 1st, 1906, he raised himself in bed, stared round the ward, gasped a few times, and fell back dead.

Dr. Wise made a careful attempt to cultivate organisms from the blood of this case, but failed to do so.

Necropsy.

Necropsy at 3 p.m. Weather cool. Body of a well-developed Indian. There is marked general anasarca. All superficial veins bleed freely on section. Rigor mortis is partial. The eyes are slightly bloodshot. Pupils are equal and regular. There are a few traumatic scars on the shins and knees. The abdominal wall is markedly oedematous. The abdominal muscles are normal-looking. The spleen is rather small, dark chocolate in colour, firm, and cuts with only slight resistance.

The liver is reduced in size. It is firm, and the anterior border is sharp. Just above the gall bladder the edge is indurated. The pulp is congested, and there is a moderate nutmeging. There is a slight grating when the pulp is incised. The walls of the gall bladder are slightly thickened. The ducts are free, and the bile runs freely.

The kidneys are healthy-looking. The capsules peel readily, leaving a smooth reddened surface. The cortices are normal-looking. The cut surfaces retract slightly. There is perhaps a slight cloudy swelling.

The entire mucosa of the stomach is slightly congested. Here and there are patches and rings of deep scarlet congestion, made up of distinct points of injection. These patches and rings are more numerous and closer together in the great curvature and about the pyloric ring. The mucosa has lost its transparent look and is opaque. There are no erosions or ulcerations. The duodenum and upper few inches of the jejunum are as the stomach. The rest of the small gut is slightly congested, and there are some haemorrhagic injections of the valvulae conniventes. The entire large gut is normal. No worms seen.

Fig. 1.—Gastric mucosa in acute pernicious beri-beri. Case 6 of publication 4.

Both pleurae contain a moderate amount of clear yellowish fluid. The right lung is anthracosed. The lobes are glued together by slight and recent adhesions. The apex and anterior border are acutely emphysematous. The cut surfaces exude on pressure a slight amount of blood-stained mucus. The left lung is the same as the right.

The pericardium contains a moderate amount of clear yellowish fluid. There is a slight general cloudiness of the visceral pericardium, most marked along the lines of the vessels. There are also in this layer numerous small and large patches of petechiae.

The right heart is enormously dilated, and full of dark semi-fluid *post mortem* clot. The muscle tissue looks coarse-grained and spongy. It does not look grossly fatty. The right auriculo-ventricular ring is stretched. The left ventricle is slightly dilated. The wall is 13 mm. thick. The cut surfaces are granular, pale, and spongy-looking. No gross fatty change to be made out. The coronary arteries are slightly atheromatous at their origins. The large vessels and all cardiac valves are normal.

The brain exhibits no gross lesion.

The muscles of the lower limbs are moderately oedematous. The subcutaneous tissues of the lower limbs, abdomen, scrotum, and back are markedly oedematous.

Microscopical Examination.

Several portions of the stomach, duodenum, and other parts of the small and large gut were microscopically examined. The large intestine was normal. The mucosa of all parts of the stomach and duodenum exhibited a more or less marked necrosis (see Fig. 1, which shows this necrosis in a case studied in the Malay Peninsula; and Fig. 2, which is of the pyloric mucosa of this case). Only the cells of the bodies of the glands were natural-looking. A small amount of fibrin was shown by Weigert's method. All the vessels were over-full; and in the congested patches, though there was no actual extravasation, the vessels were nevertheless distended to bursting point. The necrosis of the mucosa in the congested areas was not more marked than in the intercongested patches. This would seem to indicate that the necrosis of the mucosa was not due to the congestion. All sections showed a certain amount of superficial sloughing of the mucosa. In the duodenum the villi were often seen almost detached; in the stomach the sloughing was unevenly distributed. Most of the sloughs contained a bacillus which corresponded with the bacillus already described by me as having been found in other cases of

Fig. 2.—Gastric mucosa of this case.

acute beri-beri. Occasionally this bacillus was seen in the deeper layers of the mucosa, but never below the necrotic line. It was not found in any other part of the gut of this case, or in the tissue of the other organs. The bacillus measures 4 to $9 \mu \times 1.05 \mu$. It is Gram-positive, and stains readily by the ordinary basics. (See Fig. 3, which shows this bacillus in the mucosa of a Malayan case; and Fig. 4, which shows the bacillus recovered from the case presented here.) More of this organism later.

The cerebral and cerebellar cortical cells were normal. The cells of the sensori-motor cranial nuclei were likewise normal, except those of the vagi. In all levels of the ambiguous nuclei the great majority of the cells showed a marked chromatolysis and dislocation of nuclei. In the median levels not a normal cell was to be seen. The group of cells just anterior to the hypoglossal nuclei were similarly changed. The spinal cord was not removed, so that I cannot state the condition of its motor cells.

Several strands of nerve fibres dissected from the right cardiac auricle exhibited the effects of the beri-beri toxin. In a few fibres there were truly Wallerian signs of degeneration. But the majority of the fibres were in what I have previously classified as the first and second stages of change—that is, the medulla at the nodes and at points between the nodes was blackened by osmic acid, but had not broken into globules. The main trunks of the cervical and thoracic cardiac branches of the vagi were healthy, so that in this, as in other cases of acute beri-beri which I have examined, the change in the nerve fibres is purely peripheral.

Both stellate ganglia contained a minority of cells which showed dislocation of nuclei. A few cells were seen which also exhibited some chromatolysis. This would appear to indicate that a minority of the post-ganglionic accelerator cardiac fibres were poisoned by the beri-beric toxin. I did not determine the state of the post-ganglionic accelerator fibres themselves, but the "axonal reaction" in the stellate cells would appear to indicate that they, as well as the inhibitory fibres, had been involved by the beri-beric virus. On these findings the cardiac irregularity, distension, and final failure are explainable.

Only a few poisoned fibres were found in the twigs to the anterior tibial and peroneal muscles. The cutaneous branches of the musculo-cutaneous in the legs also showed a few poisoned fibres. There were not enough of them, however, to warrant a view that there was large disturbance of the sensori-motor functions of the lower limbs. It will be remembered that this was a doubtful point in the clinical examination.

Fig. 3.—Bacilli in mucosa of Fig. 1.

The central cells were stained by Nissl's method; the nerve fibres by Marchi's and Busch's methods.

The cardiac muscle stained by Marchi's method, and with Soudan III exhibits a marked fatty degeneration. The degeneration was not uniformly distributed, except in the papillary muscle. Rather it occurred in small, well-marked patches, separated by a few fibres which showed only a slight fatty degeneration. The right heart appeared to be more extensively and more severely involved than the left.

There was a moderate induration of the liver, with some fatty infiltration. The vessels were full, and occasionally small extravasations were met with. Thick frozen sections showed more free blood. But by this method blood is readily drawn from the vessels, and so spread as to resemble extravasations. The spleen exhibited nothing of note. The kidneys were over-full of blood, but otherwise normal.

BACTERIOLOGY.

Broth, ordinary agar, and glucose agar, were inoculated from the auricular blood, from the liver, and from the contents of the stomach and duodenum; also from scrapings of the mucosa of the latter parts. The cardiac blood proved to be sterile, and no growth occurred in the media inoculated from the liver.

All of the media inoculated from the contents and scrapings of the stomach and duodenum showed growths by the end of twenty-four hours. Plates were made from all surface colonies, but no organism was seen which corresponded with that found in the mucosa of the stomach and duodenum.

In the glucose agar stabs, however, there was found in almost pure culture a bacillus which corresponded morphologically and in staining reaction with that found in the gastric and duodenal mucosa. It was recovered in pure culture in Bulloch's apparatus and in Buchner's tubes. By these methods it grew slowly and aberrantly; but when stabbed into glucose agar it extended rapidly, first, in discrete colonies which soon coalesced, and then by diffusion of products clouded the whole column of media.

The bacillus at twenty-four hours ranged from 4 to 8 or 9 μ in length, and was 1.05 μ in breadth. The ends are almost square, and the sides regular but slightly curved. Older cultures were longer, stouter, and tended to become clubbed. By both Weigert-Gram and Loeffler the staining was irregular, many of the organisms resembling the granular forms of the Klebs-Loeffler bacillus.

INOCULATION EXPERIMENTS.

A rat proved to be refractory to 15 c.cm. of a broth emulsion of a ten-day-old culture subcutaneously injected. A monkey subcutaneously injected with two similar doses showed signs of paralysis of the hind limbs; but it developed a small septic abrasion in the thigh, and so had to be excluded. A guinea-pig injected subcutaneously with 15 c.cm. of a broth emulsion on January 25th and with a similar dose on February 1st exhibited first a general weakness and then paresis of the hind limbs on February 12th. On February 16th the paralysis of the hind legs was more pronounced, and on February 17th the animal died. All cultures from blood and tissues

Fig. 4.—Bacilli. Ten-day-old culture from duodenum of present case.

proved sterile. There was no lesion at the site of inoculation. No pathological change was found in any of the tissues, except in the nerve twigs to the flexors of the hind ankles, which contained a few changed fibres—about 1 to 4 or 5. Degenerated fibres are not unusual in these animals, so that this observation cannot be regarded as satisfactory.

Not being in a position to carry these experiments further Dr. Dudgeon has consented to do so, and all cultures have been handed to him. I am indebted to Dr. G. C. Low and Dr. Wenyon for having inoculated the above animals for me.

CONCLUSION.

All that need be said in conclusion is that here we had a case of acute beri-beri, dying in the seventh week of the disease from cardiac failure, and exhibiting the lesions which were constantly found during my studies of this disease in the Far East. The train of events during life were an acute onset with complaint of gastro-intestinal symptoms, followed or accompanied by other symptoms that marked the effect of the beri-beric virus on the cardiac and vaso-motor nervous systems; then death from cardiac failure.

PREVIOUS PAPERS BY THE AUTHOR.

¹ Changes in the Neuronal Centres in the Beri-beric Neuritis, *BRITISH MEDICAL JOURNAL*, June 29th, 1901. ² *The Etiology and Pathology of Beri-beri*, Churchill, London, May, 1902. ³ Beri-beri in Monkeys, *Brain*, vol. xxvi, 1903. ⁴ *The Pathology and Classification of Beri-beri*, London, 1903. ⁵ Result of Preventive Measures against Beri-beri, *Journal of Hygiene*, April, 1905. ⁶ The Pathology of Beri-beri, *Review of Neurology and Psychiatry*, September, 1905. ⁷ The Cause, Course, and Treatment of Beri-beri, *American Medicine*, October, 1905.