these facts: that the lymph paths of the parietal, peritoneal, and pleural serosae are in continuous communication with those of the diaphragmatic serosae; that there exist in the diaphragm numerous perforating lymph vessels, and that these pass through the diaphragm not only in the direction from peritoneum to pleura, but also, contrariwise, from pleura to peritoneum. The anatomical disposition is thus given for a spread of infection in the direction illustrated by the present case. On the other hand, the assumption of Burkhardt that the bacteria pass through the membranes and muscle by a process of active growth, and not by passive transportation, cannot be disproven. Probably the metastasis occurs in both ways.

REFERENCES.

- ¹ Gruenisen, Ueber die subphrenischen Abscesse mit Bericht ueber 60 operierte Faelle, *Arch. f. klin. Chir.*, 1903, Band lxx, Heft 1.
² Perutz, Der subphrenische Abscess, *Centralbl. f. d. Grenzgebiete d. Med. u. Chir.*, 1905, Nos. 4 to 9.
³ Stephen Paget, *The Surgery of the Chest*.
⁴ Kuettner, Die perforierenden Lymphgefasse des Zwerchfells und ihre pathologische Bedeutung, *Beitr. z. klin. Chir.*, 1903, Band xl, Heft 1.

AN IMPROVED METHOD OF PERFORMING THE "PANCREATIC" REACTION IN THE URINE.

By P. J. CAMMIDGE, M.B.LOND., D.P.H.CAMB.

IN March, 1904, I described a new method of examining the urine in suspected cases of pancreatic disease, the results of which, when considered in conjunction with the clinical symptoms, had been found by Mr. Mayo Robson and myself to be of considerable assistance in diagnosis.¹ Since then numerous experiments have been made with a view to freeing the test of disturbing factors which, either from errors in manipulation or from inherent defects in the method, were liable to obscure the issue or render its interpretation at times a matter of difficulty, especially in inexperienced hands. Before describing the improved method, I should like to correct an erroneous impression which some appear to have received from my original paper. It was not my intention to suggest that the presence of glycerine in the urine had been proved in pancreatic disease, nor did I wish it to be thought that the methods of examination I described were tests for that substance; the glycerine theory was merely mentioned to make clear the genesis of the reaction finally described.

Examination of the phenylhydrazin precipitate derived from the urine in cases of pancreatic inflammation, after treatment with hydrochloric acid, showed that it consisted of a phenylhydrazin compound of glycuronic acid and the osazone of a sugar. Investigation of the precipitate obtained from the urine after treatment with mercuric chloride indicated that it consisted entirely, or almost entirely, of a glycuronic acid compound of phenylhydrazin, so that the difference noticed between the original *a* and *b* reactions in characteristic cases of pancreatic disease was probably dependent on some substance other than glycuronic acid. If, therefore, the glycuronic acid could be removed previous to employing the phenylhydrazin test, the result might be expected to be more definite and the necessity for two comparative tests be done away with. This was found to be possible by taking advantage of the fact that glycuronic acid is precipitated by basic lead acetate from acid solutions, while the sugars are only thrown out when the reaction of the fluid is alkaline. The details of the improved method based upon this fact are as follow:

A specimen of the twenty-four hours' urine, or of the mixed morning and evening secretions, is filtered several times through the same filter paper and examined for albumen, sugar, bile, urobilin, and indican. A quantitative estimation of the chlorides, phosphates, and urea is also made and the centrifugalized deposit from the urine examined microscopically for calcium oxalate crystals. If the urine is found to be free from sugar and albumen, and of an acid reaction, 1 c.cm. of strong hydrochloric acid (specific gravity 1.16) is mixed with 20 c.cm. of the clear filtrate, and the mixture gently boiled on the sand-bath in a small flask, having a long-stemmed funnel in the neck to act as a condenser. After ten minutes' boiling the flask is well cooled in a stream of water, and the contents made up to 20 c.cm. with cold distilled water. The excess of acid present is neutralized

by slowly adding 4 grams of lead carbonate. After standing for a few minutes to allow of the completion of the reaction, the flask is again cooled in running water and the contents filtered through a well-moistened, close-grained filter-paper until a perfectly clear filtrate is secured. The filtrate is then well shaken with four grams of powdered tribasic lead acetate and the resulting precipitate removed by filtration, an absolutely clear filtrate being obtained by repeating the filtration several times if necessary. Since the large amount of lead now in solution would interfere with the subsequent steps of the experiment, it is removed either by treatment with a stream of sulphuretted hydrogen or, what I have found to be equally satisfactory and less disagreeable, by precipitating the lead as a sulphate. For this purpose the clear filtrate is well shaken with 2 grams of finely-powdered sodium sulphate, the mixture heated to the boiling point, then cooled to as low a temperature as possible in a stream of cold water, and the white precipitate removed by careful filtration; 10 c.cm. of the perfectly clear transparent filtrate is made up to 18 c.cm. with distilled water and added to 0.8 gram of phenylhydrazin hydrochlorate, 2 grams of powdered sodium acetate, and 1 c.cm. of 50 per cent. acetic acid contained in a small flask fitted with a funnel condenser. The mixture is boiled on a sand-bath for ten minutes, and then filtered hot through a filter paper moistened with hot water into a test tube provided with a 15 c.cm. mark. Should the filtrate fail to reach the mark, it is made up to 15 c.cm. with hot distilled water. In well-marked cases of pancreatic inflammation a light-yellow, flocculent precipitate should form in a few hours, but it may be necessary to leave the preparation to stand overnight before a deposit occurs. Under the microscope the precipitate is seen to consist of long, light-yellow, flexible, hair-like crystals, arranged in sheaves which, when irrigated with

Crystals obtained from the urine by the improved method in a case of pancreatitis in which there was a gall stone in the pancreatic portion of the common duct.

33 per cent. sulphuric acid, melt away and disappear in ten to fifteen seconds after the acid first touches them. The precipitate should always be examined microscopically, as it may be difficult to determine the characters of a small deposit with the naked eye, and so cases giving only a slight reaction may be overlooked. To exclude traces of sugar, undetected by the preliminary reduction tests, a control experiment is carried out by treating 20 c.cm. of the urine in the same way as in the test described excepting for the addition of the hydrochloric acid.

The urine employed for the experiment should be fresh, and not have undergone fermentative changes. If alkaline in reaction it should be made acid with hydrochloric acid before the test is commenced; any glucose that may be present should be removed by fermentation after the urine has been boiled with the acid, and the excess neutralized. The administration of calcium chloride has been found to interfere with the success of the reaction.

I have arranged the cases examined by this improved method in seven groups, corresponding to the clinical

diagnosis and the condition found at operation. For this information I am indebted to those who kindly supplied me with particulars of the cases I have examined for them in my laboratory, and particularly to Mr. Mayo Robson, who has operated on many of the cases included in Groups 1, 2, and 4. The urines here dealt with form a consecutive series of 100 specimens, derived from 94 persons. All have been examined under the same conditions by the method I have described. From 29 specimens a more or less abundant deposit of crystals was obtained, but in 71 no crystalline deposit could be found even with the aid of the microscope. In 9 cases (forming Group 1) a stone was found in the pancreatic portion of the common duct at operation, and the pancreas was said to show evidence of disease. In 19 (Group 2), although no stone was found at the time of operation, there was said to be enlargement and hardening of the head of the pancreas. I found a gall stone in a specimen of faeces sent to me from one of these cases, and in another case included in this group I am informed that biliary calculi had been discovered in the stools shortly before I was asked to examine the urine. No reaction was obtained in 4 cases diagnosed as cancer of the pancreas (Group 3). One case in which the clinical symptoms and after-history of the case pointed to malignant disease of the pancreas yielded an abundant deposit. A preparation made from the urine of this case by the original α method showed a mixture of fine crystals dissolving in 33 per cent. sulphuric acid in fifteen to twenty seconds, and many coarse, broad crystals that took five to six minutes to disappear. The urine was also found to contain a very large amount of bile and urobilin, and the excretion of chlorides was much diminished, while the urea and phosphates were approximately normal. In addition, the faeces when examined by the method I have recently described² were found to be acid in reaction, to contain a large amount of undigested muscle fibre, and to yield not a trace of stercobilin. An estimation of the fats showed a total of 64.7 per cent., of which 56.3 per cent. was "neutral fat" and only 8.4 "fatty acid." So that the laboratory results, taken as a whole, confirmed the clinical diagnosis and pointed to cancer of the head of the pancreas, with complete obstruction of the common duct and a considerable degree of secondary inflammation. Of the remaining 67 specimens which gave no reaction, 9 were from cases of jaundice in which a calculus was found at operation in the gall bladder or in the common duct above the pancreatic portion (Group 4). In all these cases the pancreas was said to be normal in appearance and consistency. Fifty-two were examined as controls. Of these, 30 were from patients suffering from diseases in which there was no reason to expect that the pancreas was involved (Group 5). They included cases of cancer of the stomach, colon, rectum, or liver, gastric ulcer, duodenal ulcer, gastritis, colitis, appendicitis, tuberculosis of the intestine, intestinal obstruction, cirrhosis of the liver, hepatic abscess, nephritis, floating kidney, tuberculosis of the kidney, cystitis, and Addison's disease. Twenty-two were from healthy individuals (Group 6). I have placed six cases in Group 7 which were under the care of Mr. Mayo Robson, and in which I had the opportunity of examining the urine one to two weeks after a cholecystenterostomy had been done for the relief of pancreatitis. All had given a well-marked reaction when the urine was examined shortly before the operation, and are included in Groups 1 and 2, but when the second examination was made no reaction could be obtained.

I am not at present prepared to contend that at all times and under all conditions the results given by this improved method of examining the urine are indicative of the presence or absence of pancreatic inflammation, as my experience of it is not as yet sufficiently great, but I think that the evidence available suggests that it may be of considerable assistance in diagnosis, and that by means of this and other methods of investigation I have described, the 50 per cent. of cases of pancreatic disease now said to be only recognized *post mortem* might be reduced to a much more creditable figure.*

My experience suggests that it is always advisable to

* Since writing this paper I have examined 95 specimens of urine, most of which were sent to me for the diagnosis of pancreatic disease, by this method, and in 54, where the after-history has been obtainable, my conclusions, based upon the examination of the excreta, have been confirmed. In the remaining 29 no reply has been received, or the case is too recent to allow of a definite opinion being formed.

control the urine examination by an investigation of the faeces, for if the results agree, the chance of a mistaken opinion is considerably reduced and is probably small. The examination of the faeces is particularly useful in cancer cases, as the instance quoted in a former part of the paper clearly shows. It is of the utmost importance that malignant and inflammatory conditions of the pancreas should be clearly differentiated. Clinically it is often a matter of great difficulty; but if the results of the examination of the urine and faeces, carefully and conscientiously performed, are considered in conjunction with the history and symptoms, the chance of an erroneous diagnosis is materially reduced. I should, however, like to emphasize the fact that the laboratory investigations require a considerable amount of care, and that it is necessary to weigh all the evidence obtainable in forming a conclusion. The "pancreatic" reaction is only one factor that has to be taken into account in making a diagnosis in suspected cases of pancreatic disease or of jaundice in which it is sought to determine whether there is a gross obstruction to the free flow of bile into the intestine, and whether this obstruction when present is of a simple or malignant nature.

REFERENCES.

- ¹ *Lancet*, March 19th, 1904. ² *BRITISH MEDICAL JOURNAL*, October 28th, 1905.

EXTROVERSION OF THE BLADDER :

ITS TREATMENT BY EXTRAPERITONEAL IMPLANTATION OF THE URETERS INTO THE RECTUM.

By C. J. BOND, F.R.C.S.,

SURGEON TO THE LEICESTER INFIRMARY.

I WISH to add another (hitherto unpublished) case of extraperitoneal rectal implantation of ureters for ectopia vesicae to the eight cases recorded by Dr. H. Simpson Newland in his interesting paper in the *BRITISH MEDICAL JOURNAL* of April 28th.

As Dr. Lendon says in his description of his original operation in an article in the same issue of the *JOURNAL*, it seems probable that extraperitoneal implantation of the ureters into the rectum will eventually become the method of the future in dealing with these distressing cases. And, indeed, any one who is familiar with the repeated operations, the imperfect immediate results, and the very unsatisfactory ultimate state of the patients when treated by the older plastic methods, will certainly endorse this opinion.

Having tried Maydl's intraperitoneal method unsuccessfully in a child of 8 weeks, in January, 1903, I operated on May 1st, 1903, on a healthy lad, aged 17, the subject of well-marked ectopia vesicae. After isolating each ureter with a small button of vesical mucous membrane, surrounding its orifice, as in Dr. Lendon's method, I separated the lower attachment of the vesical wall and inserted the mucous flaps containing the ureter endings into the wall of the rectum extraperitoneally by two separate incisions, one on either side of the gut. The lad is now, three years later, a fairly healthy young man of 20. He has practically no difficulty with the urine, which is voided from the bowel four or five times in the 24 hours. Although his condition is so far satisfactory, there is no doubt that he passed, shortly after the operation, through a serious attack of pyelitis, first of one kidney and then of the other, and even now has occasional attacks of pain in the left kidney, with purulent urine.

It appears probable that every patient whose ureters are made to open into the colon or rectum undergoes some degree of ascending infection of those canals, and bearing in mind the presence of ascending currents in the ureters among other mucous canals, under certain conditions, as described in my Address on Surgery at the Leicester meeting of the Association last July,¹ it is difficult to see how things could be otherwise. While under these new conditions the ureters must open into a septic cavity, yet the preservation of the normal valvular opening at the ureteral papillae, and hence the avoidance of any cicatrization of the outlet must be our chief safeguard in preventing stagnation of the contents of the ureter, the most potent contributory factor in the production of ascending infection. It will be interesting to learn whether the transplantation of the bladder wall and