For the fainting attacks it is well for the parents to keep at hand a rapidly-acting diffusible stimulant, such as is prescribed for patients who live in dread of an attack of angina pectoris. Sal volatile, ether, and spirits of chloroform makes a useful combination. A warm mustard bath is valuable for an infant in this state of collapse, and oxygen is also of assistance. When the heart failure is more gradual and a result of overdistension of the right side of the heart the application of leeches over the liver brings relief.

AORTIC INSUFFICIENCY.

By SEYMOUR TAYLOR, M.D., F.R.C.P.,

PHYSICIAN TO THE WEST LONDON HOSPITAL.

DR. MITCHELL BRUCE'S clinical lecture, published in the *BRITISH MEDICAL JOURNAL* of January 6th last, will amply repay careful study. There is one point, however, about which, as he admits, "there is a division of opinion," since it is "maintained by some authorities on diseases of the heart that digitalis should not be given in aortic incompetence," but he justifies the use of digitalis, because his patient, who had "failure of the heart in association with aortic incompetence, was beneficially, as well as safely, treated with . . . digitalis."

I am in thorough accord with so distinguished an authority when he makes his deduction from clinical experience. The theories which are evolved from laboratory experiment are not always justified or proved by bedside observations. The records of a physiological observation and the evidence of a sphygmograph or cardiograph may not repeat themselves, as we expect they should, when we arrive at the bedside.

There would appear to be a tendency in laboratory experiment to record results with a somewhat biased mind. A sphygmographic curve intended to show the effect of this or that drug may mislead, and I think often has done so. I have worked with this instrument for a number of years, but I cannot satisfy myself as to the clinical accuracy of its statements. Hence the more patients I see the more I am convinced that clinical knowledge and experience constitute the most valuable form of medical observation. I would, therefore, meet Dr. Mitchell Bruce on the same clinical grounds, by recording my experience of the use of digitalis in aortic regurgitation.

It is not a little singular that some of the most eminent authorities on cardiac affections, who advocate the use of digitalis in this grave valvular affection, do so with qualifications and reserve. They, with some notable exceptions, appear to be somewhat lukewarm in its praises, or at least to resort to some saving clause, as though they feared that their writings and opinions should be falsified by the test of accumulated experience. Dr. Oliver, when advocating the use of digitalis in this disease, admits that the cases must be carefully selected. Dr. Barrs says that digitalis may aggravate symptoms, and in some cases, although much temporary relief has been afforded, sudden death had followed its use. But his opinion is decidedly in favour of the drug when there are signs of ventricular failure. Sir William Broadbent, as becomes a physician with his profound clinical knowledge, is cautious when speaking of the use of this drug. He points out the advantage of digitalis when dropsy supervenes, but shows that if its exhibition be persisted in after this, it is not uncommon for patients to die suddenly, sometimes in bed, more frequently when they have begun to get up. He says, "There are grounds for suspicion that digitalis contributes to this. In the absence of mitral symptoms, it is rarely that digitalis is called for in aortic incompetence, or is of service, and it may undoubtedly do harm."

Probably the controversy may never be settled in our time. Digitalis will have its advocates and its opponents. Many of the former will say that the latter are influenced by the teaching of pharmacology; and the latter will retort that the former are imbued with the experiments in physiology and physics. It may be as well, therefore, if short records of a few cases which have come under my notice should be published:

1. Kate F., aged 35, sempstress, was under my care as an out-patient for fourteen years. She suffered from aortic regur-

gitation of rheumatic origin. It is true that the first sound over the aortic area was attended by a slight murmur indicating that there was some obstruction co-existing with incompetence. During the whole of the long period she was under my care she never had digitalis in any form. The treatment consisted of periodic rests in bed, a careful dieting in which animal food was as predominant as her means would allow, and for medicine the tincture of nux vomica at intervals. This line of treatment was resorted to when attacks of dyspnoea and other signs of cardiac failure appeared from time to time; but she was able to earn her living by fairly hard work, though accompanied by anxiety. Then she was transferred to another physician during one of her bad periods, and was given digitalis. After the ninth dose (1/4 x of the tincture) she suddenly died.

2. Clara G., aged 31, had acute rheumatism when she was 17 years old. A lady of independent means, she could hence rest when necessary, and follow medical advice without anxiety as to her livelihood. She consulted me seventeen years before her death, complaining of dyspnoea on slight exertion. I found that she had aortic regurgitation, pure and simple. I saw her from time to time and from year to year, at intervals of about three months, and never during that long period did she take digitalis in any form whilst she was under my care. Shortly before her death she went to reside in Surrey, where she again suffered from an attack of breathlessness, and requested her local medical adviser to write to me, which he did, giving me an excellent account of her symptoms and his treatment. His remedies contained digitalis in one form or another, and I wrote to him to express my opinion that this drug was harmful in her case. She died suddenly on the fifth day after digitalis was first prescribed.

3. A young practitioner, having charge of a ward in a poor-house infirmary, came to me in distress, bringing the following story: Three old men were under his care suffering from aortic disease, regurgitation being the principal or dangerous condition. As they were all showing signs of heart failure and had oedema of the lower extremities, my friend ordered each to take a mixture containing digitalis three times a day. One man died on the third day thereafter, another on the fifth, and the other on the seventh day—all somewhat suddenly.

4. A gentleman, aged about 60 years, was desirous of effecting a "short term" (a few months) assurance of his life. I found on examination that he had a blowing diastolic murmur over the aortic area and a reflux pulse. Seeing that I hesitated over his proposal, he volunteered this statement: "I suppose you are frightened at my heart. I know it is wrong, but it is no worse now than it has been for some years, and I have just been to my doctor, who has given me this prescription." It contained digitalis, whereupon his proposal was rejected. Within a few weeks—not more than three—a notice of his death claim was received.

5. Miss N., aged 17, consulted me in 1888 on account of shortness of breath. She had had two attacks of rheumatic fever since she was 14 years old. I found that she had all the signs and symptoms of aortic regurgitation, but the mitral valve was "still holding." I prescribed a diet and regimen similar to that mentioned in the first case, and gave her my opinion as to the inadvisability of her taking digitalis. She has obeyed this injunction to the letter. She married, against my advice, in 1892, being then aged 21 years, and she has now three healthy children. She called to see me in February last in apparent good health but with obvious signs of aortic regurgitation.

6. At a clinical lecture some few years ago I exhibited six patients, all adults. Three were suffering from aortic regurgitant disease accompanied also by mitral insufficiency, whilst the other three had only aortic disease (regurgitation). Owing to some mistake on the part of the hall porter the patients, all six, were allowed to climb a deep staircase to the theatre, although orders had been given that they should be taken up by lift. The three patients who had the most extensive valvular lesions—aortic regurgitation and mitral regurgitation—performed the journey with ease; the other three, who had aortic regurgitation only, were so distressed on their arrival at the demonstration that they could not be examined for quite an hour; and, indeed, I deemed it wise to retain two of them in the wards over night.

These last cases may be somewhat away from the subject I am discussing, but I quote them as illustrating the advantages which will accrue to a patient with aortic insufficiency when he shall also have mitral insufficiency coexisting. I could relate other cases in which sudden death has occurred within short periods of the administration of digitalis in aortic insufficiency pure and simple, but these will suffice.

And herein lies one of the points of my contention. Not only would it appear that a patient suffering from aortic regurgitation has less distress if he also has mitral regurgitation, but he is also capable of greater exertion. There is a safety-valve action through the mitral orifice comparable to the tricuspid safety valve on the right side during prolonged muscular strain. It is true that the one is a pathological condition and the other a physiological

one, but none the less they both exist under the differing conditions.

But in pure aortic regurgitation, the mitral valve still holding, the physics of the disease are vastly altered. The left ventricle receives the diastolic shock—sometimes more, sometimes less, according to the amount of completeness of the regurgitation. Two crowds—namely, an oncoming rush and a backward retreat—thus meet and collide in a limited space, with no outlet or retreat. Injury must accrue to something. Either the constituents of the crowd must be injured, or the boundaries of the containing space stretched or broken down.

In the case of the diseased heart it is the ventricular walls which, after bearing the strain, eventually give way. And, as is almost universally admitted, the diastolic pause is lengthened under the influence of digitalis. The period is augmented during which all this extra strain is being borne. Centrifugal pressure is enough to try the strength of the ventricular cavity in ordinary circumstances. But in Dr. Bruce's case the conditions are not the same. It is true that he describes a patient having dropsy and other signs of heart failure, the chief lesion being aortic incompetence. It is true, also, that his patient was relieved by a course of digitalis; but it is equally true that he was suffering from mitral disease, and I contend that, by supporting a failing ventricular compensation by digitalis, there is no actual proof that such treatment is to be pursued in pure aortic regurgitation. I will at once admit that I have seen cases of aortic regurgitation much benefited by the use of digitalis; but they were all of them cases in which mitral regurgitation was also present.

These contentions are obviously difficult of proof. Very little in therapeutics can be actually proved. I am still open to conviction if the weight of evidence is against me, and, as Dr. Clifford Allbutt wisely observes, "it is best that each observer should give the results of his own impressions as simply as possible," but if I may formulate an opinion, which has been forced on my mind by a somewhat extensive experience, I would at once say that although digitalis is the most powerful remedy in mitral disease which we possess, in cases of pure aortic regurgitation digitalis is not only a dangerous drug, but actually a poison.

DEGENERATION OF THE MYOCARDIUM IN HOT CLIMATES.

By W. CARNEGIE BROWN, M.D., M.R.C.P.

A SOMEWHAT unfamiliar result of residence in the tropics, but one which is by no means infrequent, is a peculiar form of degeneration of the cardiac muscle. Clinically and pathologically the condition is of much interest and importance, for the evidences of myocardial change are apt to be misinterpreted, and sudden and unanticipated death occurs not infrequently after an apparently trivial illness.

The patient is usually a man who has been some years in the tropics; he is confident that he is organically sound, and though he confesses he is not well, he is sure that there is little the matter. In answer to leading questions he says he is tired, yet sleepless; he is disinclined for exertion, mentally or bodily; he yawns unaccountably; he has flushings of the face, tingling and itching in the extremities; he perspires more readily and copiously than is his habit; and he has a vague feeling of uneasiness and apprehension. There is no faintness, no dyspnoea, no cough, no pre-cardiac oppression. Palpitation is absent, or it is slight, and felt only after exertion. There is perhaps some indigestion, and he has remarked that the feeling of weariness comes on coincidentally with discomfort in the stomach. He is thirsty, and wants more than his usual quantity of stimulant. He is not depressed, but he worries unnecessarily about his affairs, and is kept awake at nights by a stream of thought and indefinite anxiety. The pupils are large, and not actively responsive to volitional or reflex stimuli. The pulse is irregular, but it is full, soft, and deliberate; the vasomotor system is inactive, and there is no hardening of the arteries. The skin is moister and more lax than usual. The temperature is normal; the tongue is clean, and gives no indication of any morbid condition; the urine is

scanty, and loaded with urates, though otherwise normal. The rhythm of the natural cardiac cycle is unimpaired, but the heart's action is somewhat hesitating, and an occasional flutter or tremor may be detected at the apex; the impulse is softer, perhaps more diffuse, than normal, but it is not displaced. The clearness of note in the sounds is diminished; they seem a little distant, and the first is somewhat weak. Little further aid is to be obtained from the stethoscope; there are no adventitious murmurs, and examination of the pulmonary area and *conus arteriosus* gives no indication of deficient or defective blood.

In the absence of any definite cardiac symptoms, the malaise is apt to be set down as nervous or perhaps imaginary. The patient is told that there is nothing serious the matter, and he is dismissed with words of comfort and caution. In the great majority of instances, he reports in a day or two that he is quite well, but in some cases, and not very infrequently, the event is different, and the termination is tragic and startling. Generally without apparent cause, sometimes during a transitory attack of fever or indisposition, the undefined restlessness and weariness return. After an uncomfortable day, the patient goes to bed overtired; possibly he takes some sulphonal or veronal and falls asleep. In the middle of the night he starts up with a feeling of inexpressible alarm, there is a spasm of angina, and he falls back dead.

Contrasted with the commoner forms of myocardial decay the salient features of the disease are the absence of fainting fits, the inconsiderable impairment of the circulation, the vagueness of the physical signs, and the fact that fatal syncope usually supervenes in the absence of unwonted exertion. It is perhaps only a coincidence, but one the importance of which will not be overlooked, that death frequently occurs at night and after the administration of a soporific.

The *post-mortem* appearances are fairly uniform. On opening the pericardium (which contains no excess of fluid) the heart is found to be somewhat larger than normal. It lacks the resistance of a healthy organ, and is lax and flaccid. Epicardial fat is markedly deficient. If the viscus is removed from the thorax and held up by the great vessels, it sags and droops; in extreme cases the parietes are so loose that it collapses in mushroom shape over the hand that grasps it. Section into the ventricles shows no dilatation and no endocarditis; on the contrary, the lining membrane is blanched and bloodless. A similar condition prevails in the muscle; there are no infarcts, no points of haemorrhage, no deposits of pigment, no fat, no reaction to iodine, no serous atrophy. The coronary arteries are dilated and soft, and show no atheromatous change. The myocardium is pale, almost to whiteness, soft and friable. Microscopically, the intermuscular connective tissue is seen to be swollen, and apparently increased in quantity, but rounded infiltration and vascular congestion are absent. The condition of the individual fibres suggests cloudy swelling; the section looks waterlogged, it appears to have a general boggy turbidity, the striae are hazy and indistinct, and there is occasional rupture and fragmentation of the ultimate strands. From other indications it is obvious that the general nutrition is profoundly impaired, and that the myocardial lesions, as is the case in other retrogressions of tissue, are only prominent incidents of a general dyscrasia. The liver is small, and its section is pale and pasty-looking. There is no definite atrophy of the intestines, but their coats are thin and the villi and mucosa are shrunken and deficient in lymphoid tissue. The body fat is not diminished in quantity; in places it seems to be in excess, but it is dull and discoloured, and its distribution is irregular. In two necropsies in which a careful examination of all the organs was made, it was noted that no parasites, systemic or intestinal, were found, and that blood and bone marrow were normal.

The disorder appears to have no racial preference. It is questionable, indeed, if it is "tropical," in the sense that it is limited to tropical countries. Under the name of "soft heart" a similar macroscopic condition was long ago described by Louis, Laennec and Stokes, but its histo-pathology, which was worked out by v. Recklinghausen, is altogether different. The cloudy swelling of the myocardium seen after death from acute fevers has,