followed by salines, and a mixture containing magnesia sulphate, acetate of ammonia, and sal ammonia, to act on bowels, skin and liver, is often useful.

Where the pain is due to old uterine or perimetrial inflammation, rest and other suitable means must be adopted to promote absorption of the inflammatory products and return of uterine mobility, but as some of these cases require surgical treatment, either by vaginal or abdominal section, I cannot go into details now.

The treatment of *ovarian dysmenorrhoea* is more difficult, but when the pain is due to pre-existing perioöphoritis much may be done to lessen discomfort on the lines laid down for congestive dysmenorrhoea. The actual pain is relieved by many of the drugs alluded to under spasmodic dysmenorrhoea, but the bromides of potassium and sodium and nitroglycerine are particularly indicated. Turpentine stupes or hot bran poultices over the abdomen are very soothing.

Operative treatment is unsatisfactory, and, in my opinion, is not justifiable except when this form of pain occurs in women with absent or undeveloped uteri. Some of you will remember a typical case of this in Golding Ward about five months ago. She was a woman of 28 years of age, who had true epileptic seizures and dysmenorrhoea once a month, with absolute primary amenorrhoea. On examination under anaesthesia the vagina was found to be short but patent, no uterus could be felt, but what appeared to be an ovary was felt on one side. It appeared to be a case of ovarian dysmenorrhoea with absent uterus. At the urgent request of the patient and her friends I opened the abdomen, and found that each ovary (one was inside the inguinal canal of its own side) was attached to a small solid muscular nodule which represented one half of a double uterus, and on each side there was a much hypertrophied round ligament. I removed these structures. The patient had no dysmenorrhoea at her next period, but had an epileptic fit of lessened severity, with little or no subsequent coma. Since then she has only had one slight fit, and is much improved mentally.

The treatment of "Mittelschmerz" is most unsatisfactory. I cured one case by removing a vaginal cyst which became tense and caused pain on the tenth day after each period. Cases due to chronic salpingitis have also been cured by operation. Cases due solely to the ovulation cycle occurring at a different time to the menstrual cycle, are relieved somewhat by a blister over the ovarian region, but as a rule sedatives have to be given, and nothing short of removal of the offending organ will cure. Needless to say, this is not often to be recommended.

You will see that the treatment of dysmenorrhoea depends not only upon the recognition of pelvic diseases and abnormalities, but also upon a due regard being paid to constitutional or general conditions. The practitioner who devotes all his attention to local causes, to the exclusion of the general state, is as little likely to relieve complex cases of dysmenorrhoea as he who neglects to adequately investigate pelvic conditions, and who treats such cases by sedatives, which merely mask the symptoms without curing the condition which causes them.

REFERENCE.

- ¹ BRITISH MEDICAL JOURNAL. EPITOME, August 30th, 1902, No. 121, and October 25th, 1902, No. 255.

A SERIES OF FIFTY CONSECUTIVE ABDOMINAL SECTIONS.*

By R. P. RANKEN LYLE, M.D.,

LECTURER IN MIDWIFERY AND DISEASES OF WOMEN AND CHILDREN,
UNIVERSITY OF DURHAM COLLEGE OF MEDICINE.

In December, 1902, the Samaritan Hospital was instituted in Newcastle-upon-Tyne for the practice of gynaecology. There are six beds, including one emergency bed, for patients, and an out-patient department, which is open two days a week. All the operations are performed in an ordinary attic, under the most restricted conditions, and it has not been specially fitted up in any way. Since its opening I have performed a large number of major and minor operations, and my object now is to show you the result of the first 50 abdominal sections, and to draw certain conclusions therefrom.

The period during which they were performed extends

* A paper read before the British Gynaecological Society.

from October 1st, 1903, to January 18th, 1906, and the conditions necessitating them were as follows:

Ovarian cystomata, including dermoids	...	18
Fibromyomata of the uterus	...	11
Ectopic gestation	...	6
Broad-ligament cyst	...	3
Adherent retroversion	...	3
Pyosalpinx	...	3
Tuberculous disease of the peritoneum	...	2
Malignant disease of the cervix uteri	...	2
General abdominal carcinosis	...	1
Parovarian cyst	...	1
Total	...	50

The actual operations were:

Ovariectomy	...	17
Supravaginal hysterectomy	...	8
Salpingectomy	...	5
Salpingo-oöphorectomy	...	4
Enucleation of cyst	...	4
Coeliotomy	...	4
Myomectomy	...	3
Ventrifixation	...	3
Panhysterectomy	...	2

I have, unfortunately, to record one death. The case will be fully described.

THE TECHNIQUE ADOPTED.

The technique adopted in all these operations was of the simplest nature, as I am of opinion that complicated technique or elaborate preparations are not only unnecessary but may be even a source of danger to the patient, inasmuch as minor essentials may be sometimes accidentally overlooked when the preparations are too elaborate.

Antisepsis was very carefully carried out, the hands and forearms were thoroughly washed in hot lysol solution with a nailbrush, then turpentine and methylated spirit were used, and the hands and forearms well washed in 1 in 1,000 corrosive solution. The instruments were boiled and placed in 1 in 40 carbolic solution. The sponges, which were of gauze, were boiled for twenty minutes, and were then wrung out of hot saline. All buried ligatures were of catgut, which was prepared by ether, methylated spirit, and corrosive sublimate solution. Silkworm gut was used for all superficial sutures. The abdomen was prepared the night previous by washing with soap and water, then sponged with turpentine and methylated spirit, and the application of a corrosive pad over night, and washed with 1 in 1,000 corrosive sublimate solution just previous to the operation. No vaginal antisepsis was carried out even in cases of panhysterectomy. The abdominal incision was restricted to the smallest dimensions consistent with the proper performance of the operation, and there was consequently a marked absence of shock following the operation, and a diminished risk of ventral hernia in later years. All cut surfaces of peritoneum were properly adjusted—such as suturing the anterior and posterior layers of the peritoneum after hysterectomy and the removal of broad ligament cysts, and suturing the peritoneum over the pedicle in cases of ovariectomy and salpingectomy.

The abdominal wall was carefully sewn up in three and sometimes four layers, care always being taken to properly adjust corresponding structures. A small strip of plain sterilized gauze wrung out of hot saline and covered with plain absorbent wool was used as a dressing to the wound. The dressing was kept in position by strapping and flannel bandages. Ether, with one or two exceptions, was the anaesthetic. In three cases of pyosalpinx, in which a small amount of purulent material escaped during the operation, it was carefully mopped up and did not cause any further trouble. In those cases where a large wounded surface of peritoneum occurred, for example, ventral fixation for adherent retroversion, as much saline solution as possible was left in the abdomen to float the intestines from the wounded surface and so prevent adhesions.

After-treatment.

The patient was given a drink of hot water with a little bicarbonate of soda in it (which is often effectual in relieving bilious vomiting) about three hours after the operation. This was freely continued in addition to barley water until the following morning, when she was given a cup of tea, followed during the day by milk diluted with barley

water or soda water. Early on the second morning after operation, calomel was given in two doses of 5 and 3 gr. respectively, at an interval of one hour, followed by a hot seidlitz drink half an hour later. As soon as the bowels were moved the patient was put on a liberal diet. The sutures were removed on the tenth day.

ANALYSIS OF THE CASES.

Ovarian Cystomata.—Of these there were 18 cases divided as follows: Simple unilocular, 6 cases; multilocular simple, 4 cases; multilocular undergoing colloid degeneration, 3 cases; multilocular suppurative, 1 case; multilocular malignant, 1 case; multilocular papillomatous, 1 case; dermoids, 2 cases.

In addition to these, a case of left tuberculous pyosalpinx was found, complicated with a right ovarian cyst the size of a goose egg; and a case of cervical fibromyoma had a cystoma of each ovary.

The following cases were of special interest:

CASE VII.—Suppurating multilocular ovarian cyst with purulent peritonitis. This patient was exceedingly ill, having had a temperature of 104° F. for two evenings before the operation. On opening the abdomen the peritoneum was very inflamed and oedematous, and the omentum was adherent to the tumour, which was about the size of a seven months gestation. On separating the omentum several ounces of encapsuled pus escaped, the tumour was removed, the posterior vaginal fornix was opened and a gauze drain was inserted through it into the pelvis. The temperature fell to 97° F. on the morning after the operation, but was 105° F. the following evening; after a great deal of fluctuation it gradually fell to normal six weeks later, when she was discharged. Some days after leaving the hospital she had a relapse, and unfortunately died a fortnight later.

CASE XXVI.—Ovarian dermoid with twisted pedicle. The twist occurred about one week before admission, when there were the usual symptoms of peritonitis. On opening the abdomen the tumour was found partly gangrenous. She made an uninterrupted recovery.

CASE XLII.—Malignant ovarian cyst. The whole of the peritoneum and omentum was infiltrated with hard, malignant masses, and the tumour was found adherent in all directions and commencing to disintegrate. It was not removed, but the abdominal cavity was filled with saline solution and then closed. Convalescence, as far as the operation was concerned, was uneventful.

CASE XLIV.—Enormous multilocular ovarian cyst undergoing colloid degeneration. This patient was confined nine weeks previously of a seven months child, since that time the abdomen rapidly enlarged, and on admission was about three times the size of a full term gestation. She was very emaciated and very ill, pulse 120, very feeble, and temperature 99° F. She had severe dyspnoea, and was unable to lie down during the whole night before operation. The patient had to lie on her left side on the operating table, as respiration ceased when she was turned on her back. The abdomen was opened with her on her left side, and about two gallons of ascitic fluid escaped. The tumour, which was about 1½ times the size of a pregnant uterus, was adherent to the under surface of the liver and diaphragm. The pedicle was clamped and divided, and the tumour removed from below upwards. The other ovary was about the size of an ostrich's egg. It was also removed, and the abdomen filled with about 1½ gallons of normal saline. During the greater part of the operation the patient was practically pulseless, and respiration temporarily ceased on two occasions, the anaesthetic (ether) was not pressed, and during the closing of the abdomen she was partly conscious. She got three injections of ⅓ grain of strychnine. When returned to bed the pulse was just perceptible at 180, shortly afterwards it fell to 160, and in the evening it fell to 140. She made a good recovery, and six months later was reported to be in good health.

CASE XLV.—Right ovarian cyst. In addition this patient had a left ovarian abscess which contained 4 or 5 oz. of very fetid yellow pus and a double hydrosalpinx. She made an uninterrupted recovery.

Fibromyomata.—Eleven operations were performed for the removal of fibromyomata, and in every case the symptoms complained of by the patient were sufficient to render the operation one of necessity. Supravaginal amputation of the uterus was performed in 8 cases, and myomectomy in 3. Myomectomy was the operation of selection, but in those cases where it was not feasible, supravaginal hysterectomy was performed, care being always taken to leave the ovaries, in order to save the patient from the ordinary consequence of their removal prior to the menopause. An artificial menopause practically means that the patient will be an invalid for many months, and it is highly desirable that this, if possible, should be prevented. It was not possible, however, to

leave the ovaries in Case XLVII, as there was a cystoma of each ovary.

The following cases were of interest:

CASE III.—Myomatous uterus and large ventral hernia—death. This patient was operated on eight years previously, the ovaries being removed in the hope of curing her. The result, however, was that she continued bleeding, making her a chronic invalid; besides the tumour increased in size, and she developed a very large ventral hernia, which reached to her knees and totally incapacitated her from work. She was very corpulent and anaemic. The abdomen was opened by an elliptical incision, extending from the umbilicus to the pubis, and surrounding the hernial sac, which contained nearly all the intestines. They were entirely adherent to the hernial ring, which was 4 in. long by 3 in. broad. The intestines were separated from the ring, which was dissected out. On examining the pelvis everything was found adherent, a myomatous uterus, the size of a large cocoanut, was freed from its adhesions and lifted out of the pelvis, the broad ligaments ligatured and divided, and the tumour removed, the cervix being left. The peritoneum was sutured over the stump. With some difficulty the intestines were returned into the abdomen, and the abdominal wall sutured in four layers with strong catgut. She developed a small faecal fistula in the lower angle of the wound, which practically ceased discharging on the seventh day. During the next two days her condition was very satisfactory, but on the morning of the tenth day, at 2 a.m., she was suddenly awakened with severe pain in the region of her heart, and much difficulty in respiration. She was deeply cyanosed, pulse very rapid, irregular and feeble. She was given stimulants internally, with strychnine and digitalin hypodermically. Considerable improvement took place till 6 a.m., when she had a similar attack, and died in ten minutes, death being due to pulmonary embolism, probably fat embolism.

CASE XXXII.—Myomatous uterus—supravaginal hysterectomy. The tumour was the size of a fetal head and impacted in the pelvis, causing partial intestinal obstruction. Patient had purulent cystitis and was seriously ill. On opening the abdomen the bowels and omentum were found adherent all over the front of the tumour. These were separated, and both tubes were found distended with pus. While removing them they both burst and some pus escaped. This was carefully mopped up. The myomatous uterus was removed and the abdomen closed. The patient made an uninterrupted recovery.

CASE XLIII.—Subserous fibromyoma—myomectomy. This patient was five months pregnant, and suffered severe pain on the left side of the abdomen. The abdomen was opened at the left border of the rectus, and a pedunculated myoma the size of a large goose egg was found in the left iliac fossa, and was removed from the surface of the pregnant uterus. The uterus then rotated from left to right through a quarter of a circle, so that the point of attachment came to lie in the middle line anteriorly. The patient went on to full term and had a normal confinement.

Ectopic Gestation.—There were 6 cases of ectopic gestation. Case XVII was admitted on August 2nd, 1904. Rupture took place at 4 p.m. the following afternoon as she was lying quietly in bed. The patient rapidly became collapsed, but fortunately before an hour passed I was able to proceed with the operation, which was in no way difficult. The abdomen was full of recent blood clot; this was not removed. All these 6 cases made uninterrupted recoveries.

Broad Ligament Cyst.—There were 3 cases; none of them presented any difficulty. The cysts were opened, the capsules enucleated, and the wounds in the broad ligaments sutured with fine catgut. They all did well.

Pyosalpinx.—In addition to the case which was associated with a myomatous uterus, and which is already described, there were 3 cases:

One of these—Case XXXIV—was tuberculous. It was single, and was associated with an ovarian cyst of the opposite side. The pyosalpinx was matted with some knuckles of intestine, which were involved in the disease, and during its removal the bowel unfortunately got injured. This was carefully and with some difficulty sutured. A strip of plain sterilized gauze was inserted from the wound in the bowel to the lower end of the abdominal incision to determine the track of a faecal fistula, should such occur. The following day the gauze was removed, and two days later a small amount of faeces came through the lower angle of the wound. This continued for six days, then ceased, and the fistula completely closed. With this exception, convalescence was quite uneventful.

CASE XXVIII.—Double pyosalpinx of old standing. On opening the abdomen the intestines were found matted with old and dense adhesions, over the tumour. They were separated with much difficulty, and during the removal of the tumours one of them burst and a quantity of pus escaped. This was very carefully mopped up and the abdomen closed without the insertion of any gauze for drainage. Three days later some discharge, which had a faecal odour, came through the lower angle of the

wound. This continued for five days, when the fistula closed spontaneously, leaving a hollow, granulating surface about 2 in. square. Three silkworm-gut sutures were inserted, pulling the granulating surfaces together; the wound healed perfectly, and the patient did well.

Malignant Cervical Disease.—Panhysterectomy was performed in two cases of advanced malignant disease of the cervix in order to relieve severe symptoms of haemorrhage and pain; both patients recovered, although one of them developed a temperature which on one occasion was as high as 102.6° F. The temperature, however, remained perfectly normal for nine days before she was discharged from the hospital.

DRAINAGE.

Gauze was used in three of the earlier cases (Nos. v, vii, and xii) for the supposed purpose of drainage, and the result in all these cases was unsatisfactory. In Case v, panhysterectomy for ulcerating malignant disease of the cervix, gauze was used from the pelvis into the vagina, and the result was a gradual rise of temperature to 102.6° F., whereas in a similar case (Case xlvi) no gauze was used, and the patient made an uninterrupted recovery. In the case of the suppurating multilocular ovarian cyst (Case vii) the posterior vaginal fornix was opened and the pelvis drained through the vagina. The result was a pelvic abscess, which had to be douched for many days afterwards, and to which, I have little doubt, her high temperature and unsatisfactory convalescence was partly if not mainly due. In Case xii gauze was also used for the supposed purpose of drainage, on account of general oozing, and although this patient had a normal convalescence for four weeks after the operation, a faecal fistula developed during the fifth week.

I have since then altogether abandoned the use of gauze for drainage purposes in abdominal and pelvic work, unless in cases of pelvic abscess, where the abscess wall cannot be removed, and have since followed the following rule: If a little pus accidentally escapes I carefully mop it up, and in case a large amount escapes and soils the intestines, I wash the intestines and pelvic cavity thoroughly with normal saline and then close the abdomen, leaving a considerable quantity of saline behind. In cases of general oozing I first secure the more important points, and then fill the pelvis with saline.

I have followed this rule in several such cases on which I have operated since, and the result has been quite satisfactory in every case. I have now no doubt that the gauze in the earlier cases did not act as a drain, but merely as a superfluous and noxious foreign body, and a local irritant to any intestinal surface with which it came in contact.

TEMPERATURE.

A study of the temperature charts of the series is most interesting, considering the circumstances under which the operations were performed. There was not a single case of "sepsis from without," and in no less than 41 cases the temperature did not exceed 101° F. even on a single occasion, while in 28 it never exceeded 100.4° F.

In 5 cases the temperature exceeded 101° F. on two occasions only, but did not exceed 102° F. These cases were:

No. xvii. Recent ruptured ectopic with severe internal haemorrhage.

No. xxi. Ovariectomy for simple ovarian cyst.

No. xxxiii. Supravaginal hysterectomy for large myomatous uterus in a very corpulent subject.

No. xlii. Double ovariectomy. The abdomen was enormously distended, and the patient was extremely ill on admission.

No. xlvi. Panhysterectomy for ulcerating malignant disease of the cervix.

In Case iii, which proved fatal, the temperature was marked between 101° F. and 102° F. on three occasions. Only in two cases was the temperature recorded above 102° F. and less than 103° F. These were:

No. v. Panhysterectomy for ulcerating malignant disease of the cervix.

No. xlviii. Advanced malignant disease of the peritoneum. The temperature was due in this case to a somewhat mild attack of left-sided pneumonia at the end of the second week after the operation.

Only in one case was there any serious degree of temperature. This case was one of suppurating multilocular ovarian cyst with purulent peritonitis, but the

temperature was 104° F. for two nights prior to the operation.

The following table summarizes the charts:

	One Occasion Only.	Two Occasions Only.	More than Two Occasions.	Total.
Not exceeding 100.4° F...	—	—	—	28
100.5° F. to 101° F. ...	9	3	1	13
101.1° F. to 102° F. ...	—	5	1	6
102.1° F. to 103° F. ...	1	1	—	2
Above 103° F. ...	—	—	1	1

GENERAL OBSERVATIONS.

From this series the conclusion appears to be deducible that attention to the following points is highly desirable in order to obtain good results:

1. A correct preliminary diagnosis.
2. Simplicity in operative technique.
3. The simplest antiseptic preparations combined with asepsis.
4. The use of few surgical instruments and appliances.
5. Limited abdominal incisions.
6. The correct adjustment of all cut peritoneal surfaces.
7. The complete closure of the abdominal cavity without drainage of any kind.
8. The free use of normal saline solution in cases where purulent material gets accidentally spilled in the pelvis or abdomen during the performance of an operation.

Finally, I desire to express my great indebtedness for these results to Mr. J. W. Leech and the many practitioners and students who assisted me with the operations. I have also to record my high appreciation of the careful preparations, assistance at the operations, and careful nursing of all these patients by the sister in charge, and on account of which the entire elimination of sepsis was mainly due.

MIDWIFERY EXPERIENCES:

A RECORD OF FORTY-SEVEN YEARS' WORK.

By JOHN LINDSAY, M.D., J.P.,

PRESIDENT OF THE GLASGOW AND WEST OF SCOTLAND BRANCH,
BRITISH MEDICAL ASSOCIATION.

SINCE 1875 I have had assistants, till within the last few years, unqualified. Since that time I may not have been in actual attendance on more than the half of the cases, but upon nearly all serious cases I have been told about anything unusual.

When I started practice in 1859 I had experience of the usual kind of about 50 maternity cases in England. I had never seen labour carried out except in bed, or on the floor when there was no bed. In our district—and I do not know how far around—it is a very common habit to go through labour out of bed, the patient on her knees holding on by a chair. I resisted this very strongly at first, but had to give way sometimes when the patient said she could not lie and those in attendance pressed the matter, saying that her old doctor would have allowed it, and she would have been better long since. I soon got to see that it was not a bad way for a poor woman with, perhaps, a very scanty supply of clothing. The whole mess—often a very ugly one—is left on the floor, and with a warm dry cloth to the parts and a quick shift to bed the patient is pretty comfortable. I daresay more than half of our patients have been delivered in this way. I once tried the forceps in this position, but did not use them well and never tried again.

There has been no use of antiseptics except some antiseptic oil or grease for the finger, hand, or instruments. Many times we have been caught on the roads with no preparation. From the first I have always been very careful in washing my hands and changing my outer coverings if we had infection on our rounds, though this was not always possible.

It would be wrong to say that I have never poisoned a woman with my finger, because I was once caught a good many miles from home on my way from a *post-mortem*