NOTE ON A FILARIAL LARVA IN THE
BLOOD OF A BLACKBIRD.By WILLIAM ST. CLAIR SYMMERS, M.B.ABERD.,
MUSGRAVE PROFESSOR OF PATHOLOGY, QUEEN'S COLLEGE,
BELFAST.

In the month of July of the present year a blackbird (*Turdus merula*) dropped dead from a tree in my garden in Belfast. I examined this bird, expecting to find trypanosomes in its blood, but instead there were very numerous tiny round "worms," which were evidently the larvae of some member of the Filaridae. Avian filariasis is, I believe, not uncommon in tropical and subtropical lands, but the occurrence of such a thing in the north of Ireland is worth mention, and may lead to further search for this condition among the birds of northern countries. I regret that I have no information as to the migratory habits of the blackbird.

As to the bird in question, the internal organs were normal to the naked eye, and the heart blood contained no bacteria that could grow on the usual media of the laboratory. A small quantity of blood was, in the first instance, withdrawn by a Pasteur pipette from the heart; a tiny droplet blown upon a glass slide, and without being covered by a cover-slip, was examined under a low power. It was at once seen that in two places in the field of vision the blood corpuscles were in active commotion, and it soon became evident that this was due to the wriggling of a worm-like parasite. This parasite was best seen after diluting the blood with normal salt solution, and by using higher powers of the microscope. The larvae were present in every small droplet of the heart blood, but none were found in the blood of the liver.

When examined in the fresh blood (diluted several times with salt solution) the animal moved rapidly, rolling itself into coiled spirals like a pile of rope, unrolling immediately, curving and twisting without ceasing, at times forming a perfect figure 8. There was practically no tendency towards progredient movement, so that the animal could be observed without much use of the movable stage of the microscope, although it occasionally rolled out of the field of vision. The movements at this stage were peculiarly smooth, continuously active, graceful, and quick. After a few hours (room temperature) the movements were slower, and gradually became more and more jerky, until after twenty-four hours the jerkiness was the chief characteristic of the motion, which then resembled the awkward action of a mechanical toy, the rope-coil position was no longer assumed, nor were figures of 8 formed. The sluggish, and doubtless dying, animal moved with jerks, as if with effort, bending stiffly; at times the anterior quarter of the body forming nearly a right angle with the posterior portions. At no stage of the movements was there any extension or contraction of the body. On several occasions an individual taken freshly from the heart blood protruded from its anterior extremity a tiny tent like pellicle or cupola, which was immediately withdrawn only to be shot out again after a few moments. This "prepuce" was more pointed at its summit than the similar arrangement seen in the larva of Bancroft's filaria, nor did it show any thread or cilium projecting from its top. But owing to the quick movements of the animal and the rapid withdrawal of this cupola (prepuce) my eye could not catch very well the anatomy of this part. The more sluggish animals did not show this phenomenon, the extremity being rounded off and having a tiny dimple indenting the (blunt) vertex. After forty-eight hours the animals (covered and vaseline-rimmed) were motionless, and lay extended in more or less straight lines—except where crowded by blood corpuscles.

The larvae were devoid of a sheath. They measured on the average 0.15 mm. in length and 0.0035 mm. in breadth, being a little thicker at the anterior than at the posterior end, but still of nearly equal thickness throughout, and the two extremities were bluntly rounded off. The cuticle was smooth. I did not notice an excretory pore. The dimple above mentioned as existing in the living animal could not be detected in the dead. When unstained, the animal was structureless internally, and was colourless and clear; when stained by Leishman's stain, numbers of small granules were seen packed together and occupying nearly the whole body; these granules stained deep blue,

and lay in a homogeneous matrix which stained less deeply. At certain places the granules were absent, leaving a clear triangular space (V-spot). Such spots were not invariable as to position, but one of them was fairly constant, and lay at the junction of the anterior and middle third of the body. Moreover, there was a striking and constant appearance—namely, a clear, unstained oval area, lying about midway between the two extremities. This area occupied the whole calibre of the body, and measured 0.01 mm. in length. These details were best seen when the preparations, after staining in the usual way with Leishman's stain, were rapidly washed alternately in alcohol and in water, so as to remove some of the dye.

The adult filaria was not found.

The larva here in question is evidently much like the *Filaria perstans*, being without a sheath, and having both extremities blunt. In size it is smaller than *perstans*, but is apparently almost identical in this respect (size) with the larva described by Manson in natives of British Guiana, and held by this distinguished observer as closely resembling, if not identical with, *F. perstans*.

I hesitate in the identification of this blackbird filaria, and consequently refrain from giving it a name.

MEMORANDUM ON
THE OBSERVATION OF SPIROCHAETES IN
YAWS AND GRANULOMA PUDENDI.By ALEX. MACLENNAN,
DISPENSARY SURGEON, WESTERN INFIRMARY, GLASGOW.

I HAVE recently had the opportunity of examining some smear preparations from yaws sent to me by Dr. Branch of St. Vincent, B.W.I., to whom I am very much indebted. The smears were fixed in methyl alcohol by Dr. Branch and were stained by me in a number of ways. In one specimen taken from a papilloma in recurrent yaws numerous spirochaetes were observed. The method of staining was by Giemsa's solution and gentian violet, as described by me in a recent number of the BRITISH MEDICAL JOURNAL. These spirochaetes possessed similar characters to those of the *Spirochaeta pallida*, as already pointed out by Castellani, but the staining was fainter. Loops in the course of the filament were observed, as in the *pallida*.

In two smears from *Granuloma pudendi*, sent also by Dr. Branch, I have found a few spirochaetes—most probably the *refringens*—while in one stained first in fuchsin, then in acetone gentian violet, a number of very long highly-refractive spirochaetes were observed. These parasites were much longer than the *Spirochaeta pallida*, and the waving was finer and closer. In one forty waves were counted. Though the staining was very prolonged, these spirochaetes were very faintly stained. There were also a number of other structures present which were less definite.

SOCIETY FOR RELIEF OF WIDOWS AND ORPHANS OF MEDICAL MEN.—A quarterly court of the above Society was held on October 10th, Dr. Blandford, President, in the chair. Fourteen directors were present. Proposals for membership were read from two medical men, and they were duly elected. It was reported that since the last court two widows, annuitants of the charity, had died, and one had become ineligible owing to an increase of income. Of the two who had died one had received in grants £621 and the other £500, their husbands having paid in subscriptions £52 10s. and £15 15s. respectively. The one who was no longer eligible had received £900, her husband having paid a composition fee of £21 for life membership at the time of his election. It was proposed to distribute the sum of £556 among the annuitants of the charity as a Christmas present, each widow to receive £10, each orphan £3, and those on the Copeland Fund £5 each. Six letters asking for relief had been received during the past quarter, but the applicants were ineligible under the by-laws of the Society, relief being only granted to widows of deceased members and to orphans below the age of 16. Any registered medical practitioner resident within a radius of 20 miles from Charing Cross is eligible for membership. Further particulars may be obtained from the Secretary, 11, Chandos Street, Cavendish Square, W.