muscular rheumatism. Chart 13 shows the course of a case treated with it.

Salicin, saligenin, and aspirin (acetyl-salicylic acid) form salicylic acid in the body, and in virtue of this have an antirheumatic action; while besides these there are numerous other compounds of salicylic acid which are active, but for the most part inferior to sodium salicylate.

There can be no reasonable doubt that compounds of salicylic acid, or substances from which it can be formed in the body, have a specific action against the poison of acute rheumatism. The number of such specifics is small in medicine, comprising quinine in malaria, mercury in syphilis, a few serums and arsenic in malaria, in pernicious anaemia, in chorea, in lymphadenoma, and in sleeping

Chart 13.—Action of methyl salicylate.

sickness. The action of arsenic in all these diseases is comparatively feeble. In all it requires prolonged administration, in some it merely lessens the vitality and virulence of the parasite, and hence improvement is often slight and relapses are very common. In the same way it is evident that salicylates in acute rheumatism frequently exert an action too feeble to satisfactorily cut short that disease, and that this is most apparent when the inflammation is seated in the fibrous tissues connected with the muscles and joints, or in the valves of the heart. It would certainly be an advantage to have a new remedy acting similarly but more powerfully, and, if possible, without its toxic effects on the central nervous system. There is no reason why such specific remedies should not be discovered for all the ordinary fevers if a large series of likely substances was systematically tested in our fever hospitals.

THE RECOGNITION OF MYELOPATHIC ALBUMOSE IN THE URINE.

By T. R. BRADSHAW, B.A., M.D., F.R.C.P.,

PHYSICIAN, LIVERPOOL ROYAL INFIRMARY; LECTURER AND EXAMINER IN CLINICAL MEDICINE, UNIVERSITY OF LIVERPOOL, ETC.

THE detection in the urine of the peculiar substance first described by Bence Jones in the year 1847, is one of the most significant clinical observations which it can fall to the lot of the physician to make. Its presence is one of the few signs of disease which can be strictly called pathognomonic, admitting, as far as our present knowledge goes, of only one interpretation, and it is usually for many months the only indication of the true nature of the disease from which the patient is suffering, and may for a long time be the sole evidence that he is seriously ill at all.

Multiple myeloma, the morbid condition of which it

is the sign, goes on to a fatal termination with the certainty of malignant disease, and no measures hitherto attempted appear to have the slightest influence in retarding its progress. In fact, a correct diagnosis has seldom been made until the disease has become pretty widely distributed, and clearly beyond the reach of any local measures which might conceivably be employed to arrest it. We may be certain that in this, as in other forms of new growth, any remedial measures that may hereafter be devised will be only applicable in the early stages of the disease, and their success will depend upon an early diagnosis. As this is only possible by the detection in the urine of the body which is the subject of the present paper, I think it is of the highest importance to have a clear understanding of the means by which it may be recognized.

Myelopathic albumose, as, for reasons set forth in former contributions, I designate the body in question, is a member of the great group of nitrogenous colloid substances known as proteids. Elaborate investigations have been made during the last few years, especially in Germany, with a view to determining its exact position in the group; but, as might be expected in view of our scanty knowledge of the constitution of the proteids in general, no great progress has been made in this direction. This much, however, may be taken as established—it differs essentially from all other proteids which have ever been met with either in health or disease. It is closely related to the native albumins, but at the same time presents characters which bring it near, if not within, the group of albumoses. It is remarkable in this that it is the only simple proteid derived from the human body that has ever been made to crystallize artificially. Further, it shows a remarkable readiness to undergo what I can only describe as molecular changes under conditions not easy to determine, changes which make its detection difficult if we are not aware of the possibility of their having taken place.

Myelopathic albumose is physically and chemically closely related to the albumins, and when present in urine generally gives a positive response to the common tests for albumin, for which it is consequently most likely to be mistaken. Thus it is thrown out of solution by heat in the presence of a trace of acid, by ferrocyanide of potassium and acetic acid, by picric acid, and by mercurio-potassic iodide. On the other hand, its response to these tests is, as a rule, not identical with that of albumin, the most noticeable points of difference being with regard to its solubility or otherwise at various temperatures and the greater or less readiness with which it is precipitated by various mineral acids. Some of the points of distinction can only be properly demonstrated by somewhat tedious chemical methods, and will only be briefly alluded to here. The object of this paper is to point out simple methods of analysis which are within the reach of every practitioner who pays attention to the urine at all, which require no complicated apparatus, and which, if carried out with ordinary care, may be relied on to distinguish myelopathic albumose from all other possible constituents of the urine.

I. Heat Test.

Myelopathic albumose coagulates at a much lower temperature than albumin—namely, at 58° C., or thereabouts (while serum albumin coagulates at about 75° C.)—the exact point varying a little in both cases according to the degree of acidity of the fluid, the concentration of the solution, and other conditions not thoroughly understood. Thus, in the case of albumose coagulation is sometimes accelerated by agitating the fluid. The best way of applying the heat test is on a water bath, which can be easily improvised by placing the test tube with the urine inside another of larger calibre containing water. When thus used the test is very striking, the whole column of urine becoming suddenly opaque at about 58° to 60° C., and the coagulum running together to form flakes at two or three degrees higher. Occasionally some turbidity is noticed at a much lower temperature, especially if the acidity is somewhat high. The recognition of this low coagulation point I consider conclusive evidence of the presence of albumose. I have never failed to find it in myelopathic albumosuria, and it never occurs in albuminuria. Care must, however, be taken to add only enough acid to ensure coagulation. A slight excess of acid may prevent its taking place altogether.

II. Solution on Boiling.

Another important characteristic of myelopathic albumose is that the coagulum formed at a comparatively low temperature undergoes more or less complete solution when the temperature of the fluid is raised to the boiling point. The fact of solution having occurred may be obvious from the clearing up of the turbidity; if it has only taken place to a slight extent it may be demonstrated by throwing the liquid while hot on a pleated filter, when the filtrate will become turbid on cooling. This reaction, when positive, is highly characteristic of myelopathic albumose, but it is not always obtained unless certain precautions are taken. Mere excess of albumose alone seems to interfere with it, so that it only appears after dilution. Again, it may be obtained merely by adding a little acetic acid to the hot urine. The conditions which determine the solution of the coagulum at the boiling point are not thoroughly understood. In this connexion it is worthy of remark that where little or no solution takes place on boiling the flakes of coagulum run together to form a putty-like mass, which collects on the top of the fluid, leaving the lower part almost clear. This appearance is rather characteristic of albumose, but must not be mistaken for clearing of the fluid by actual solution of the coagulum.

III. Reaction with Hydrochloric Acid.

Another characteristic of scarcely less importance even than the low coagulation point is the behaviour of this body with strong hydrochloric acid. If the albumose is abundant the acid may be poured in freely, when a dense precipitate takes place immediately, and is not dissolved unless an enormous excess is added. If the albumose is only present in small quantities the contact method should be employed, the urine being carefully floated on to the acid, when a sharp white band will appear at the line of junction of the fluids. This test is exceedingly delicate, and does not depend on the presence of excess of salts. A positive result may be obtained with it after free dilution of the urine; I have observed it after dilution with some thirty volumes of water. It is, however, more distinct if the dilution is made with a weak solution of a neutral salt. The hydrochloric acid test, though less sensitive than the nitric acid test to be presently described, is sufficiently delicate to indicate the presence of myelopathic albumose in quantities less than 0.5 per 1,000. Its great value, however, lies in the certainty with which it enables us to distinguish the albumose from other proteids. Hydrochloric acid added to albuminous urine causes no turbidity unless either the albumin is present in a very large amount or salts are in a concentrated state. If either of these conditions is present a moderate degree of turbidity may be produced by hydrochloric acid, especially on standing, but in these cases the previous dilution of the urine with three or four volumes of water will prevent the appearance of any turbidity. If these precautions are attended to, the appearance of a turbidity with hydrochloric acid is conclusive evidence of the presence of myelopathic albumose. This test is convenient for ordinary bedside use; it requires no special apparatus or manipulative dexterity, and is but little, if at all, less conclusive than the determination of the coagulation point.

IV. Reaction with Nitric Acid.

Myelopathic albumose is thrown down by strong nitric acid in the cold, and so far behaves like albumin. On raising the temperature to the boiling point, however, the coagulum dissolves more or less completely and reappears on cooling. This reaction, which is commonly accepted as one of the most characteristic reactions of the primary albumoses, is of great theoretical interest, and in my opinion justifies the provision inclusion of the body we are considering among the albumoses, notwithstanding the fact that it differs in some of its properties from the other members of the group. When the test is applied directly to the untreated urine the solution on boiling is generally only partial, and sometimes does not take place at all. If, however, the urine is first diluted, it almost invariably takes place to a very large extent. The same conditions which interfere with the solution of the heat coagulum already referred to seem also to interfere with the solution of the coagulum formed with nitric acid. The best way to apply the test is to add the acid drop by drop until

the maximum degree of precipitation is produced, which may be recognized by a tendency for the precipitate to dissolve where the acid collects in excess at the bottom of the test tube. The whole column of liquid is then boiled for a few seconds while stirred with a glass rod and is thrown on a pleated filter. A positive result is shown by the filtrate, first clear, becoming densely turbid on cooling, and it may be confirmed by finding that clearing again takes place on heating. This process may be repeated several times. If a negative result is obtained at first the test should be repeated with the urine diluted with three or four volumes of water or of normal urine. The coagulum with hydrochloric acid is likewise soluble on boiling.

V. Reaction with Potassium Ferrocyanide.

Myelopathic albumose is coagulated by potassium ferrocyanide and acetic acid, but in a manner somewhat different from albumin. With the latter complete coagulation takes place immediately, but with albumose it often takes time to develop, and unless a good deal of acid is used it may not appear at all. If this test is the one commonly used for the detection of proteids in general, as is the practice with some physicians, the presence of albumose may be altogether overlooked.

Trustworthiness and Delicacy of the Tests.

Since the year 1897, when I first became interested in the subject of myelopathic albumosuria, I have examined the urine in 9 cases of the disease; generally more than one sample has been submitted to analysis, and in some cases I have examined a large number. In every instance I have found the foregoing tests responded to in such a way as to leave no room for doubt that the body under investigation was myelopathic albumose, and in every case which has been watched to its termination the association with disease of the bony skeleton has been confirmed. Until recently, however, I had always found the albumose, if present at all, in enormous quantities, sometimes as much as 20 parts per mille, and I have often felt a doubt whether, if it were excreted in small amount—say as little as 0.5 per mille—the tests which I rely upon would be equally effectual for its recognition and differentiation from other proteids.

It is true that after free dilutions of the urine the same reactions were noted, but in the case of a substance of so doubtful a nature, and one whose behaviour seems so much influenced by environment, one felt that it was not safe to predicate what its behaviour would be under new conditions; that if it were excreted only in traces it might undergo some molecular or other changes which would interfere with its reactions to the various tests. In the latter part of 1904, however, my friend and former house-physician, Dr. Cyril Moss, who was then in the service of the P. and O. Steamship Company, was kind enough to bring me two bottles of urine from a woman who was in hospital in Sydney, N.S.W., with well-marked symptoms of multiple myeloma, and who has since been proved by autopsy to have suffered from that disease. This urine was high coloured, was amphoteric, and had a specific gravity of 1030. It gave the heat test for albumose distinctly. I acidulated it faintly with acetic acid, and heated it on a water bath, when a turbidity, moderate in amount, came on quite suddenly, just about 60° C. It disappeared on further heating, and reappeared on cooling. Similarly a turbidity was produced by dropping in nitric acid; it disappeared on heating and returned on cooling. A white band also appeared in contact with hydrochloric acid.

The turbidity was about the same intensity as would be yielded by a trace of albumin, or less than 0.5 part per mille.

Estimation by means of Esbach's picric acid albuminometer also indicated that the quantity of albumose was not more than 0.5 per 1,000. By comparison with gravimetric estimates of the quantity of albumose present in other cases, I have found that the readings of this instrument are approximately correct for this body. I conclude, therefore, that the tests I have advocated can be relied on to distinguish albumose from albumin in any case in which the former is present in sufficient amount to be capable of recognition at all.

I have stated that albumose is liable to undergo certain molecular changes which are difficult to explain and may puzzle the inexperienced. One of the most remarkable

and constant of these is that it becomes deposited in an insoluble form on standing for a longer or shorter time. This insoluble dysalbumose, as it has been called, is generally to be found in a specimen that has stood for some weeks.

I have by me a bottle of urine which I filtered quite clear twelve months ago, and which now contains a dense deposit. This deposit may take place very rapidly; in my first case it sometimes took place in the patient's bladder, and consequently the urine looked like milk when voided. This change is apparently of the same nature as the coagulation by heat.

Other Tests.

Before concluding I should like to allude to certain tests for myelopathic albumose which have been mentioned by various writers, but which have proved in my hands absolutely worthless. Of these the best known is the so-called biuret test, performed by treating the solution with caustic potash or soda and a trace of copper sulphate. The result is supposed to be that in the case of albumin a violet colour is produced; in the case of albumoses and peptones a rose red. Professor Huppert of Prague, however, states somewhat emphatically that the biuret reaction given by peptone is in no way distinguishable from that of albumin.¹ If this be true with regard to peptone, we may suspect that it will *a fortiori* be also true of albumoses, which chemically stand much closer to albumin. My own experience with the biuret test convinces me that it is of no use whatever as a means of distinguishing the different proteids as they occur in the urine. A rose pink is always obtained if the amount of copper is small, no matter what is the kind of proteid under examination. I have therefore no hesitation in saying that in practice the biuret reaction is absolutely worthless in the examination of urine.

Another test has been brought out by Boston of Philadelphia, and is described at length in his book on clinical diagnosis.² It depends on the fact that albumose contains loosely combined sulphur, which in the presence of alkali unites with lead salts to form black sulphide. Boston says that many chemists regard this reaction as one of the most characteristic features of the body we are considering, and that it distinguishes it from the digestive albumoses. I may remark in passing that in practice we are hardly likely to meet with difficulty in distinguishing myelopathic from digestive albumose; the real problem is to distinguish it from albumin. However, I have applied Boston's test to urine containing albumin, digestive albumose (Witte's peptone), and myelopathic albumose respectively, and I find that the results obtained in all three cases are quite indistinguishable.

Again, Dixon Mann states,³ on the authority of Grutterink and Graaf, that with an equal volume of 50 per cent. acetic acid the albumose will become solid, so that the test-tube can be inverted without spilling. This in my experience is certainly not the case. However, I maintain that the simple tests I have already described are sufficient to enable us to identify this body, and that from the point of view of the practising physician there is no practical advantage in adding to their number.

There are many points with regard to the nature of this interesting body which I have not touched upon in the foregoing remarks. If I were to deal with them I should have to enter into various chemical problems which are outside the scope of the present communication, the purpose of which is to place clearly before my readers the means by which the presence of myelopathic albumose in the urine may be determined.

REFERENCES.

¹ Neubauer and Vogel, *Analyse des Harns*; 3te Aufl., p. 478. ² Boston, *A Textbook of Clinical Diagnosis*, p. 187. ³ J. Dixon Mann, *Physiology and Pathology of the Urine*, p. 128.

FRENCH CONGRESS OF GYNAECOLOGY AND OBSTETRICS.—The fifth periodical Congress of Gynaecology, Obstetrics, and Paediatrics, will be held at Algiers, April 1st to 8th, 1907, under the general presidency of Professor Queirel, of Marseilles, who will also preside over the Section of Obstetrics. Dr. Boursier, of Bordeaux, will be President of the Gynaecological Section, and Dr. Guinon, of Paris, of that of Paediatrics. The General Secretary is Professor Jules Rouvier, 52, rue Daguerre, Algiers.

RHEUMATIC FEVER AND AMYLOID DEGENERATION.

By JAMES M. BEATTIE, M.A., M.D.,

LECTURER ON PATHOLOGICAL BACTERIOLOGY AND SENIOR ASSISTANT TO THE PROFESSOR OF PATHOLOGY, UNIVERSITY OF EDINBURGH.
[From the Pathological Laboratory, University of Edinburgh.]

THOUGH the causes of amyloid degeneration are not clearly understood, still it is generally recognized that in the great majority, if not all cases, the condition is secondary to some infective agent. This infective agent is generally bacterial, and has been acting for a considerable length of time. The experimental work of Krakow, Maximow, Davidsohn and others has all gone to support the bacterial agent as the cause, and if we exclude the experiments of Czerny, which have not been confirmed by others who carried out similar work, the degeneration has been produced artificially only by bacteria or their products. It is true that cases of amyloid degeneration have been recorded as following cancer, leukaemia, malaria, and lymphadenoma; one of these is due to a parasite which can produce toxic products, very much allied to the products of bacteria, and, while the cause of the other three diseases is still unknown, I do not think I am stating too much in saying that the majority of pathologists favour the parasitic origin of at least two of them. In view of these facts, it seemed to me that in investigating and upholding the bacterial origin of acute rheumatism some help might be obtained by a study of the more chronic lesions which all recognized as sequelae of acute rheumatism, and the possibility of their experimental production by the streptococcus which has been claimed as causal.

At the commencement of my observations my attention was directed to the presence of amyloid degeneration in various organs of a case of cardiac disease following acute rheumatism. Since that time three other cases have been noted by me.

CASE I.

A. P., female, aged 28, was admitted to the Royal Infirmary, Edinburgh, on November 9th, 1904, complaining of breathlessness.

History.—Seven months previously she had arthritis in the right ankle and was treated for acute rheumatism. This was followed a month later by arthritis in the knees and ankles, and in October, a month before admission, she had pains over the heart and she became breathless. There was no history of any previous attack of acute rheumatism, and till June, 1904, her health had been perfectly satisfactory.

Condition on Admission.—There was a well-marked systolic murmur over the mitral area and other signs of definite cardiac disease.

After-history.—November 13th. There was a purpuric rash over the chest and upper part of the abdomen. On the flexor surface of the left forearm, about $\frac{1}{2}$ in. below the bend of the elbow, there was an egg-shaped swelling $1\frac{1}{2}$ in. long by $\frac{1}{2}$ in. wide. This was more definite on the radial than on the ulnar aspect. The left radial pulse at the wrist was hardly perceptible. The liver was slightly enlarged and tender on palpation. On November 16th there was marked diarrhoea. The diarrhoea continued. Tenderness over the liver increased, and very acute tenderness developed over the right lower ribs, especially the fifth and sixth anteriorly, and in the axilla, and continued till the death of the patient on December 4th, at 6.10 a.m.

Autopsy (December 4th, 11 a.m.).

Heart and Pericardium.—There was 1 oz. of clear serous fluid in the pericardial sac, no adhesions, and no recent pericarditis. The endocardium showed a few thickened patches of chronic endocarditis. Over the right ventricle there was an excess of fat. The right auricle was filled with recent blood clot and was slightly dilated. The right ventricle was also slightly dilated. The tricuspid and pulmonary valve segments showed no special change. The left auricle was dilated and filled with recent blood clot. The left ventricle was hypertrophied and slightly dilated. The aortic valve segments were slightly thickened, especially at their attached margin, but there was no evidence of any recent aortic endocarditis. On the inferior surface of the anterior segment of the mitral valve there was a large mass of fairly firm vegetations, in the form of irregular nodules. The largest of these was $\frac{1}{2}$ in. in diameter. On the auricular surface of the posterior segment there were some small recent warty vegetations, which spread upwards for $\frac{1}{2}$ in. over the wall of the auricle. There was neither ulceration nor rupture of any of the segments. There was chronic thickening of all the mitral segments; but no stenosis apart from that caused by the vegetations. From the vegetations a culture corresponding in all characters with the *Streptococcus rheumaticus* was isolated.

Lungs.—There was slight emphysema with collapse at the